

DOI: 10.4081/aiol.2015.5470

SUPPLEMENTARY MATERIAL

Spatial distribution of microphytobenthos, meiofauna and macrofauna in the north-western Adriatic Sea: a synoptic study

Annalisa Franzo, Tamara Cibic, Paola Del Negro, Cinzia De Vittor

Sezione Oceanografia, Istituto Nazionale di Oceanografia e Geofisica Sperimentale (OGS), via A. Piccard 54, I-34151 Trieste, Italy

*Corresponding author: afranzo@ogs.trieste.it

Non commercial use only

Supplementary Tab. 1. Floristic lists of diatoms observed in the study area and during the two sampling periods. The corresponding living forms and abundance range are indicated. A distinction was made considering the dimensions of different species belonging to the genus *Navicula*: spp. 1 <20 µm; 20 µm < spp. 2 <40 µm; spp. 3 >40 µm.

September 2010			Range min-max ABU (cells cm ⁻³)
<i>Achnanthes</i> spp.	Bory de Saint-Vincent, 1822	Benthic, epiphytic or epipsammic	0-300
<i>Amphora</i> spp.	Ehrenberg ex Kützing, 1844	Benthic, mostly epipsammic	0-300
<i>Bacillaria paxillifera</i>	(O.F.Müller) T.Marsson, 1901	Tychopelagic	0-6300
<i>Caloneis</i> cfr. <i>leptosoma</i>	(Grunow) Krammer, 1985	Benthic, epipelagic	0-2700
<i>Caloneis liber</i>	(W.Smith) Cleve, 1894	Benthic, epipelagic	0-1200
<i>Diploneis bombus</i>	(Ehrenberg) Ehrenberg, 1853	Benthic, epipelagic	0-1200
<i>Ellobium</i> spp.	Röding, 1798	Benthic, epiphytic	0-300
<i>Entomoneis</i> spp.	Ehrenberg, 1845	Benthic, epipelagic	0-1500
<i>Gyrosigma acuminatum</i>	(Kützing) Rabenhorst, 1853	Benthic, epipelagic	0-2100
<i>Gyrosigma attenuatum</i>	(Kützing) Rabenhorst, 1853	Benthic, epipelagic	0-1200
<i>Gyrosigma balticum</i>	(Ehrenberg) Rabenhorst, 1853	Benthic, epipelagic	0-600
<i>Gyrosigma fasciola</i>	(Ehrenberg) J.W.Griffith & Henfrey, 1856	Benthic, epipelagic	0-300
<i>Gyrosigma macrum</i>	(W.Smith) J.W.Griffith & Henfrey, 1856	Benthic, epipelagic	0-300
<i>Gyrosigma</i> spp.	(Kützing) Rabenhorst, 1853	Benthic, epipelagic	0-3300
<i>Navicula directa</i>	Hassall, 1845	Benthic, epipelagic	0-1500
<i>Navicula mollis</i>	(W.Smith) Ralfs, 1861	Benthic, epipelagic	0-2700
<i>Navicula</i> spp. 1	(W.Smith) Cleve, 1896	Epiphytic	0-4500
<i>Navicula</i> spp. 2	Bory de Saint-Vincent, 1822	Benthic	0-3900
<i>Navicula</i> spp. 3	Bory de Saint-Vincent, 1822	Benthic	0-7800
<i>Nitzschia fasciculata</i>	Bory de Saint-Vincent, 1822	Benthic	0-900
<i>Nitzschia longissima</i>	(Grunow) Grunow, 1881	Benthic, epipelagic	0-10200
<i>Nitzschia lorenziana</i>	(Brébisson) Ralfs, 1861	Benthic, epipelagic	0-300
<i>Nitzschia lorenziana</i>	Grunow, 1879	Benthic, epipelagic	0-3900
<i>Nitzschia lorenziana</i> var. <i>densestriata</i>	(Persoon) A. Schmidt et al., 1874	Benthic, epipelagic	0-300
<i>Nitzschia sigma</i>	(Kützing) W.Smith, 1853	Benthic, epipelagic	0-3000

September 2010			Range min-max ABU (cells cm ⁻³)
<i>Nitzschia sigma</i> var. <i>intercedens</i>	Grunow, 1878	Benthic, epipellic	0-300
<i>Nitzschia sigma</i> var. <i>rigida</i>	Grunow ex Van Heurck, 1880	Benthic, epipellic	0-2700
<i>Nitzschia sigma</i> var. <i>sigmatella</i>	Grunow, 1878	Benthic, epipellic	0-300
<i>Nitzschia tryblionella</i>	Hantzsch, 1860	Benthic, epipellic	0-600
<i>Nitzschia</i> spp.	Hassall, 1845	Benthic	0-7200
<i>Paralia sulcata</i>	(Ehrenberg) Cleve, 1873	Tychopelagic	0-33600
<i>Pinnularia</i> spp.	Ehrenberg, 1843	Benthic, epipellic	0-3600
<i>Pleurosigma aestuarii</i>	(Brébisson ex Kützing) W.Smith, 1853	Benthic, epipellic	0-2700
<i>Pleurosigma elongatum</i>	W.Smith, 1852	Benthic, epipellic	0-300
<i>Pleurosigma</i> spp.	W.Smith, 1852	Benthic, epipellic	0-1800
<i>Psammodictyon panduriforme</i>	(W.Gregory) D.G.Mann, 1990	Benthic	0-300
<i>Rhopalodia gibba</i>	(Ehrenberg) Otto Müller, 1895	Benthic	0-10800
<i>Surirella minuta</i>	Brébisson, 1849	Benthic	0-300
<i>Surirella</i> spp.	Turpin, 1828	Benthic	0-300
<i>Toxarium hennedyanum</i>	(Gregory) Pelletan, 1889	Benthic, epiphytic	0-300
<i>Tryblionella punctata</i>	W. Smith, 1853	Benthic, epipellic	0-300

March 2011			Range min-max ABU (cells cm ⁻³)
<i>Amphora ostrearia</i>	Brébisson, 1849	Benthic, epipsammic	0-300
cfr. <i>Biddulphia rhombus</i>	(Ehrenberg) W.Smith, 1854	Planktonic	0-300
<i>Caloneis liber</i>	(W.Smith) Cleve, 1894	Benthic, epipellic	0-300
<i>Ceratoneis closterium</i>	Ehrenberg, 1839	Tychopelagic	0-900
cfr. <i>Cymbella</i> spp.	C.Agardh, 1830	Benthic, epipellic	0-300
<i>Diploneis bombus</i>	(Ehrenberg) Ehrenberg, 1853	Benthic, epipellic	0-2100
<i>Ellobiums</i> spp.	Röding, 1798	Benthic, epiphytic	0-300
<i>Entomoneis alata</i>	(Ehrenberg) Ehrenberg, 1845	Benthic, epipellic	0-900
<i>Gyrosigma acuminatum</i>	(Kützing) Rabenhorst, 1853	Benthic, epipellic	0-1200
<i>Gyrosigma balticum</i>	(Ehrenberg) Rabenhorst, 1853	Benthic, epipellic	0-300
<i>Gyrosigma macrum</i>	(W.Smith) J.W.Griffith & Henfrey, 1856	Benthic, epipellic	0-300
<i>Gyrosigma spencerii</i>	(Kützing) Rabenhorst, 1853	Benthic, epipellic	0-900
<i>Gyrosigma</i> spp.	Hassall, 1845	Benthic, epipellic	0-300
<i>Navicula directa</i>	(W.Smith) Ralfs, 1861	Benthic, epipellic	0-300
<i>Navicula</i> spp. 1	Bory de Saint-Vincent, 1822	Benthic	0-3900
<i>Navicula</i> spp. 2	Bory de Saint-Vincent, 1822	Benthic	0-3600
<i>Navicula</i> spp. 3	Bory de Saint-Vincent, 1822	Benthic	0-300
<i>Nitzschia fasciculata</i>	(Grunow) Grunow, 1881	Benthic, epipellic	0-300
<i>Nitzschia lorenziana</i>	Grunow, 1879	Benthic, epipellic	0-300
<i>Tryblionella punctata</i>	W.Smith, 1853	Benthic, epipellic	0-300
<i>Nitzschia sigma</i> var. <i>sigmatella</i>	Grunow, 1878	Benthic, epipellic	0-900
<i>Nitzschia tryblionella</i>	Hantzsch, 1860	Benthic, epipellic	0-600
<i>Nitzschia</i> spp.	Baer, 1826	Benthic	0-300
<i>Paralia sulcata</i>	(Ehrenberg) Cleve, 1873	Tychopelagic	0-39300
<i>Pinnularia</i> spp.	Ehrenberg, 1843	Benthic, epipellic	0-1200
<i>Pleurosigma normanii</i>	Ralfs, 1861	Benthic, epipellic	0-300

March 2011			Range min-max ABU (cells cm ⁻³)
<i>Rhopalodia gibba</i>	(Ehrenberg) Otto Müller, 1895	Benthic	0-1500
<i>Skeletonema costatum</i>	(Greville) Cleve, 1873	Planktonic	0-99600
<i>Thalassionema nitzschioides</i>	(Grunow) Mereschkowsky, 1902	Planktonic	0-1200
<i>Thalassionema</i> spp.	Grunow ex Mereschkowsky, 1902	Planktonic	0-900

Non commercial use only

Supplementary Tab. 2. Complete list of macrofauna taxa observed in the present study and during both sampling periods. The abundance range is indicated.

September 2010 Taxa		Range min-max	ABU (ind. m ⁻²)
<i>Abra alba</i>	(W. Wood, 1802)		0-30
<i>Abra prismatica</i>	(Montagu, 1808)		0-70
<i>Abra</i> spp.	Lamarck, 1818		0-30
<i>Acanthocardia paucicostata</i>	(G. B. Sowerby II, 1834)		0-10
<i>Acteon tornatilis</i>	(Linnaeus, 1758)		0-10
Actiniaria indet.			0-10
<i>Alitta succinea</i>	(Leuckart, 1847)		0-10
<i>Amage adspersa</i>	(Grube, 1863)		0-50
<i>Ampelisca spinipes</i>	Boeck, 1861		0-1270
<i>Ampelisca</i> spp.	Krøyer, 1842		0-320
<i>Amphictene auricoma</i>	(O.F. Müller, 1776)		0-320
Amphipodaindet.	Latreille, 1816		0-10
<i>Amphiura chiajei</i>	Forbes, 1843		0-20
<i>Amphiura filiformis</i>	(O.F. Müller, 1776)		0-20
<i>Ampithoe ramondi</i>	Audouin, 1826		0-210
<i>Anadara transversa</i>	(Say, 1822)		0-250
<i>Anapagurus</i> spp.	Henderson, 1886		0-20
<i>Antalis dentalis</i>	(Linnaeus, 1758)		0-10
Aphroditidae indet.	Malmgren, 1867		0-10
<i>Aponuphis bilineata</i>	(Baird, 1870)		0-10
<i>Aporrhais pespelecani</i>	(Linnaeus, 1758)		0-10
<i>Apseudopsis latreillii</i>	(Milne Edwards, 1828)		0-950
<i>Arenicola marina</i>	(Linnaeus, 1758)		0-20
<i>Arenicola</i> spp.	Lamarck, 1801		0-20
<i>Aspidosiphon (Aspidosiphon) muelleri muelleri</i>	Diesing, 1851		0-10

September 2010		
Taxa		Range min-max ABU (ind. m ⁻²)
<i>Brada villosa</i>	(Rathke, 1843)	0-20
<i>Callianassa</i> spp.	Leach, 1814	0-10
<i>Calyptraea chinensis</i>	(Linnaeus, 1758)	0-30
<i>Capitella giardi</i>	(Mesnil, 1897)	0-20
<i>Caprella</i> cfr. <i>andreae</i>	Mayer, 1890	0-10
<i>Caprella</i> spp.	Lamarck, 1801	0-10
Cirratulidae indet.	Carus, 1863	0-10
<i>Cirratulus</i> spp.	Lamarck, 1818	0-80
<i>Condylactis aurantiaca</i>	(Delle Chiaje, 1825)	0-10
<i>Corbula gibba</i>	(Olivi, 1792)	0-2380
<i>Cossura soyeri</i>	Laubier, 1964	0-20
<i>Cylichna cylindracea</i>	(Pennant, 1777)	0-20
<i>Dischides politus</i>	(S. Wood, 1842)	0-10
<i>Echinocardium cordatum</i>	(Pennant, 1777)	0-10
<i>Elasmopus rapax</i>	Costa, 1853	0-10
<i>Epitonium</i> spp. juv.	Röding, 1798	0-10
<i>Euclymene palermitana</i>	(Grube, 1840)	0-40
<i>Eunice vittata</i>	(Delle Chiaje, 1828)	0-10
<i>Euspira guilleminii</i>	(Payraudeau, 1826)	0-10
Flabelligeridae	de Saint-Joseph, 1894	0-10
<i>Glycera alba</i>	(O.F. Müller, 1776)	0-30
<i>Glycera fallax</i>	Quatrefages, 1850	0-10
<i>Glycera</i> spp.	Lamarck, 1818	0-10
<i>Glycera tridactyla</i>	Schmarda, 1861	0-20
<i>Golfingia</i> spp.	Lankester, 1885	0-20
<i>Goneplax rhomboides</i>	(Linnaeus, 1758)	0-10
<i>Harmothoe</i> spp.	Kinberg, 1856	0-10
<i>Heteromastus filiformis</i>	(Claparède, 1864)	0-40

September 2010		
Taxa		Range min-max ABU (ind. m ⁻²)
<i>Hydrobia acuta</i>	(Draparnaud, 1805)	0-220
<i>Idotea</i> spp.	Fabricius, 1798	0-20
<i>Inermonephtys inermis</i>	(Ehlers, 1887)	0-10
<i>Iphinoe serrata</i>	Norman, 1867	0-10
<i>Kurtiella bidentata</i>	(Montagu, 1803)	0-10
<i>Labioleanira yhleni</i>	(Malmgren, 1867)	0-10
<i>Leptopentacta elongata</i>	(Düben & Koren, 1846)	0-10
<i>Leucothoe venetiarum</i>	Giordani- Soika, 1950	0-90
<i>Liocarcinus corrugatus</i>	(Pennant, 1777)	0-10
<i>Liocarcinus depurator</i>	(Linnaeus, 1758)	0-10
<i>Lumbrineriopsis paradoxa</i>	(Saint-Joseph, 1888)	0-20
<i>Lumbrineris adriatica</i>	Fauvel, 1940	0-170
Maldanidae indet.	Malmgren, 1867	0-50
<i>Mangelia costulata</i>	Risso, 1826	0-10
<i>Mimachlamys varia</i>	(Linnaeus, 1758)	0-20
<i>Moerella distorta</i>	(Poli, 1791)	0-40
<i>Monia patelliformis</i>	(Linnaeus, 1761)	0-10
<i>Musculus subpictus</i>	(Cantraine, 1835)	0-10
Mysida indet.	Boas, 1883	0-10
<i>Mysta picta</i>	(Quatrefages, 1866)	0-10
<i>Namalycastis brevicornis</i>	(Audouin & Milne Edwards, 1833)	0-10
<i>Nassarius reticulatus</i>	(Linnaeus, 1758)	0-10
<i>Nassarius</i> spp.	Duméril, 1805	0-20
Natantia indet.	Boas, 1880	0-10
<i>Neanthes kerguelensis</i>	(McIntosh, 1885)	0-10
<i>Necallianassa acanthura</i>	(Caroli, 1946)	0-10
<i>Nephtys hombergii</i>	Savigny in Lamarck, 1818	0-40
<i>Nephtys</i> spp.	Cuvier, 1817	0-20

September 2010		
Taxa		Range min-max ABU (ind. m ⁻²)
<i>Nereis lamellosa</i>	Ehlers, 1864	0-20
<i>Nereis rava</i>	Ehlers, 1864	0-10
<i>Notomastus</i> spp.	Sars, 1850	0-10
<i>Nucula sulcata</i>	Bronn, 1831	0-190
<i>Nuculana pella</i>	(Linnaeus, 1767)	0-20
<i>Oestergrenia thomsonii</i>	(Herapath, 1865)	0-50
<i>Ophiura</i> spp.	Lamarck, 1801	0-20
Orbiniidae indet.	Hartman, 1942	0-10
<i>Ova canaliferus</i>	(Lamarck, 1816)	0-10
<i>Paradoneis lyra</i>	(Southern, 1914)	0-20
<i>Phaxas adriaticus</i>	(Coen, 1933)	0-40
<i>Pherusa monilifera</i>	(Delle Chiaje, 1841)	0-10
<i>Phoronis muelleri</i>	Selys-Lonchamps, 1903	0-10
<i>Phyllodoce</i> spp.	Lamarck, 1818	0-10
<i>Pitar rudis</i>	(Poli, 1795)	0-20
<i>Polititapes rhomboides</i>	(Pennant, 1777)	0-30
Priapulida indet.	Théel, 1906	0-20
<i>Processa</i> spp.	Leach, 1815 [in Leach, 1815-1875]	0-10
<i>Pseudoleiocapitella fauveli</i>	Harmelin, 1964	0-80
<i>Pseudostichopus peripatus</i>	(Sluiter, 1901)	0-20
<i>Scoletoma fragilis</i>	(O.F. Müller, 1776)	0-20
Sigalionidae indet.	Malmgren, 1867	0-20
<i>Spisula subtruncata</i>	(da Costa, 1778)	0-10
<i>Sternaspis scutata</i>	Ranzani, 1817	0-140
Syllidae indet.	Grube, 1850	0-10
<i>Tellimya ferruginosa</i>	(Montagu, 1808)	0-430
<i>Tellina pulchella</i>	Lamarck, 1818	0-50
<i>Tellina</i> spp.	Linnaeus, 1758	0-10

September 2010		
Taxa		Range min-max ABU (ind. m ⁻²)
Tellinidae indet.	Blainville, 1814	0-30
Terebellidae indet.	Johnston, 1846	0-40
<i>Tharyx</i> spp.	Webster & Benedict, 1887	0-180
<i>Timoclea ovata</i>	(Pennant, 1777)	0-10
<i>Tisbe</i> spp.	Lilljeborg, 1853	0-10
<i>Turritella communis</i>	Risso, 1826	0-240
<i>Turritella triplicata</i>	(Brocchi, 1814)	0-20

Non commercial use only

March 2011		
Taxa		Range min-max ABU (ind. m ⁻²)
<i>Abra alba</i>	(W. Wood, 1802)	0-10
<i>Abra prismatica</i>	(Montagu, 1808)	0-10
<i>Acanthocardia</i> spp.	J.E. Gray, 1851	0-10
Actiniaria indet.		0-50
<i>Alpheus glaber</i>	(Olivi, 1792)	0-10
<i>Amathia semiconvoluta</i>	Lamouroux, 1824	0-10
<i>Ampelisca spinipes</i>	Boeck, 1861	0-310
<i>Ampelisca</i> spp.	Krøyer, 1842	0-110
<i>Amphictene auricoma</i>	(O.F. Müller, 1776)	0-10
<i>Amphipholis squamata</i>	(Delle Chiaje, 1828)	0-20
<i>Amphiura chiajei</i>	Forbes, 1843	0-40
<i>Anadara transversa</i>	(Say, 1822)	0-90
<i>Anapagurus</i> spp.	Henderson, 1886	0-20
Aphroditidae indet.	Malmgren, 1867	0-40
<i>Aponuphis bilineata</i>	(Baird, 1870)	0-30
<i>Aporrhais pespelecani</i>	(Linnaeus, 1758)	0-40
<i>Apseudopsis latreillii</i>	(Milne Edwards, 1828)	0-60
<i>Arenicola</i> spp.	Lamarck, 1801	0-20
<i>Brada villosa</i>	(Rathke, 1843)	0-10
<i>Brissopsis lyrifera</i>	(Forbes, 1841)	0-10
<i>Callianassa</i> spp.	Leach, 1814	0-10
Capitellidae indet.	Grube, 1862	0-40
<i>Caprella</i> spp.	Lamarck, 1801	0-80
<i>Capulus ungaricus</i>	(Linnaeus, 1758)	0-10
<i>Caulleriella</i> spp.	Chamberlin, 1919	0-140
<i>Cereus pedunculatus</i>	(Pennant, 1777)	0-10
Cirratulidae indet.	Carus, 1863	0-50

March 2011		
Taxa		Range min-max ABU (ind. m ⁻²)
<i>Cirratulus</i> spp.	Lamarck, 1818	0-20
<i>Clymenura</i> spp.	Verrill, 1900	0-40
<i>Condylactis aurantiaca</i>	(Delle Chiaje, 1825)	0-10
<i>Corbula gibba</i>	(Olivi, 1792)	0-450
<i>Cylichna cylindracea</i>	(Pennant, 1777)	0-50
Decapoda indet. Larva	Latreille, 1803	0-10
<i>Dischides politus</i>	(S. Wood, 1842)	0-10
<i>Dosinia lupinus</i>	(Linnaeus, 1758)	0-10
<i>Epitonium</i> spp. Juv.	Röding, 1798	0-10
<i>Euclymene palermitana</i>	(Grube, 1840)	0-40
<i>Eulima glabra</i>	(da Costa, 1778)	0-10
<i>Eunoe nodosa</i>	(M. Sars, 1861)	0-10
<i>Euspira guilleminii</i>	(Payraudeau, 1826)	0-20
<i>Galathea intermedia</i>	Lilljeborg, 1851	0-10
<i>Glycera alba</i>	(O.F. Müller, 1776)	0-20
<i>Glycera</i> spp.	Lamarck, 1818	0-10
<i>Glycera tridactyla</i>	Schmarda, 1861	0-30
<i>Glycera unicornis</i>	Lamarck, 1818	0-10
<i>Golfingia</i> spp.	Lankester, 1885	0-30
<i>Goneplax rhomboides</i>	(Linnaeus, 1758)	0-10
<i>Heteromastus filiformis</i>	(Claparède, 1864)	0-30
<i>Hydrobia acuta</i>	(Draparnaud, 1805)	0-140
Hydrozoa indet.	Owen, 1843	0-10
<i>Iphinoe serrata</i>	Norman, 1867	0-10
<i>Labioleanira yhleni</i>	(Malmgren, 1867)	0-10
<i>Leptopentacta elongata</i>	(Düben & Koren, 1846)	0-10
<i>Leucothoe venetiarum</i>	Giordani- Soika, 1950	0-10

March 2011		
Taxa		Range min-max ABU (ind. m ⁻²)
<i>Liocarcinus pusillus</i>	(Leach, 1816)	0-10
<i>Lumbriclymene minor</i>	Arwidsson, 1906	0-10
<i>Lumbrineris adriatica</i>	Fauvel, 1940	0-110
<i>Lumbrineris</i> spp.	Blainville, 1828	0-20
Maldanidae indet.	Malmgren, 1867	0-110
<i>Mangelia costulata</i>	Risso, 1826	0-10
<i>Melinna palmata</i>	Grube, 1870	0-90
Mysida indet.	Boas, 1883	0-10
<i>Nassarius pygmaeus</i>	(Lamarck, 1822)	0-90
<i>Nassarius reticulatus</i>	(Linnaeus, 1758)	0-50
Natantia indet.	Boas, 1880	0-10
<i>Neanthes kerguelensis</i>	(McIntosh, 1885)	0-30
<i>Necallianassa acanthura</i>	(Caroli, 1946)	0-30
Nemertea indet.		0-10
<i>Nephtys hombergii</i>	Savigny in Lamarck, 1818	0-20
<i>Nephtys</i> spp.	Cuvier, 1817	0-50
<i>Notomastus</i> spp.	Sars, 1850	0-60
<i>Nucula</i> spp.	Lamarck, 1799	0-100
<i>Nucula sulcata</i>	Bronn, 1831	0-130
<i>Oestergrenia thomsonii</i>	(Herapath, 1865)	0-70
<i>Othomaera schmidtii</i>	(Stephensen, 1915)	0-10
<i>Ova canaliferus</i>	(Lamarck, 1816)	0-10
<i>Owenia fusiformis</i>	Delle Chiaje, 1844	0-20
<i>Paradoneis lyra</i>	(Southern, 1914)	0-10
<i>Pherusa monilifera</i>	(Delle Chiaje, 1841)	0-10
<i>Polititapes rhomboides</i>	(Pennant, 1777)	0-10
Polychaetae indet.		0-20

March 2011		
Taxa		Range min-max ABU (ind. m ⁻²)
Polynoinae indet.	Kinberg, 1856	0-90
Priapulida indet.	Théel, 1906	0-20
<i>Priapulus caudatus</i>	Lamarck, 1816	0-10
<i>Processa</i> spp.	Leach, 1815 [in Leach, 1815-1875]	0-10
Processidae indet.	Ortmann, 1896	0-10
<i>Pseudoleiocapitella fauveli</i>	Harmelin, 1964	0-10
<i>Raphitoma linearis</i>	(Montagu, 1803)	0-10
<i>Saccella commutata</i>	(Philippi, 1844)	0-10
Sipunculidae indet.	Rafinesque, 1814	0-10
<i>Sipunculus (Sipunculus) nudus</i>	Linnaeus, 1766	0-10
Spatangoida indet.	L. Agassiz, 1840	0-10
Spionidae indet.	Grube, 1850	0-10
<i>Sternaspis scutata</i>	Ranzani, 1817	0-160
<i>Streblospio shrubsolii</i>	(Buchanan, 1890)	0-20
Syllidae indet.	Grube, 1850	0-20
<i>Tellimya ferruginosa</i>	(Montagu, 1808)	0-450
<i>Tellina pulchella</i>	Lamarck, 1818	0-20
Terebellidae indet.	Johnston, 1846	0-30
<i>Tharyx</i> spp.	Webster & Benedict, 1887	0-20
<i>Timoclea ovata</i>	(Pennant, 1777)	0-50
<i>Tisbe</i> spp.	Lilljeborg, 1853	0-10
<i>Turritella cingulata</i>	G. B. Sowerby I, 1825	0-10
<i>Turritella communis</i>	Risso, 1826	0-320
<i>Turritella triplicata</i>	(Brocchi, 1814)	0-10