



pISSN: 2037-7452

eISSN: 2037-7460

<https://www.pagepressjournals.org/index.php/bam/index>

Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. The Early Access service lets users access peer-reviewed articles well before print / regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

The European Journal of Translational Myology is, therefore, e-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one.


The final version of the manuscript will then appear on a regular issue of the journal.

E-publishing of this PDF file has been approved by the authors.

Eur J Transl Myol 2024 [Online ahead of print]

To cite this Article:

Fallah A, Dabbaghian FH, Jamshidi A, Akhtari E. ***Placebo-controlled effect of topical Qust (Costus) oil on postmenopausal women's sexual desire disorder: a double-blind, randomized clinical trial.*** Eur J Transl Myol doi: 10.4081/ejtm.2024.12660

 ©The Author(s), 2024
Licensee PAGEPress, Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.



Placebo-controlled effect of topical Qust (Costus) oil on postmenopausal women's sexual desire disorder: a double-blind, randomized clinical trial

Akram Fallah,¹ Fattaneh Hashem Dabbaghian,² Amirhossein Jamshidi,^{3,4} Elham Akhtari⁵

¹Traditional Medicine Fellow, Faculty of Traditional Medicine, Iran University of Medical Sciences, Tehran, Iran; ²Community Medicine Specialist, Department of Traditional Medicine, Institute for Studies in Medical History, Persian and Complementary Medicine, School of Persian Medicine, Iran University of Medical Sciences, Tehran, Iran; ³Institute for Studies in Medical History, Persian and Complementary Medicine, Iran University of Medical Sciences, Tehran, Iran; ⁴Department of Traditional Pharmacy, School of Persian Medicine, Iran University of Medical Sciences, Tehran, Iran; ⁵Department of Medicine, Institute for Studies in Medical History, Persian and Complementary Medicine, School of Persian Medicine, Iran University of Medical Sciences, Tehran, Iran

Abstract

Decreased libido and anorgasmia are common problems for women after menopause that reduce the quality of life of couples. This study examined the effect of topical Qust oil on sexual desire disorder in postmenopausal women. In this double-blind, randomized, clinical trial, 110 postmenopausal women with decreased sexual desire visiting a Traditional Medicine Center and Hazrat Rasool Akram Hospital (affiliated to Iran University of Medical Sciences) were selected by convenience consecutive sampling and randomly assigned to experimental and control groups. The experimental group received qust oil, while the control group was given a placebo (liquid paraffin); they were instructed to massage the product topically on their pubic area and perineum daily. The sexual function of both groups was assessed and compared before the intervention and four weeks after the intervention using the Female Sexual Function Index. The mean and standard deviation of the improvement of sexual function post-intervention were $37.66 \pm 32.52\%$ and $11.96 \pm 11.18\%$ in the experimental and control groups, respectively ($p < 0.001$). In terms of the improvement of components of sexual function, a significant difference was observed between the two groups in the subscales of sexual desire [$57.05 \pm 42.99\%$ vs. $21.25 \pm 27.85\%$, $p < 0.001$], arousal, orgasm, and satisfaction ($p < 0.001$ for all); however, no significant difference was observed in terms of

lubrication ($p=0.25$) and pain during intercourse ($p=0.776$). In postmenopausal women with sexual dysfunction, massaging the pubic area and perineum with qust oil for at least four weeks significantly improves desire, arousal, orgasm, and sexual satisfaction.

Key words: sexual desire, menopause, costus oil, Persian medicine.

Most women spend more than a third of their life with menopause, and the population of postmenopausal women in the world will increase to 1,000,300,000 by 2030¹. Hot flashes, circadian rhythm disorders, mood disorders such as decreased concentration, depression and anxiety, tachycardia, headache, musculoskeletal pain, and sexual disorders are common problems during menopause². The onset of menopause is associated with changes in sex hormones, including a decrease in estrogen, increase in FSH (Follicle-Stimulating Hormone), decrease in LH (Luteinizing Hormone) and consequently a decline in progesterone and cessation of menstruation³. Despite the numerous advances in the prevention and treatment of menopause complications, decreased libido and anorgasmia are still among the most common problems experienced by postmenopausal women that can decrease their quality of life⁴. Sexual functioning is an important component of quality of life⁵. The high prevalence of sexual dysfunction, its undeniable impact on marital disputes and the ensuing social problems, and the patients' refusal to discuss their sexual problems out of shame highlight the need to delve deeper into this subject⁶.

The prevalence of sexual dysfunction is reported to be about 72% in postmenopausal women in Iran, with 75% affecting the women's arousal and 62% their desire⁷. Sexual dysfunction leading to divorce has been reported to be about 40% in Iran. Meanwhile, lack of sexual satisfaction can lead to problems such as depression and deterioration of the family's mental health⁸. Sexual responses and desires are multifactorial, influenced by psychological, social, environmental, and learned factors. Sexual dysfunction is often accompanied by other psychological disorders such as anxiety, depression, personality disorder, and schizophrenia⁴. Currently, two modalities, namely sex therapy and drug therapies, are used to treat female sexual dysfunction⁹. Drug therapies include estrogen and testosterone compounds and anti-anxiety and anti-depressant medications, which have side effects and may interact with other drugs taken by the patient during menopause¹⁰. The American Endocrine Society has recommended against excessive prescription of androgens for treating female sexual

problems, citing side effects such as increased risk of breast cancer and cardiovascular diseases¹¹. In traditional medicine references, the reasons for poor sexual desire include weaknesses of the main body organs (heart, brain, and liver)¹². The corroboration of the comorbidity in the increased prevalence of sexual dysfunction and psychological problems based on new statistical studies is somewhat aligned with this view held by traditional medicine¹³. The treatments mentioned in accessible sources of traditional medicine include a wide range of commonly-used substances, from libido-enhancing single ingredients to compounds containing numerous elements.

Despite the increasing availability of effective conventional medical treatments, herbal remedies continue to be a popular alternative for men and women looking to improve their sexual functioning. Nevertheless, the efficacy of most herbal agents in treating sexual problems remains unclear¹⁴. Out of the proposed treatments, this study examines a simple and accessible drug that can be used topically to avoid interfering with the drugs used by postmenopausal women, who are often middle-aged and might be taking multiple drugs to manage other chronic diseases. Oil massage has been mentioned in many references of traditional medicine as a mechanism for stimulating sexual desire¹⁵. Qust (Costus) oil is derived from Iranian books of traditional medicine. It contains bitter qust root, Chinese cinnamon, marjoram, valerian root, cloves, nutmeg fruit, *Acorus calmus* root, *Ferula gummosa* gum resin, olive oil, grape vinegar, and water¹⁵. According to traditional medicine, this product is *muhatal* and *musakhan* and a tonic for the nerves, organs, and sexual desire¹⁵. When topically used, simple qust oil, which is the main part of this product, is a tonic for the nerves, organs, and sexual desire, a strong *muhatal* and reliever of melancholic and phlegmatic fever and chills¹⁵. Although it is mostly used to strengthen the nerves and organs, it was chosen in this study due to the effects of its constituent plants on sexual desire.

Materials and Methods

This was a double-blind, randomized, clinical trial on postmenopausal women aged 40-65 years with sexual dysfunction visiting a traditional medicine center and Hazrat Rasool Akram Hospital, affiliated to Iran University of Medical Sciences.

Sample size was calculated based on type-I error of 0.05 and study power of 80%. Since there was no information about the effect of this drug, and assuming the equality of the standard deviation (SD) of changes in sexual desire in the two groups, the effect size index and Cohen's formula were used to determine the sample size, and the effect size was taken as 0.6.

$$n = \frac{2 \left(\frac{Z_{\alpha} + Z_{\beta}}{2} \right)^2}{d^2} d = \frac{\mu_1 - \mu_2}{\sigma}$$

The sample size was thus estimated as 44 per group. After adding 20% attrition, the required sample size was increased to 55.

Inclusion criteria

Menopause (at least 1 year since the last menstruation), age 40 to 65 years, complaining of decreased sexual desire, having an emotionally satisfying relationship with a stable and sexually active husband within at least the last 6 months, normal pelvic examination, and providing informed consent for participation in the study.

Exclusion criteria

Vaginal infection, vaginal bleeding, a serious physical medical condition or chronic illness requiring constant care, consuming any chemical or herbal medicines known to affect sexual desire, smoking, or any substance abuse.

Attrition criteria

Breakdown of the satisfactory emotional relationship with the husband during the study, unwillingness to continue participating in the study, using other treatments for sexual dysfunction or other problems during the study, low compliance or adherence to instructions, and developing skin allergy or other intolerable complications.

The participants were selected from eligible women using convenience consecutive sampling. After explaining the objectives of the study, the women who were willing to participate first reported their age, age of marriage, education level, and sexual functioning. Then, by block randomization, they were randomly assigned to the experimental or control groups.

In the experimental group, qust oil (Tooba Company) was delivered to the patients, who were instructed to massage 20 drops on the pubic area, twice a day for 4 weeks, in a back and forth motion for 5 minutes (superficial and without pressure) and 3 drops on the perineum for 1 minute in a circular motion. In the control group, liquid paraffin (manufactured by Merck) was delivered to the patients, who were instructed to massage 20 drops on the pubic area,

twice a day for 4 weeks, in a back and forth motion for 5 minutes (superficial and without pressure) and 3 drops on the perineum area for 1 minute in a circular motion. Both groups were followed up by phone every week to monitor their regular use of the drugs and to check any side effects. At the end of the fourth week, the two groups were again examined in terms of sexual functioning and severity of menopausal symptoms. The primary outcome was change in sexual desire, and the secondary outcome was arousal, lubrication, orgasm, satisfaction, pain during intercourse, and overall sexual function, as evaluated by the FSFI four weeks after the start of the intervention¹⁴.

Data analysis

SPSS v. 27 was used for the data analysis. Chi-square test and Fisher's exact test were used in order to compare the two groups in terms of qualitative variables (education level, occupation, and complications). In order to compare the group in terms of sexual functioning, the normality of the data distribution was first checked using the Kolmogorov-Smirnov test. Based on the test result and rejection of the null hypothesis of normal data distribution, Mann-Whitney U test was used to compare the two groups. The significance level was set at 95% ($P < 0.05$).

Results

Among 120 postmenopausal women with sexual dysfunction, ten were excluded due to not meeting the inclusion criteria, and 110 women were included in the study in two groups of 55. Three women from the experimental group and six from the control group dropped out during the study. Finally, 52 women in the experimental group and 49 in the control group remained until the final analysis (Figure 1).

The mean and SD of the age of the participants was 54.35 ± 5.06 years (range: 43-65 years) and the duration of marriage was 31.66 ± 8.21 years (range: 1.47 years). In terms of education level, in both experimental and control groups, most of the participants had a high school diploma, followed by higher education. In terms of occupation, in both groups, most women were homemakers. No significant difference was observed between the experimental and control groups in terms of occupation and education level (Table 1).

There was no significant difference between the two groups in terms of sexual function before the treatment, except for satisfaction. The level of sexual satisfaction of the women

was significantly lower in the experimental compared to the control group ($p=0.039$). After the intervention, significant differences were observed between the experimental and control groups in sexual functioning in the domains of sexual desire ($p<0.001$), arousal ($p<0.001$), orgasm ($p<0.001$), satisfaction, and overall score ($p<0.001$). Meanwhile, no significant difference was observed in lubrication and pain during intercourse (Table 2, Figure 2).

One woman (2.0%) from the experimental group had complications (hives), which improved with oral treatment. No complications were observed in the control group. No significant difference was observed between the two groups in the frequency of complications ($p=0.485$).

Discussion

In the present study, the topical use of qust oil by postmenopausal women with sexual dysfunction for four weeks significantly increased sexual desire, arousal, orgasm, satisfaction, and overall sexual functioning; however, no significant difference was observed in lubrication and pain during intercourse. The researchers' search of electronic databases yielded no relevant or similar papers that had used qust oil topically to treat women's sexual dysfunction. Two human studies, *i.e.*, carried out by Sadeghi *et al.* (2020) on the oral effect of carrot seed and by Akhtari *et al.* (2014) on the effect of *Tribulus terrestris* extract on reduced sexual desire in women of reproductive age, reported favorable results in terms of increasing libido^{14, 16}. Similarly, two human studies were conducted by Preedy *et al.* (2021) and Akbar *et al.* (2020) on the effect of olive oil and valerian (a component of qust oil), respectively. Preedy *et al.* (2021), who examined the effect of olive oil on physical health, observed that short-term massaging with olive oil can increase sexual desire¹⁷. Akbar *et al.* (2020) examined the effects of *Valeriana jatamansi* and found that the root of this plant increases libido while reducing anxiety, depression, seizures, headaches, and menstrual cramps¹⁸. In-vitro animal studies have investigated the medicinal effects of qust oil or each of its components separately, yielding promising results¹⁹⁻²². Muchtaromah (2020) examined the combined protective and therapeutic effects of three herbs, including garlic, turmeric, and *A. calmus* (a component of qust oil), on mice. This combination increased the animal subjects' endometrial thickness, secretion of sex hormones, and fertility¹⁹. Mehanna *et al.* (2020) examined the oil extract of marjoram (a component of qust oil) and observed that with its strong antioxidant effects, this substance increases spermatogenesis and testosterone levels in

male rats suffering tissue damage and necrosis of the testicular cells²⁰. Mishra *et al.* (2016) examined clove extract (a component of qust oil) and showed that its administration at low doses (15 mg) has androgenic effects, increases testosterone levels, and rejuvenates sex genes²¹. As can be observed, each component of qust oil has been examined in human and animal studies, showing androgenic effects and influencing the sex hormones, testosterone levels, fertility, and sexual function. Our study was conducted on a human sample, and instead of examining the individual effects of the mentioned medicinal plants, we studied their combined effect (valerian, marjoram, cloves, *F. gummosa*, olive, nutmeg, *A. calmus*, bitter qust, and Chinese cinnamon) through a product known as qust oil, which significantly improved the sexual functioning of postmenopausal women; it can be argued that these substances have had a synergistic effect.

Another difference between the present study and most of the cited studies was the different method of qust oil administration (topical use via massage versus oral use). Regarding the positive effect of massage therapy on sexual desire, Chia *et al.* (2011) examined the therapeutic effect of massage on sexual desire and reported that massaging the pubic area increases libido and sexual desire.²² Therefore, the therapeutic effects of qust oil alone in sexual dysfunction and its massaging on the pubic area may increase the effectiveness of this compound. Meanwhile, this route of administration will have no side effects potentially arising through oral consumption. As for other uses of qust oil, the results of studies have shown that rubbing qust oil on the pubic area is effective in treating bed wetting in children²³ and urinary incontinence in women²⁴. In two review studies and meta-analyses, Karimi *et al.* and Zalchour *et al.* investigated the effect of medicinal plants on sexual functioning. The review study by Karimi *et al.* (2023) in Iran carried out a systematic review of clinical trials to investigate the effect of medicinal plants on sexual functioning in Iranian postmenopausal women. The studied plants included *Ginkgo biloba*, fennel, *Hypericum perforatum*, date palm pollen, Aphrodite, fenugreek, common hop, black cohosh, lavender, ginseng, saffron, red clover, *T. terrestris*, *Nigella sativa*, and *Vitex agnus-castus*, and combined aromatherapy (fennel, *Heracleum persicum*, common sage, bitter orange, lavender, and geranium). The results showed that most plants were effective in improving sexual functioning²⁵. In a review and meta-analysis by Zalchour *et al.* (2023), the effect of medicinal plants on the sexual functioning and satisfaction of postmenopausal women in Iran and other countries was investigated. The meta-analysis showed that herbal medicines significantly increase the overall sexual functioning score and the sexual satisfaction score of

postmenopausal women²⁶. To investigate the pharmacological and phytochemical properties of qust oil, Kumari *et al.* (2024) carried out a phytochemical analysis of the product in India and showed that *Saussurea costus* (Falc) Lipsch contains a large number of bioactive compounds, including sesquiterpenes, flavonoids, and essential oils, which are responsible for its numerous medicinal properties. Pharmacological studies have shown anti-inflammatory, antioxidant, anti-cancer, hepatoprotective, and immune-modulating effects for this product. As a result, it is widely used in the treatment of diseases such as asthma, digestive disorders, and skin diseases. *S. costus* has significant therapeutic potential, mainly owing to its rich phytochemical composition. The convergence of its traditional uses and modern pharmacological findings offers promising avenues for future research, especially in drug development and understanding its mechanism of action in various diseases²⁷. According to the study by Bahrami *et al.* (2018) on the characteristics of thick vessels, energy channels, meridians, and chakras, it can be concluded that massaging and stimulating the second pair of the thick vessels lead to connection and balance of the cephalo uterin axis and favorable results in the treatment of decreased sexual desire²⁸. In the present study, no complications (itching, rash, burning, or hives) were observed as a result of qust oil massaging on the pubic and perineal area, which suggests the safety of the topical use of qust oil to treat sexual dysfunction in postmenopausal women.

Conclusions

In postmenopausal women with sexual dysfunction, massaging the pubic area and perineum with qust oil for at least four weeks has no side effects and significantly improves the components of desire, arousal, orgasm, and sexual satisfaction.

List of abbreviations:

FSFI, Female sexual function index

FSH, Follicle-Stimulating Hormone

LH, Luteinizing Hormone

Correspondence: Elham Akhtari, Associate professor, Department of Traditional Medicine, Institute for Studies in Medical History, Persian and Complementary Medicine, School of Persian Medicine, Iran University of Medical Sciences, Tehran, Iran.

E-mail: eli.akhtari@gmail.com

ORCID: 0000-0003-4491-6443

Co authors:

Akram Fallah

E-mail: Akram.fallah@gmail.com

ORCID:0009-0006-0356-4890

Fattaneh Hashem Dabbaghian

E-mail: dabaghian.f@iums.ac.ir

ORCID:0000-0003-0005-5792

Amirhossein Jamshidi

E-mail: jamshidi.ah@iums.ac.ir

ORCID:0000-0003-1310-1516

Conflict of interest: the authors declare no potential conflict of interest, and all authors confirm accuracy.

Ethics approval: the Ethics Committee of Iran University of Medical Sciences under the code of ethics : IR.IUMS.REC.1402.378 . The Iranian Registers of Clinical Trial (IRCT) code :IRCT 2023730058969NI.The study is conformed with the Helsinki Declaration of 1964, as revised in 2013, concerning human and animal rights.

Informed consent: all patients participating in this study signed a written informed consent form for participating in this study.

Patient consent for publication: written informed consent was obtained from a legally authorized representative(s) for anonymized patient information to be published in this article.

Availability of data and materials: all data generated or analyzed during this study are included in this published article.

Acknowledgements

the authors express their gratitude to the Research and Technology Deputy of the university, the Faculty of Traditional Medicine, the Traditional Medicine Center, and the participants, for their financial and non-financial support. The authors declare that they have no conflicts of interest.

References

1. Nappi RE, Lachowsky M. Menopause and sexuality: prevalence of symptoms and impact on quality of life. *Maturitas* 2009;63:138-41.
2. Berek J. Berek & Novak Gynecology. Translated by Ghazijahani B, Zonuzi A, Bahrami N. Tehran, Iran: Golban publication; 2007: pp. 471-501.
3. Bagheriani N, Bahrami M, Kamalinejad M, et al. The effect of rheum ribes root (rhubarb) on menopausal hot flashes: a randomized double-blind, placebo-controlled trial. *Res J Pharmacogn* 2023;10:41-50.
4. Nappi RE, Cucinella L, Martella S, et al. Female sexual dysfunction (FSD): Prevalence and impact on quality of life (QoL). *Maturitas* 2016;94:87-91.
5. Koliji T, Keshavarz Z, Mojab F, et al. Comparison of Cinnamon and Vitex on improving sexual function in postmenopausal women: a triple-blind randomized clinical trial. *Jundishapur J Natural Pharmaceut Products* 2022;17(4).
6. Danforth DN. *Danforth's obstetrics and gynecology*: Lippincott williams & wilkins; 2008.
7. Niazi A, Moradi M. Sexual satisfaction and function in postmenopausal women treated with herbal medicines: a review of clinical trials. *Evidence Based Care* 2019;9:7-16.
8. Beigi M, Fahami F, Hassanzahraei R, Arman S. Associative factors to sexual dysfunction in menopause women. *Iranian J Nursing Midwifery Res* 2008;13(1).
9. Wright JJ, O'Connor KM. Female sexual dysfunction. *Med Clin* 2015;99:607-28.
10. Bradshaw K, Corton M, Halvorson L, et al. *Enfermedad mamaria-Williams Ginecología* [Internet]. New York (NY): McGraw-Hill Global Education Holdings LLC; 2017.
11. Balon R, Clayton AH. Female sexual interest/arousal disorder: a diagnosis out of thin air. *Arch Sexual Behavior* 2014;43:1227-9.
12. Aghili Khorasani M. *Summary of Wisdom (Kholasat Alhekma)*. Tehran. Ismaeilian Publications; 2006.

13. Velayati A. An introduction to the history of medicine in Islam and Iran. *Med J Islamic Republic of Iran (MJIRI)* 1988;2:131-6.
14. Akhtari E, Raisi F, Keshavarz M, et al. Tribulus terrestris for treatment of sexual dysfunction in women: randomized double-blind placebo-controlled study. *DARU J Pharmaceut Sci* 2014;22:1-7.
15. Ibn Sīnā A. The canon of medicine. 1025.
16. Sadeghi S, Bahrami R, Raisi F, et al. Evaluation of the effect of carrot seed (*Daucus Carota*) in women of fertile age with hypoactive sexual desire disorder: A randomized double-blind clinical trial. *Complement Ther Med* 2020;54:102543.
17. Preedy VR, Watson RR. Olives and olive oil in health and disease prevention: Academic press; 2020.
18. Fleming T. PDR for herbal medicines. 1998.
19. Muchtaromah B, Muti'ah R, Yusmalasari DR, et al. Efficacy of *Allium sativum*, *Curcuma mangga* and *Acorus calamus* extract combination on rat fertility. *Pharmacognosy J* 2020;12(1).
20. Mehanna S, Issa MY, Hassan NH, et al. *Origanum majorana* essential oil improves the rat's sexual behavior and testicular oxidative damage induced by imidacloprid via modulating the steroidogenesis pathways. *Saudi Pharmaceut J* 2022;30:1315-26.
21. Mishra R, Singh S. Biphasic effect of *Syzygium aromaticum* flower bud on reproductive physiology of male mice. *Andrologia* 2016;48:1011-20.
22. Chia M. *Karsai Nei Tsang: Therapeutic Massage for the Sexual Organs*: Simon and Schuster; 2011.
23. Motaharifard MS, Effatpanah M, Nejatbakhsh F. Nocturnal enuresis in children and its herbal remedies in medieval persia: a narrative review. *J Pediatr Rev* 2020;8:15-22.
24. Niktabe Z, Eftekhaar T, Mirabzadeh Ardakani M, et al. Efficacy of *Saussurea costus* (qost) oil as an Iranian traditional medicine product on female urinary incontinence; double blinded randomized clinical trial. *Res J Pharmacogn* 2018;5:47-55.
25. Karimi FZ, Salehian M, Hosseini H, et al. The Effect of the medicinal plants on sexual function in Postmenopausal Women in Iran: a systematic review of clinical trials. *J Sabzevar University Med Sci* 2023;30:144-62.
26. Zalpour A, Zare E, Nasiri M. The Effects of Herbal Medicines on Sexual Function and Sexual Satisfaction among Women: A Systematic Review and Meta-Analysis. *Jundishapur J Natural Pharmaceut Products* 2023 (In Press).

27. Kumari R, Negi M, Thakur P, et al. *Saussurea costus* (Falc.) Lipsch.: a comprehensive review of its pharmacology, phytochemicals, ethnobotanical uses, and therapeutic potential. *Naunyn-Schmiedeberg's Arch Pharmacol* 2024;397:1505-24.
28. Bahrami M, Mastery Farahani R, Nazem E, et al. The body organs and their reconstruction power (regeneration) from the viewpoint of Iranian-islamic physicians. *Iran Red Crescent Med J* 2014;16:e15193.




Figure 1. CONSORT diagram of sample selection




Figure 2. The mean and standard deviation of the post-intervention overall score of sexual functioning by group

Table 1. Frequency of the participants' education level and occupation divided by group

Variable	Group		P-value
	Experimental Number (%)	Control Number (%)	
Education level			
Below high school diploma	5 (9.6)	9 (18.4)	0.302
High school diploma	28 (53.8)	20 (40.8)	
Associate or bachelor's degree	9 (17.3)	13 (26.5)	
Master's degree or higher	10 (19.3)	7 (14.3)	
Total	52 (100)	49 (100)	
Occupation			

Homemaker	33 (63.5)	35 (71.4)	0.120
Self-employed	7 (13.5)	1 (2.0)	
Governmental	12 (23.1)	13 (26.5)	
employment	52 (100)	49 (100)	
Total			

Table 2. The mean and standard deviation of the pre-intervention sexual functioning score by group

Variable	Group		P-value
	Experimental Mean±sd	Control Mean±sd	
Pre-intervention			
Desire	4.23±1.47	3.76±1.45	0.103
Arousal	7.40±3.08	8.73±2.74	0.283
Lubrication	9.47±2.73	9.68±2.95	0.667
Orgasm	7.07±2.28	6.65±2.72	0.293
Satisfaction	7.44±2.48	6.44±2.57	0.039
Pain during intercourse	8.03±2.71	7.67±2.54	0.380
Overall	44.67±11.59	41.94±13.51	0.277
Post-intervention			
Desire	6.13±1.31	4.41±1.53	<0.001
Arousal	12.37±2.42	7.80±2.80	<0.001
Lubrication	11.85±2.19	11.14±3.20	0.250
Orgasm	9.21±1.64	6.37±2.52	<0.001
Satisfaction	10.08±1.66	7.32±2.24	<0.001
Pain during intercourse	8.92±2.71	8.80±2.21	0.776
Overall	58.83±8.44	45.84±11.84	<0.001

*. Mann-Whitney U test

Submitted: 13 May 2024

Accepted: 13 June 2024

Early access: 16 September 2024