



eISSN 2039-4772

<https://www.pagepressjournals.org/index.php/chest/index>

**Publisher's disclaimer.** E-publishing ahead of print is increasingly important for the rapid dissemination of science. The **Early Access** service lets users access peer-reviewed articles well before print / regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

**Chest Disease Reports** is, therefore, e-publishing PDF files of an early version of manuscripts that have undergone a regular peer review and have been accepted for publication, but have not been through the typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one. The final version of the manuscript will then appear in a regular issue of the journal.

E-publishing of this PDF file has been approved by the authors.

*All legal disclaimers applicable to the journal apply to this production process as well.*

Chest Disease Reports 2024 [online ahead of print]

To cite this article:

Gaurav Pandey, Sonisha Gupta, Parul Singhal, et al. Germ cell tumor impersonating as a pleuroperecardial cyst: a rare phenomenon. Chest Disease Reports. 2024;12:12119. doi:10.4081/cdr.2024.12119

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*



©The Author(s), 2024

Licensee *PAGEPress*, Italy

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

## Germ cell tumor impersonating as a pleuropericardial cyst: a rare phenomenon

Gaurav Pandey<sup>1</sup>, Sonisha Gupta<sup>2</sup>, Parul Singhal<sup>3</sup>, Syed Haider Mehdi Rizvi<sup>1</sup>, Shubham Pandey<sup>4</sup>, Shweta Mishra<sup>5</sup>

<sup>1</sup>Student, Dept. Respiratory Medicine, School of Medical Sciences & Research, Gautam Budh Nagar;

<sup>2</sup>Professor, Dept. Respiratory Medicine, School of Medical Sciences & Research, Gautam Budh Nagar;

<sup>3</sup>Professor, Dept. of Pathology, Saraswati Institute of Medical Sciences, Hapur;

<sup>4</sup>Junior Resident Doctor, Dept. Of Emergency Medicine, Satyewadi Raja Harishchanda Hospital, Narela, New Delhi;

<sup>5</sup>Consultant Cranio-Facial Surgeon, Sharva Cosmoaesthetiq Clinique, Rohini, New Delhi, India;

**Corresponding author:** Shubham Pandey, Dept. Of Emergency Medicine, Satyewadi Raja Harishchanda Hospital, Narela, New Delhi 110040, India.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

Tel. 7838492484. E-mail: [drshubhampandey2286@gmail.com](mailto:drshubhampandey2286@gmail.com)

**Key words:** teratoma, mediastinum, chest wall.

**Authors' contributions:** all the authors made a substantive intellectual contribution.

All the authors have read and approved the final version of the manuscript and agreed to be held accountable for all aspects of the work.

**Conflict of interest:** the authors declare no potential conflict of interest.

**Funding:** none.

**Patient's consent for publication:** the patient gave his written consent to use his personal data for the publication of this case report and any accompanying images.

**Informed consent:** written informed consent was obtained from a legally authorized representative for anonymized patient information to be published in this case report.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

**Availability of data and materials:** all data generated or analyzed during this study are included in this published article.

**Acknowledgments:** the authors would like to thank Dr. Saurav for the constant support and guidance.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

## **Abstract**

A Mixed Germ Cell Tumor (MGCT) is a rare tumor with a low degree of differentiation. The most common sites are yolk sac and astrocytoma. Usually, it is gonadal, but in 5% of cases, it can present as extragonadal. The most common extragonadal site is the mediastinum, where the mediastinal mass sometimes mimics thymoma, lymphoma, pericardial cyst, and also occasionally causes pericardial tamponade. The presentations vary, ranging from accidental findings on routine radiography to life-threatening respiratory and cardiovascular compromise. We present an extragonadal MGCT case presenting as a mediastinal mass with symptoms of 1 year duration. The diagnosis was confirmed on Histopathological Examination (HPE) of surgically excised specimen. Given a possible life-threatening condition, a timely diagnosis is required.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

## **Introduction**

A Mixed Germ Cell Tumor (MGCT) is a rare tumor form of two or more germ cell components, with a low degree of differentiation.<sup>1</sup> Usually, MGCT occurs in ovary and testis, and is rarely found at extragonadal sites like head and neck, mediastinum, and abdomen.

Germ cell tumors usually originate from the gonads and are derived from reproductive cells embryologically. But in approximately 4-5% of all the cases, they have extragonadal origin, with the most common extragonadal site being the mediastinum. Primary mediastinal non-seminomatous germ cell tumor is very rare, approximately 4-5% of all germ cell tumors. The presentations vary, ranging from accidental findings on routine radiography to life-threatening respiratory and cardiovascular compromise.<sup>2</sup>

Pericardial cysts are rare, with an incidence rate of 1 in 100000. Usually, they are congenital but, in some cases, can present as acquired pericardial anomalies (*e.g.* post-inflammatory, hydatid, neoplastic). Pericardial cysts are induced by an incomplete coalescence of fetal lacunae during the development of the pericardium.<sup>3</sup> Pericardial cysts are usually unilocular, well marginated spherical or teardrop shaped, and may be attached to the pericardium directly or by a pedicle. Of all pericardial cysts, 70 to 75% are located at the right cardio-phrenic angle, and the rest are on the left side of the mediastinum.<sup>4</sup>

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

The mediastinal mass sometimes mimics thymoma, lymphoma, pericardial cyst, and also occasionally causes pericardial tamponade, which needs proper investigation and timely treatment.

Herein, we report a case of mediastinal MGCT. We also reviewed the relevant literature, focusing on the Computed Tomography (CT) and clinical characteristics of mediastinal MGCT.

### **Case Report**

A 42-year-old male came in our Outpatient Department (OPD) in June 2022, with complaints of chronic dry cough, progressive shortness of breath on exertion, and weight loss for the past 1 year.

The patient had visited multiple medical setups for these complaints, and received different treatments, including Anti-Tuberculosis Treatment (ATT), which he stopped on his own after further decline in his health. Eventually he came to our OPD. On physical examination, the patient was found to be in good clinical condition, conscious and oriented to time, place and person, with Blood Pressure (BP) 120/80 mmHg, Heart Rate (HR) 82/min & SPO2 97% on room air.

On respiratory examination, diminished breath sounds on the left lower lung area were

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*



found on auscultation. Chest X-Ray (Figure 1) showed high density, oval shaped opacity, situated in the right para-cardiac area with anterior projection. Ultrasound of the thorax revealed a pleuro-pericardial cyst located tangentially to the right chest wall.

Contrast Enhanced Computed Tomography (CECT ) thorax (Figure 2) described a well-defined, non-enhancing hypodense (25-30 HU) lesion (55\*55\*95mm) in the anterior mediastinum right paracardiac region, at the level of right atrium, abutting the pericardium with mild indentation. It was also causing smooth indentation on the underlying lung parenchyma. The lesion shows peripheral intermittent thick calcified rim with non-enhancing hyper dense content in the inferior portion. Multiple nodules soft tissue attenuating clustered lesion are seen in the vicinity of the aforementioned lesion in right paracardiac fat, measuring 51\*49\*32 mm. Laboratory reports were normal. Two Sputum smears for Acid Fast Bacilli (AFB) were negative. IgG antibodies specific for Echinococcus was negative. A provisional diagnosis was made as right pleuropericardial cyst, and the patient was advised for surgical intervention.

The patient underwent surgical intervention and right para mediastinal mass was taken out (Figure 3). On Histopathological Examination (HPE), it was diagnosed as a MGCT consisting of seminoma, embryonal carcinoma and mature teratoma. (Figure 4, Figure 5) Hematoxylin and eosin-stained sections showed features of seminoma and embryonal cell carcinoma, respectively. Pus culture taken from para-mediastinal mass

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

shows no growth. Post procedural Chest X-Ray and CECT Chest showed no visible space occupying mass lesion.

Post Procedural chest X-Ray (Figure 6), CT chest (Figure 7) and PET CT scan (Figure 8) were done. Positron Emission Tomography-Computed Tomography (PET-CT) was done mainly to rule out any metastatic possibilities, which showed mildly Fluorodeoxyglucose (FDG) avid soft tissue thickening in the anterior mediastinal region with non-FDG avid nodular densities in upper lobes of both lungs and in left lung lingula with mild right sided pleural effusion. The immunoassay done for Alpha Fetoprotein (AFP) (3.65 ng/ml) and Beta Human Chorionic Gonadotropin (HCG) (7.02 IU/L) was found within normal range.

The patient was advised a reexamination after 6 months, and regular follow up of the case was done till that time.

## **Discussion**

Germ cell tumors develop embryologically from the reproductive cells. They primarily arise from the gonadal organs. Since only 5% cases of the germ cell tumors are extragonadal in origin, it is uncommon to identify these tumors.<sup>5</sup> The most common extragonadal sites include the mediastinum, retro-peritoneum, vagina, and brain.<sup>6</sup>

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

Clinical characteristics range from life-threatening respiratory and cardiovascular impairment, to unintentional discoveries on routine radiography. Such massive tumors can cause symptoms like coughing, shortness of breath, and chest pain, or they can perforate and cause other symptoms, including pleural and pericardial effusion.<sup>7,8</sup>

Mediastinal tumor are of different types, *e.g.* teratoma, seminoma, embryonal carcinoma, choriocarcinoma, MGCT.

The Teratoma is classified into: i) mature teratoma; ii) immature teratoma – benign; iii) immature teratoma - with malignant component.

Immature teratoma with malignant component is further classified into: i) Type I, with another Giant Cell Tumor (GCT) [seminoma, Yolk Sac Tumor (YST), Embryonal Carcinoma (EC), choriocarcinoma]; ii) Type II, with a malignant epithelial neoplasm (adenocarcinoma, squamous cell carcinoma, *etc.*); iii) Type III, with a malignant mesenchymal component (rhabdomyosarcoma, angiosarcoma, *etc.*); iv) Type IV, any combination of previous types.<sup>9</sup>

Anterior mediastinal tumors account for 50% of all the mediastinal masses, that includes thymoma, teratoma, thyroid tumors and lymphomas.<sup>10</sup>

Thymoma is the most commonly occurring anterior mediastinal mass in the adult population, although it accounts for less than 1% of adult malignancies.<sup>11</sup> Thymomas mainly afflict people over the age of 40; they are uncommon in youngsters and equally

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

affect both males and women. Between 20% and 30% of thymoma patients experience symptoms as a result of pressure.<sup>12</sup>

The anterior mediastinum is where primary mediastinal lymphoma typically develops. Nearly 20% of all mediastinum neoplasms in adults and 50% in children are malignant lymphomas. The most frequent cause of masses in the pediatric mediastinum is lymphomas.<sup>13</sup>

Hodgkin Disease (HD) has a bimodal distribution of incidence, peaking in young adulthood and again after the age of 50 years.<sup>14</sup> The T-cell lymphoblastic lymphoma is a form of mediastinal Non-Hodgkin Disease (NHD) that mainly occurs in children and adolescents.<sup>15</sup> A differential diagnosis for mediastinal masses are pericardial cysts which are often unilocular, well-marginated, spherical or teardrop-shaped lesions that may be linked to the pericardium directly or by a pedicle. At the time of diagnosis, 85% of patients with mediastinal Non Seminomatous Germ Cell Tumor (NSGCTs) are symptomatic. The most frequent presenting symptoms are chest discomfort, hemoptysis, cough, fever, or weight loss. There may occasionally be pericardial tamponade or superior vena cava syndrome.<sup>4,16</sup> Anterior mediastinal NSGCT presenting as early cardiac tamponade is rarely seen in clinical practice.<sup>17</sup>

At the time of presentation, cough, chest discomfort, fever/chills, and dyspnea are common symptoms. Systemic symptoms are often brought on by the production of too

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

many hormones, antibodies, or cytokines, whereas localizing symptoms are secondary to tumor invasion (respiratory compromise, paralysis of the limbs, diaphragm, and vocal cords, Horner syndrome, superior vena cava syndrome).<sup>18</sup>

In our case, the presenting complaints of chronic dry cough, progressive shortness of breath on exertion and weight loss for past 1 year were either due to tumor-induced compression or the total atelectasis of the lung, following recurrent pulmonary infections. Benign mediastinal teratomas may be secondarily infected following rupture into nearby structures. Chest X-Ray and CECT scan are the preferred investigation for screening and diagnosis.

But, in this situation, a preoperative biopsy would have been useful, although it was not recommended due to the location of the tumor. The acknowledged standard practice for confirming a histological diagnosis is fine needle core biopsy.<sup>19</sup> In germ cell tumors, tumors markers such as beta-HCG, AFP, and Lactate Dehydrogenase (LDH) are typically increased. When evaluating the effectiveness of chemotherapy, especially in chemo-sensitive GCT, tumors marker testing is essential. Raised serum AFP levels suggest that mixed germ cell tumors contain yolk sac and embryonal components.<sup>20</sup>

The preferred surgical intervention is a thoracotomy, which is done from the side of the tumor where it has more extension. The median sternotomy can also be used when the tumor is tiny and there are few adhesions to nearby important organs.<sup>21</sup> Surgical

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

excision was done and on HPE, a diagnosis of mixed germ cell tumor having components of seminoma, embryonal cell carcinoma and teratoma was made.

## **Conclusions**

Although benign mediastinal teratomas are rare, they may present as a chest wall mass producing bulging from mediastinum towards the chest wall. This slow growing tumor is usually asymptomatic and is often detected incidentally on chest X-Ray. Echocardiography and chest CT is helpful to evaluate compression of adjacent structures. In cases of cystic masses, containing fat and calcific densities presenting in unusual locations, the diagnosis of teratoma should be considered. The treatment is surgical excision along with proper investigations to be done to make the final diagnosis. Though they cannot be removed totally, their recurrence rate is very low.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

## References

1. Hu X, Li D, Xia J, et al. Mediastinal mixed germ cell tumor: A case report and literature review. *Open Medicine*. 2021;16: 892-8.
2. Hayati F, Ali NM, Kesu Belani L, et al. Giant mediastinal germ cell tumor: an enigma of surgical consideration. *Case Reports in Surgery*. 2016;2016.
3. Akbayrak H, Yildirim S, Simsek M, Oc M. A rare giant pericardial cyst mimicking a paracardiac mass. *Cardiovascular Journal of Africa*. 2016;27:7-9.
4. Nina VJ, Manzano NC, Mendes VG, Salgado Filho N. Giant pericardial cyst: case report. *Brazilian Journal of Cardiovascular Surgery*. 2007;22:349-51.
5. McKenney JK, Heerema-McKenney A, Rouse RV. Extragonadal germ cell tumors: a review with emphasis on pathologic features, clinical prognostic variables, and differential diagnostic considerations. *Advances in Anatomic Pathology*. 2007;14:69-92.
6. Lin S, Li X, Sun C, et al. CT findings of intrarenal Yolk sac tumor with tumor thrombus extending into the inferior vena cava: a case report. *Korean Journal of Radiology*. 2014;15:641-5.
7. Miyazawa M, Yoshida K, Komatsu K, et al. Mediastinal mature teratoma with rupture into pleural cavity due to blunt trauma. *The Annals of Thoracic Surgery*.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

2012;93:990-2.

8. Yoshida K, Yamada T, Aoki T, et al. A case of mature teratoma perforated into the pericardial cavity. [Zasshi][Journal]. Nihon Kyobu Geka Gakkai. 1997;45:1107-10.

9. Kalhor N, Moran CA. Primary germ cell tumors of the mediastinum: a review. Available from: <https://med.amegroups.org/article/view/3961/4719>

10. Tomiyama N, Honda O, Tsubamoto M, et al. Anterior mediastinal tumors: diagnostic accuracy of CT and MRI. European journal of radiology. 2009;69:280-8.

11. Juanpere S, Cañete N, Ortuño P, et al. A diagnostic approach to the mediastinal masses. Insights into imaging. 2013;4:29-52.

12. Benveniste MF, Rosado-de-Christenson ML, Sabloff BS, et al. Role of imaging in the diagnosis, staging, and treatment of thymoma. Radiographics. 2011;31:1847-61.

13. Franco A, Mody NS, Meza MP. Imaging evaluation of pediatric mediastinal masses. Radiologic Clinics. 2005;43:325-53.

14. Duwe BV, Sterman DH, Musani AI. Tumors of the mediastinum. Chest. 2005;128:2893-909.

15. Takahashi K, Al-Janabi NJ. Computed tomography and magnetic resonance imaging of mediastinal tumors. J Magn Reson Imaging. 2010;32:1325-39.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*



16. Mishra S, Majumdar SK, Sable M, Parida DK. Primary malignant mediastinal germ cell tumors: a single institutional experience. *South Asian Journal of Cancer*. 2020;9:27-9.
17. Skelton WP, Mahtta D, Welniak S, et al. Pericardial effusion as an atypical initial presentation of extra-gonadal nonseminomatous germ cell tumor: a case report and literature review. *Oxford Medical Case Reports*. 2018;2018:omx097.
18. Juanpere S, Cañete N, Ortuño P, et al. A diagnostic approach to the mediastinal masses. *Insights Imaging*. 2013;4:29-52
19. Chhieng DC, Lin O, Moran CA, et al. Fine-needle aspiration biopsy of nonteratomatous germ cell tumors of the mediastinum. *American Journal Of Clinical Pathology*. 2002;118:418-24.
20. Sirohi B, Huddart R. The management of poor-prognosis, non-seminomatous germ-cell tumors. *Clinical Oncology*. 2005;17:543-52.
21. Soysal O, Sarac K, Kutlu R, et al. A case of mediastinal teratoma presenting as a cystic lesion on chest wall. *East J Med*. 1998;3:32-3.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*




**Figure 1.** Pre-procedural chest X-Ray.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*



**Figure 2.** Pre-procedural Computed Tomography (CT) of chest.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*




**Figure 3.** Post procedural specimen.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*




**Figure 4.** Histopathological Examination (HPE) showing seminoma.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*




**Figure 5.** Histopathological Examination (HPE) showing embryonal cell CA.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*




**Figure 6.** Post procedural chest X-Ray.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*




**Figure 7.** Post procedural Computed Tomography (CT) of chest.

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*





**Figure 8.** Post Procedural Positron Emission Tomography-Computed Tomography (PET-CT).

*The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.*

*All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers.*

*Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*