

# The importance of psychological intervention in emergencies for patients with anorexia nervosa

Giacomo Fiacco, Gennaro Catone

Department of Educational, Psychological and Communication Sciences, University Suor Orsola Benincasa, Naples, Italy

Dear Editor,

We consider it appropriate to highlight the importance of early psychological intervention in emergencies for patients with eating disorders. In the contemporary age, characterized by incessant socio-cultural pressures and aesthetic ideals, eating disorders emerge as a complex and critical challenge to mental and physical health in individuals during developmental age. This psychiatric condition can have fatal outcomes. Approximately one-fifth to one-third of adolescent patients with Anorexia Nervosa (AN) require intensive care during the course of the illness. Access to critical care areas in healthcare facilities due to significant emaciation-related organic decompensation is increasing significantly, particularly after the COVID-19 pandemic. Typically, emergency room access for these patients is ensured by family members or caregivers who can recognize the critical developments of the eating disorder. However, many patients, despite being in clinically severe conditions, are unable to identify the seriousness of their

clinical condition. The treatment proposed by healthcare personnel often encounters opposition from the patient, who may refuse initial care, thereby risking their own life. Clinical emergency conditions related to anorexia nervosa include electrolyte imbalances with cardiac arrhythmias, dysproteinemia with the potential for pericardial effusion, and multi-organ functional compromise. Recent systematic reviews highlight a lack of criteria, objectives, and treatment modalities, sometimes providing an ambiguous account of hospitalization practices. We believe that a consensus conference should be initiated to outline a systematic, shared approach to the emergency treatment of critically ill patients with eating disorders. In this context, the role of an expert clinical psychologist can be fundamental, even in critical care settings. The presence of such professionals could: i) better facilitate the proposed therapeutic approach, which is often life-saving; ii) interact with the family and caregiver; iii) liaise with emergency personnel treating the patient. This collaboration aims at a more rapid clinical treatment, reducing possible oppositional attitudes that hinder care. The presence of an expert psychologist could optimize the assistance and treatment times for patients with AN in critical care.

Correspondence: Giacomo Fiacco, Department of Educational, Psychological and Communication Sciences, University Suor Orsola Benincasa, Naples, Italy  
E-mail: [giacomofcc@outlook.com](mailto:giacomofcc@outlook.com)

Key words: anorexia, eating disorders, emergency.

Contributions: all the authors made a substantive intellectual contribution. All the authors have read and approved the final version of the manuscript and agreed to be held accountable for all aspects of the work.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: none.

Availability of data and materials: all data generated or analyzed during this study are included in this published article.

Received: 16 June 2024.  
Accepted: 20 August 2024.  
Early view: 17 September 2024.

This work is licensed under a Creative Commons Attribution 4.0 License (by-nc 4.0).

©Copyright: the Author(s), 2024  
Licensee PAGEPress, Italy  
Emergency Care Journal 2024; 20:12726  
doi:10.4081/ecj.2024.12726

*Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

## References

1. Herpertz-Dahlmann B. Intensive treatments in adolescent Anorexia Nervosa. *Nutrients* 2021;13:1265.
2. Vaccaro CM, Guarino G, Conte D, et al. Social networks and eating disorders during the Covid-19 pandemic. *Open Med (Wars)* 2021;16:1170-4.
3. Toppino F, Martini M, Longo P, et al. Inpatient treatments for adults with anorexia nervosa: a systematic review of literature. *Eat Weight Disord* 2024;29:38.