



eISSN 2282-2054

<https://www.pagepressjournals.org/index.php/ecj/index>

**Publisher's Disclaimer.** E-publishing ahead of print is increasingly important for the rapid dissemination of science. The **Early Access** service lets users access peer-reviewed articles well before print / regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

The **Emergency Care Journal** is, therefore, e-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear on a regular issue of the journal.

E-publishing of this PDF file has been approved by the authors.

Emerg Care J 2024 [Online ahead of print]

*To cite this Article:*

Poggiali E, Giosia V, Coen D. **A red and warm abdomen. What's the matter?** *Emerg Care J* doi: 10.4081/ecj.2024.13175

 ©The Author(s), 2024

Licensee [PAGEPress](#), Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

## **A red and warm abdomen. What's the matter?**

Erika Poggiali,<sup>1</sup> Valeria Giosia,<sup>1</sup> Daniele Coen<sup>2</sup>

<sup>1</sup>Emergency Medicine Unit, Fatebenefratelli Hospital, ASST Fatebenefratelli Sacco, Milan;

<sup>2</sup>Academy of Emergency Medicine and Care, Pavia, Italy.

**Correspondence:** Erika Poggiali, M.D. Emergency Medicine Unit, Fatebenefratelli Hospital, ASST Fatebenefratelli Sacco, Milan, Italy.

E-mail: [poggiali.erika@gmail.com](mailto:poggiali.erika@gmail.com)

**Key words:** skin, cellulitis, skin infection, Streptococcus pyogenes, Staphylococcus aureus.

**Contributions:** All the authors approved the final version and equally contributed to the work.

**Conflicts of interest:** EP and DC are members of the editorial board of ECJ. The authors declare no conflict of interest.

**Availability of data and materials:** All data underlying the findings are fully available upon reasonable request to Erika Poggiali, [erika.poggiali@asst-fbf-sacco.it](mailto:erika.poggiali@asst-fbf-sacco.it)

**Ethics approval and consent to participate:** As this was a descriptive case report and data was collected without patient identifiers, ethics approval was not required under our hospital's Institutional Review Board guidelines.

**Informed consent:** The patient provided consent for the access to medical records at the time of admission.



### **Descriptive legend**

An 83-year-old woman presented to our emergency department for a fever with nausea and an itching skin rash on her abdomen for two days. She suffered from obesity, Parkinson disease, and depression in treatment with Levodopa/Benserazide and Pramipexole, respectively. She underwent mastectomy for left breast cancer ten years earlier. She denied any allergies or use of antibiotics. Her vital signs were blood pressure 120/65 mmHg, heart rate 72 bpm, peripheral oxygen saturation 94% in room air,

respiratory rate 18/min, and body temperature 36.2 °C. Examination of her abdomen revealed an extensive red warm rash without blisters or blebs, with a slight skin oedema confirmed by point-of-care ultrasound, that excluded a subcutaneous abscess, pneumoniae, liver and kidney damage, and cholecystitis. Laboratory findings showed WBC 12,400/mm<sup>3</sup> (N 59.5%) and CRP 131 mg/L (n.v. < 5). Procalcitonin was 0.3 ng/mL (n.v. < 0.5). SARS-CoV-2 infection was excluded.

**Question:** Given the patient history and the clinical presentation, what is the most likely diagnosis?

- 1) Lipodermatosclerosis
- 2) Cellulitis
- 3) Papular urticaria
- 4) Allergic contact dermatitis

**Answer**

The right answer is cellulitis. Cellulitis is a common, potentially serious bacterial infection that involves the deep dermis and subcutaneous tissue. It represents the most common reason for skin-related hospitalization,<sup>1</sup> and it can be a diagnostic and therapeutic challenge for emergency clinicians.<sup>2</sup> Specific criteria do not exist for the diagnosis, that is always clinical with an acute onset of redness, warmth, swelling, tenderness, and pain.<sup>3</sup> Patients can complain of fever, chills, and general malaise. Cellulitis usually affects the lower legs, but it can occur on the face, arms, and other body areas. Risk factors include skin injury (any cut, burn, scrape) or skin condition (atopic dermatitis,

athlete's foot), lymphedema, venous insufficiency, overweight, immunodepression, and a previous history of cellulitis.<sup>4</sup> Treatment may include antibiotics and pain management, based on paracetamol or ibuprofen for mild to moderate pain, and morphine if severe. The majority of non-purulent, uncomplicated cases of cellulitis are caused by  $\beta$ -haemolytic streptococci, followed by methicillin-sensitive *Staphylococcus aureus*.<sup>2</sup> Failure to improve with appropriate first-line antibiotics should prompt consideration for resistant organisms - methicillin-resistant *Staphylococcus aureus* (MRSA)<sup>5</sup> - or other conditions that can mimic cellulitis.<sup>6,7</sup> If untreated or misdiagnosed, cellulitis can become an emergency for the development of septic shock, endocarditis, osteomyelitis, gangrene, and toxic shock syndrome. In some rare cases, the infection can spread to the fascial lining, causing necrotizing fasciitis.<sup>8</sup> Our patient was admitted to the Internal Medicine Unit and treated with Piperacillin/Tazobactam, with complete resolution of the cellulitis in seven days.

## References

1. Peterson RA, Polgreen LA, Cavanaugh JE, Polgreen PM. Increasing incidence, cost, and seasonality in patients hospitalized for cellulitis. *Open Forum Infect Dis* 2017;4:ofx008.
2. Boettler MA, Kaffenberger BH, Chung CG. Cellulitis: a review of current practice guidelines and differentiation from pseudocellulitis. *Am J Clin Dermatol* 2022;23:153-65.
3. Keller EC, Tomecki KJ, Alraies MC. Distinguishing cellulitis from its mimics. *Cleve Clin J Med* 2012;79:547-52.
4. Björnsdóttir S, Gottfredsson M, Thórisdóttir AS, et al. Risk factors for acute cellulitis of the lower limb: a prospective case-control study. *Clin Infect Dis* 2005;41:1416-22.
5. Raff AB, Kroshinsky D. Cellulitis: A Review. *JAMA* 2016;316:325-37.
6. Nazarko L. Red legs: how to differentiate between cellulitis, venous eczema and lipodermatosclerosis. *Br J Community Nurs* 2022;27:486-94.

7. Goldenberg M, Wang H, Walker T, Kaffenberger BH. Clinical and immunologic differences in cellulitis vs. pseudocellulitis. *Expert Rev Clin Immunol* 2021;17:1003-13.
8. Poggiali E, Santilli G, and Vercelli A. A rapidly progressive painful skin lesion. *Emerg Care J* 2023;19:11046.