



eISSN: 2281-7824

<https://www.pagepressjournals.org/index.php/hls/index>

Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. The **Early Access** service lets users access peer-reviewed articles well before print / regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

The **Healthcare in Low-resource Settings** is, therefore, e-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear on a regular issue of the journal.

E-publishing of this PDF file has been approved by the authors.

Healthc Low-resour S 2024 [Online ahead of print]

To cite this Article:

Setiyowati E, Firdaus F, Wardani EM, Mulyadi M. **Monitoring medication adherence using smart digital technology in patients with pulmonary tuberculosis.** *Healthc Low-resour S* doi: 10.4081/hls.2024.11995

 ©The Author(s), 2024

Licensee [PAGEPress](#), Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Monitoring medication adherence using smart digital technology in patients with pulmonary tuberculosis

Eppy Setiyowati,¹ Firdaus Firdaus,¹ Erika Martining Wardani,¹ Mulyadi Mulyadi^{2,3}

¹Faculty of Nursing and Midwifery, Universitas Nahdlatul Ulama, Surabaya; ²Faculty of Medicine, Universitas Nahdlatul Ulama, Surabaya; ³Jemursari Islamic Hospital, Surabaya, Indonesia

Correspondence: Eppy Setiyowati, Faculty of Nursing and Midwifery, Universitas Nahdlatul Ulama, Surabaya, Indonesia.

E-mail: eppy@unusa.ac.id

Key words: pulmonary tuberculosis, smart digital, taking medicine.

Contributions: ES, conceptualization, data curation, formal analysis, methodology, validation, visualization, writing - original draft, review and editing; FF, conceptualization, investigation, methodology, validation, and writing - original draft, review and editing; EMW, conceptualization, methodology, formal analysis, validation, and writing - original draft, review and editing; MM, methodology, visualization, writing - review and editing.

Conflict of interest: the authors declare no conflict of interest.

Ethics approval and consent to participate: this research has received ethical approval from the Health Research Ethics Commission, chakra Brahmanda Lentera Institution, based on ethical certificate 027/011/V/EC/KEP/LCBL/2023. During the research, the researchers paid attention to the ethical principles of information to consent, respect for human rights, beneficence and non-maleficence.

Patient consent for publication: written informed consent was obtained for anonymized patient information to be published in this article.

Funding: this research was supported by a research grant from Universitas Nahdlatul Ulama Surabaya with contract number 458/UNUSA/Adm-LPPM/IV/2023.

Availability of data and materials: all data generated or analyzed during this study are included in this published article.

Acknowledgments: the authors would like to thank Rector Universitas Nahdlatul Ulama Surabaya for their valuable insights and contributions to this study.

Abstract

Pulmonary tuberculosis cases can arise due to various factors, including treatment fatigue resulting from prolonged therapy, insufficient patient knowledge about the condition, geographical distance between patients' residences and healthcare facilities, a lack of reminders from healthcare workers regarding medication adherence, and the perception that

treatment provided at community health centers may be inadequate. This study aimed to investigate the effectiveness of digital smart applications in enhancing medication adherence among pulmonary tuberculosis patients. Utilizing a trial survey approach, the research focused on patients seeking treatment at the Surabaya Community Health Center. Smart digital applications were employed to monitor medication adherence, with patients inputting their data *via* cell phones. The study recorded data over one month and found that while some patients struggled to operationalize the smart digital application, many were able to utilize it effectively. Overall, the findings suggest that smart digital applications can be valuable tools for monitoring medication adherence among pulmonary tuberculosis patients, particularly for older adults.

Introduction

Tuberculosis (TB) remains a significant global health challenge, particularly prevalent in developing countries.^{1,2} Factors contributing to the high number of TB cases include treatment fatigue resulting from prolonged therapy,³ inadequate patient knowledge about pulmonary TB,^{4,5} geographical barriers to accessing healthcare facilities, a lack of reminders from healthcare providers regarding treatment adherence,⁶ and perceptions of treatment quality at community health centers.^{7,8} These factors can lead to treatment non-adherence among TB patients.^{9,10}

The compliance of TB patients with their treatment regimen plays a pivotal role in the management of the disease.¹¹ Studies have shown that non-adherence to treatment increases the risk of TB recurrence by about 2.5 times, highlighting challenges from medical, immunological, and psychological perspectives.¹² Treatment non-adherence contributes to a higher rate of treatment failure among pulmonary TB patients and exacerbates the burden of TB within communities.¹³

Factors influencing adherence to pulmonary TB medication include understanding of the treatment instructions,¹⁴ quality of interactions with healthcare providers, family support, personal beliefs, attitudes, and personality traits.¹⁵ Additionally, the lengthy and rigorous treatment regimen for pulmonary TB, spanning 6-8 months, poses another significant challenge.¹⁶

To enhance compliance with TB treatment, the use of directly observed treatment (DOT) by drug supervisors has been implemented.¹⁷ However, traditional DOT methods have not effectively addressed TB management challenges.¹⁸ Therefore, there is a need for innovative approaches to drug supervision, such as utilizing electronic and communication media to directly monitor drug intake by patients in the presence of healthcare workers. While health education and telenursing have been utilized to increase knowledge among individuals with pulmonary TB,¹⁹ they have not fully addressed long-term drug adherence challenges. In order to ensure sustained drug adherence over the entire treatment duration, which typically spans at least 6 months, the development and implementation of smart digital applications for TB patients are essential.^{19,20}

Previous studies have examined the use of digital smart applications for assessing medication adherence in various settings.^{21,22} However, these studies did not reveal the knowledge, attitudes, and behaviors of TB patients. This study aimed to specifically analyze the use of digital smart applications for controlling drug adherence in pulmonary TB patients.

Materials and Methods

Design and sample

This study employed a trial survey approach utilizing digital smart applications to monitor medication adherence. It was conducted at a community health center in Surabaya, targeting TB patients who regularly seek treatment there each month. Initially, there were eleven patients

identified at the health center and its branches in the area. However, after coordination and collaboration with the TB program team, it was found that three patients had already been declared cured. Subsequently, the study focused on the remaining nine pulmonary TB patients who possessed digital media in the form of cell phones. Data collection commenced from May to August 2023, beginning with the assessment of respondents' behaviors (knowledge, attitudes, and actions) related to medication intake and adherence.

Instruments

The methods used to measure adherence, including patient self-reports, pill counts, refill levels, biological monitoring, and electronic monitoring, have their limitations, particularly concerning reminder sounds displayed on digital smart devices. The reports displayed typically rely on patients' personal data, memory, and medication schedules provided by the treating doctor. Although the name and quantity of the medication are recorded, they are not always factored into the data analysis and processing. Reminder sounds are programmed to alert patients at the same time each day, but some respondents may not be sensitive to these reminders, leading to instances where medications are not taken as prescribed. All data entered into the smart digital system is recorded for one month, from the 1st to the 31st of August. Monitoring medication intake through digital smart devices includes recording patient names, ages, the names of the medications taken, and the quantity of medications consumed through electronic monitoring.

Medication adherence behavior was assessed based on two indicators: adherence and non-adherence. Adherence is recorded if there is consistent charging on the smart digital device for one consecutive month. Each respondent's cell phone was equipped with a digital smart application called Pamio TB. Following installation, researchers provided detailed explanations of the mechanism and operation of the Pamio TB smart digital application to the respondents.

Development of adherence to taking medications

Some methods aim to improve medication adherence among patients with pulmonary TB. However, this study specifically focuses on the implementation of a digital smart application called Pamio TB (short for "compliant to taking TB medicine"). This application includes an alarm that sounds every day at 7 a.m., reminding respondents to take their medication. When using the digital smart application, respondents input their identity, the name of the drug, and the quantity of medication taken. Additionally, the application features a questionnaire designed to assess the knowledge, attitudes, and actions of individuals with pulmonary TB. Knowledge levels are categorized as high, medium, or low, while attitudes and actions are assessed using similar indicators.

Habit of reminding

In patients with pulmonary TB, the medication-taking process requires attention from both healthcare workers and family members. Often, a family member serves as a medication supervisor to ensure adherence. However, the use of digital smart applications can potentially reduce the need for this role. When it is time to take medication, the alarm in the digital smart app will ring, prompting the respondent to press a button on their cell phone. This alarm serves as a reminder for individuals with pulmonary TB to take their medication, thus minimizing their reliance on external supervision.

Analysis

Utilizing a digital smartphone application represents a novel approach to enhancing medication adherence among individuals with pulmonary TB. This application offers constant accessibility, actively engages patients, and serves as a repository for pertinent medication adherence-related information. With the presence of alarms on patients' smartphones, reminders to take medication are automatic, eliminating the need for family members to prompt medication intake.

By employing digital smart applications for monitoring medication adherence in pulmonary TB patients, there is a shift towards a more patient-centered approach. This transition highlights the potential value of such applications as effective tools for monitoring medication adherence, reducing reliance on external supervisors, and empowering individuals with pulmonary TB to take charge of their treatment regimen.

Results and Discussion

The discussion on medication adherence among patients with pulmonary TB encompasses various aspects, drawing insights from existing literature and research findings. However, in this paper, the authors focus on areas related to improving drug adherence through digital smart applications to improve medication adherence and reduce the risk of anti-TB drug withdrawal. Developing a digital smart application to monitor medication intake aims to prevent patients from discontinuing their anti-TB drugs.²¹ This app ensures adherence to medication schedules and minimizes confusion, particularly among elderly patients. Therefore, this paper introduces a smart digital application designed to monitor adherence to TB treatment in patients with pulmonary TB.²¹ The proposed application features audible alerts from the patient's mobile alarm set for specific times.

Table 1 shows a multidisciplinary approach aimed at establishing a medication adherence monitoring system using an effective and reliable digital smart application. The integration of Internet of Things and wireless body area networks represents a significant advancement in healthcare, particularly for patients with pulmonary TB, enhancing the quality of healthcare services and the overall quality of life for these individuals.²³ Essentially, smart digital applications facilitate healthcare services by facilitating the collection, aggregation, transmission, and analysis of medical data.²⁴

This digital smart application proposes a design for monitoring medication adherence in patients with pulmonary TB, with implementation trials planned for 2023. The implementation of this application is intended to address the concerns of caregivers, ensuring the security of full treatment. Through automatic reminders, the application will prompt patients at the correct time for medication intake, minimizing the risk of errors in dosage and medication type.

Conclusions

This study proposes and implements smart digital applications to monitor medication adherence in elderly patients with pulmonary TB, providing significant assistance. The smart digital application automatically prompts patients at the onset of pulmonary TB, reminding them of the timing for medication intake. This feature is particularly beneficial for elderly patients, ensuring they remember to take their medicine on time.

References

1. Kemenkes RI. Laporan hasil riset kesehatan dasar (riskesdas) Indonesia tahun 2018.

- Riset Kesehatan Dasar. [Material in Indonesian].
2. Opperman M, Du Preez I. Factors contributing to pulmonary TB treatment lost to follow up in developing countries: an overview. *African J Infect Dis* 2023;17:60-73.
 3. Putra ON, Yulistiani Y, Soedarsono S. Scoping review: QT interval prolongation in regimen containing bedaquiline and delamanid in patients with drug-resistant tuberculosis. *Int J Mycobacteriol* 2022;11:349-55.
 4. Zhang J, Yang Y, Qiao X, et al. Factors influencing medication nonadherence to pulmonary tuberculosis treatment in Tibet, China: a qualitative study from the patient perspective. *Patient Prefer Adherence* 2020;14:1149-58.
 5. Zahroh C, Ainiyah N, Saadah N, et al. Determinant of incompliance medication people with tuberculosis disease. *Indian J Forensic Med Toxicol* 2021;15:909-14.
 6. Abqari U, van 't Noordende AT, Richardus JH, et al. Strategies to promote the use of online health applications for early detection and raising awareness of chronic diseases among members of the general public: a systematic literature review. *Int J Med Inform* 2022;162:104737.
 7. Dayer L, Heldenbrand S, Anderson P, et al. Smartphone medication adherence apps: potential benefits to patients and providers. *J Am Pharm Assoc* 2013;53:172-81.
 8. Dosi G, Pereira MC, Roventini A, Virgillito ME. Technological paradigms, labour creation and destruction in a multi-sector agent-based model. *Res Policy* 2022;51:104565.
 9. Lepuen AP, Bratajaya CNA, Rasmada S. Tuberculosis cases finding practice: the intention of Cadres. *J Keperawatan Indones* 2020;23:128-35.
 10. WHO. Primary health care on the road to universal health coverage: 2019 global monitoring report. Executive summary. Available from: <https://www.who.int/publications/i/item/9789240029040>.
 11. Rahmadi C, Efendi F, Makhfudli M. The effect of reminder system and audio-visual education to improve treatment adherence on pulmonary tuberculosis patients: a systematic review. *Mal J Med Heal Sci* 2023;19:145-55.
 12. Zein RA, Suhariadi F, Hendriani W. Estimating the effect of lay knowledge and prior contact with pulmonary TB patients, on health-belief model in a high-risk pulmonary TB transmission population. *Psychol Res Behav Manag* 2017;10:187-94.
 13. Goos M, Rademakers E, Roettger R. Routine-biased technical change: Individual-level evidence from a plant closure. *Res Policy* 2021;50:104002.
 14. Nugrahaeni DK, Rosmalaningrum L. Risk factors in pulmonary tuberculosis treatment failure. *Indones J Public Heal* 2021;16:12.
 15. Sukartini T, Purwanti ND, Mariyanti H. Family health tasks implementation and medication adherence of pulmonary tuberculosis patients: a correlational study. *J Ners* 2020;15:49-58.
 16. Putra IWGAE, Kurniasari NMD, Dewi NPEP, et al. The implementation of early detection in tuberculosis contact investigation to improve case finding. *J Epidemiol Glob Health* 2019;9:191-7.
 17. Zachariah R, Spielmann MP, Harries AD, et al. Passive versus active tuberculosis case finding and isoniazid preventive therapy among household contacts in a rural district of Malawi. *Int J Tuberc Lung Dis* 2003;7:1033-9.
 18. Gashu KD, Gelaye KA, Lester R, Tilahun B. Effect of a phone reminder system on patient-centered tuberculosis treatment adherence among adults in Northwest Ethiopia: a randomised controlled trial. *BMJ Heal Care Inform* 2021;28:e100268.
 19. Banu S, Haque F, Ahmed S, et al. Social enterprise model (SEM) for private sector tuberculosis screening and care in Bangladesh. *PLoS One* 2020;15:e0241437.

20. Eastment MC, McClintock AH, McKinney CM, et al. Factors that influence treatment completion for latent tuberculosis infection. *J Am Board Fam Med* 2017;30:520-7.
 21. Wong YJ, Ng KY, Lee SWH. Digital health use in latent tuberculosis infection care: a systematic review. *Int J Med Inform* 2022;159:104687.
 22. Sholikhah DU, Sari GM, Narendri CM, et al. The use of health technology to enhance the adherence of tuberculosis treatment: a systematic review. *J Ners* 2019;14:65-70.
 23. Thoumrungroje A, Racela OC. Innovation and performance implications of customer-orientation across different business strategy types. *J Open Innov Technol Mark Complex* 2022;8:178.
- .

Table 1. Attributes and ratings of the five apps that were installed and subjected to user testing.

Application name	Online data entry	Complex instruction	Cloud data storage	Database of medication	Multiplatform app	Final record data	Final posting record	iPhone
My med schedule Pamio TB	X	X	X	X	X	1	4	X
MyMeds	X		X	X	X	2	4	X
Med Agenda		X	X	X	X	3	5	X
Remid Me Prescription		X			X	4	4	X
Med Memory				X	X	5	4	X

Submitted: 17 October 2023

Accepted: 16 May 2024

Early access: 5 June 2024