

The role of family support in managing death anxiety among the elderly

Niko Dima Kristianingrum, Yati Sri Hayati, Annisa Wuri Kartika, Fransiska Imavike Fevriasanty, Selsa Rifziqka Heltha Novian Haryono, Ayut Merdikawati

Nursing Department, Faculty of Health Sciences, Universitas Brawijaya, Malang, East Java, Indonesia

Abstract

Death anxiety is a sentimental state experienced by individuals, specifically the elderly due to the occurrence of unpleasant events and contemplation of mortality. This condition is primarily caused by a lack of social support from family, which has a significant effect on mental health. Therefore, this study aimed to analyze the correlation between family support and the level of death anxiety in the elderly. The study procedures were carried out with a cross-sectional method, and the sample population comprised 239 elderly who were selected at a community health center in Malang Indonesia using cluster random sampling. Family support was measured using the Family Support Instrument (FSI), while death anxiety level was assessed with the Death Anxiety Scale (DAS). The relationship between the 2 variables was then tested using Spearman's Rank correlation test with p<0.05. The results showed that there was a strong correlation between the 2 variables, with family support falling into the high category for 98 participants (41%). In addition, 117 participants (49.0%) were reported to have a low category of death anxiety level. To conclude, these findings could be applied and used as a reference for future studies by exploring variables and conditions influencing family support and the death anxiety level of the elderly.

Introduction

Life expectancy is an essential factor, which has been reported to have a positive correlation with health challenges, particularly among the elderly. According to the World Health Organization (WHO), there is a substantial increase in the prevalence of both physical and mental diseases among this demographic, posing a significant burden on the society.1 In addition, the phenomenon of an aging populace and the increase in the number of the elderly in Indonesia have a significant impact on the country, families, society, and aged individuals.2 Several studies have shown that aging exerts adverse effects across various facets of life, including balance in the elderly. This demographic is often affected by biopsychosocial conditions, which interact to cause various health challenges, including psychiatric disorders, with anxiety being the most commonly diagnosed.^{3,4,5} A prevalent type of anxiety among the elderly is death anxiety, which is characterized by apprehension toward mortality and contemplation about death.^{6,7}

A previous study regarding death anxiety level was conducted at a nursing home in Bandung Indonesia using 79 elderly, with 50.6% of the participants experiencing high level. In addition, this condition was primarily caused by a lack of social support. Among the various elements of social support, the role of family is essential in determining the mental health status of the elderly. P.10 A previous study showed that the elderly living in nursing homes have a higher tendency to experience death anxiety compared to others with their families. Optimal support has also been reported to have the potential to alleviate this condition, particularly among those undergoing hemodialysis. Based on findings, there has been an increasing trend of the elderly living in nursing homes, specifically in Asian countries. Therefore, this study aimed to analyze the correlation between family support and death anxiety level among the elderly.

Materials and Methods

This study was carried out using a cross-sectional method, exploring 239 elderly taken from 4 working areas of the Ciptomulyo Community Health Center at Malang Indonesia using a cluster random sampling. Data collection was carried out in December 2022 through the use of a questionnaire. In addition, family support was measured using a modification of previous scales and analyzed for a validity test value of ≥ 0.361 as well as a reliable test with a Cronbach's Alpha value of 0.910. Death anxiety level was assessed using the Death Anxiety Scale (DAS).14 The scale consisted of 15 questions, which had been modified based on the study's needs with a validity test value of ≥ 0.361 and reliability with a Cronbach's Alpha value of 0.845. The correlation between the 2 variables was tested using the Spearman Rank test. Ethical approval letter was issued by the Health Research Ethics Committee of the Faculty of Health Sciences, Brawijaya University no. 5851/UN10.F17.10/TU/2022.

Results and Discussion

The results showed that the level of family support in this study was in the high category, as shown in Table 1. This was in line with Rahmadeni *et al.*, where the elderly did not experience depression due to optimum support from their family. ¹⁵ Moreover, the elderly typically experienced poor quality of life in the pres-

Significance for public health

The results of this study are expected to contribute to the management of death anxiety, with a particular emphasis on the role and support of family in the community. This is specifically pertinent considering the increasing number of the elderly and the growing trend of family placing the elderly in nursing homes.





Table 1. Family support frequency distribution.

Family support	n	%	
Low	60	25.1	
Moderate	81	33.9	
High	98	41.0	
Total	239	100.0	

Table 2. Elderly death anxiety level.

Death anxiety level	n	%	
Low	117	49.0	
Moderate	73	30.5	
High	49	20.5	
Total	239	100.0	

Table 3. Spearman Rank analysis

Death		Lo	Low		Anxiety level Moderate		High		Total	
		F	%	F	%	F	%	\mathbf{F}	%	
Family	support									
Low	7	2	0.84	25	10.46	33	13.81	60	25.10	
Mod	derate	26	10.88	40	16.74	15	6.28	81	33.89	
High	h	89	37.24	8	3.35	1	0.42	98	41.00	
Total	117	48.95	73	30.55	49	20.5	239	100.00		

Spearman's rho r= -0.683** p-value 0.000

ence of dysfunctional family. ¹⁶ Indonesians believed that these individuals were members of the family who must be respected. Younger members tried to meet all the needs of the elderly, including their health needs. When the physical and mental needs were met, these individuals typically felt. In the context of facing death, the elderly were often calmer in facing this challenge considering that their family was always present. ^{17,18,19}

Death anxiety level of the elderly in this study was placed in the low category, as shown in Table 2. In addition, it was reported to be influenced by factors other than family support, including spiritual health, number of hospitalizations, and the performance of health workers.9 The elderly often spend their time praying and participating in religious activities in the community. This was performed as a way to get closer to God in preparation for facing death, apart from filling free time. The better the spiritual health, the lower death anxiety level experienced by the participants. 20,21,22 In terms of the number of hospitalizations, the participants who suffered from chronic diseases and were frequently hospitalized were likely to experience higher death anxiety compared to others. This was because the elderly people who were treated in hospitals felt more lonely and uncomfortable with therapy procedures as well as a crowded hospital environment. 23,24,25 In addition, the performance of health workers could also affect death anxiety level, particularly less professional nurses. Based on these findings, there was a need for special training for nurses to increase understanding and readiness to face death, specifically for geriatric patients.²⁶

Table 3 presents the results of statistical correlation testing between family support and death anxiety in the elderly. The findings showed that there was a strong relationship between these variables with p < 0.05 and a correlation coefficient of -0.683. The higher the family support given to the elderly, the lower the death anxiety level experienced. This was in line with previous studies conducted in Arabia and Rural China, where death anxiety level experienced by the older population was in the low category while living with their families. Other studies also reported family support as a significant coping, influencing factor, and was found to have a protective effect among the elderly. 29,30,31,32

Conclusions

In conclusion, death anxiety level experienced by the elderly was influenced by physical and psychological factors, including family role in supporting the readiness to face peaceful death. In addition, active participation was needed from health facilities and health workers in providing professional service performance, particularly for geriatric patients.

References

- WHO. Ageing and health. 2022 [cited 2024 Jun 19]. Available from: https://www.who.int/news-room/fact-sheets/detail/ageing-and-health.
- Rahayu HC, Purwantoro, Wibowo M, Safitri J, Suparjito. Determinants of Welfare for the Elderly Population in Indonesia. KnE Medicine 2022;782–93.
- Apak E, Artan T, Ozucelik DN. Evaluation of bio-psychosocial and socio-cultural problems of Syrian elderly living in Turkey. Int J Soc Psychiatry 2023;69:454-66.
- Marwat DMK, Usman DK, Ullah MI, et al. Causes and Bio-Psycho-Social Changes Associated with Aging: A Review of Qualitative Literature. OEconomia 2023;6:1100–8.
- 5. Kułak-Bejda A, Bejda G, Waszkiewicz N. Mental Disorders, Cognitive Impairment and the Risk of Suicide in Older Adults. Front Psychiatry 2021;12:695286.
- Sahin D, Buken N. Death Anxiety And Concept Of Good Death In The Elderly. Turkish Journal Of Geriatrics -Turk Geriatri Dergisi 2020;23(1).
- Sharma M, Jain V. Death anxiety, religiosity and psychological

 ProQuest [Internet]. [cited 2023 Dec 23]. Available from: https://www.proquest.com/docview/2445574935?pq-origsite=gscholar&fromopenview=true&sourcetype=Scholarly%20Journals
- 8. Ningrum TP, Okatiranti O, Nurhayati S. Description of the Anxiety Level About Death in the Elderly at BPSTW Ciparay, Bandung Regency. Jurnal Keperawatan BSI 2018;6:142–9.





- Zahedi Bidgol Z, Tagharrobi Z, Sooki Z, Sharifi K. Death Anxiety and Its Predictors Among Older Adults. J Holist Nurs Midwifery 2020;101–10.
- Eslami RS, Omranian M. Social Support and Psychological Well-being of the Elderly: The Mediating Role of Death Anxiety. J Soc Sci Humanit Res 2021;9:97–107.
- 11. Ergin E, Yildirim D, Yildiz CÇ, Usenmez SY. The Relationship of Death Anxiety With Loneliness and Psychological Well-Being in the Elderly Living in a Nursing Home. Omega (Westport) 2023;88:333–46.
- 12. Dewi EI, Sriharja B, Deviantony F, et al. The Relationship Between Family Support And Death Anxiety In The Elderly Undergoing Hemodialysis. UNEJ e-Proceeding 2023;109–15.
- He AJ, Tang VFY. Integration of health services for the elderly in Asia: A scoping review of Hong Kong, Singapore, Malaysia, Indonesia. Health Policy 2021;125:351–62.
- Conte HR, Weiner MB, Plutchik R. Measuring death anxiety: conceptual, psychometric, and factor-analytic aspects. J Pers Soc Psychol 1982;43:775–85.
- Rahmadeni AS, Hayat N, Alba AD, et al. The Relationship of Family Social Support with Depression Levels of Elderly in 2019. Internat J Health Med Sci 2020;3:111–6.
- Souza Júnior EV de, Viana ER, Cruz DP, et al. Relationship between family functionality and the quality of life of the elderly. Rev Bras Enferm 2021;75:e20210106.
- Lestari MD, Stephens C, Morison T. Local knowledge and unliveable narratives: How insights from family caregiving narratives can inform locally relevant ageing policy. J Aging Studies 2023;64:101102.
- 18. Pio TMT, Prihanto JB, Jahan Y, Hirose N, Kazawa K, Moriyama M. Assessing Burden, Anxiety, Depression, and Quality of Life among Caregivers of Hemodialysis Patients in Indonesia: A Cross-Sectional Study. Internat J Environ Res Public Health 2022;19:4544.
- 19. Nugraha A, Rahmah GU, Budhiaji P. Effect of Family Support Towards Elderly Hemodialysis Patients' Anxiety Level. Basic Appl Nursing Res J 2020;1:20–4.
- Ghasemi F, Atarodi A, Hosseini SS. The Relationship Between Religious Attitudes and Death Anxiety in the Elderly People. Journal of Research and Health. 2020 May 10;10(3):135–42.
- Rababa M, Hayajneh AA, Bani-Iss W. Association of Death Anxiety with Spiritual Well-Being and Religious Coping in Older Adults During the COVID-19 Pandemic. J Relig Health 2021;60:50–63.
- 22. Soriano GP, Calong Calong KA, Martinez RCKP, Locsin RC. Religiosity, spirituality and death anxiety among older adults living at home and in institutional facilities: a comparative study. Journal of Religion, Spirituality & Aging 2022;34:284– 96.
- 23. Tadi A, Gheibizadeh M, Anjiri SGC. Death Anxiety and Associated Factors in the Hospitalized and Non-hospitalized Elderly with Chronic Diseases in Ahvaz. Jundishapur J Chronic Dis Care 2022;11(4).
- 24. Becker TD. Health care utilization as a predictor of death anxiety in older adults. Death Studies 2022;46:728–37.
- 25. Üstükuş A, Eskimez Z. The effect of death anxiety in nurses on their approach to dying patients: A cross-sectional study. Perspect Psychiatr Care 2021;57:1929–36.
- Batı S, Polat HT, Akkuş H. Determination of the Relationship Between Self-Care Agency and Death Anxiety Among Elderly Individuals. Omega (Westport) 2022:302228221095907.
- 27. Sebea DE, El-Geneidy MM, Abd-Elsalam RM, Dawood. Factors Associated with Death Anxiety among Community

- Dwelling Older Adults. Alexandria Scientific Nurs J 2021;23:1–17.
- 28. Pei Y, Cong Z, Silverstein M, Li S, Wu B. Factors Associated with Death Anxiety Among Rural Chinese Older Adults: The Terror Management Perspective. Res Aging 2022;44:65–72.
- 29. Raziani Y, Mahdavi A, Ngoc Huy DT, et al. Death Anxiety in the Iranian Elderly: A Systematic Review and Meta-Analysis Study. Omega (Westport) 2022;302228221110965.
- Laguilles-Villafuerte S. Death Anxiety and Interment Stress Family Interventions for Filipino Older Adults. Omega (Westport) 2022;302228221132905.
- 31. Laguilles-Villafuerte S, de Guzman AB. Put your hands in mine: a metaphor analysis of family support in death anxiety and interment stress among Filipino older adults. Educat Gerontol 2021;47:180–91.
- Jalal NM. Description Of Elderly Anxiety In The Face Of Death. Internat J Educat Vocational Soc Sci 2023;2:185–90

Correspondence: Fransiska Imavike Fevriasanty, Nursing Department, Faculty of Health Sciences, Universitas Brawijaya, Jl. Puncak Dieng, Kunci, Kalisongo, Kec. Dau, Malang, East Java, Indonesia, Postcode: 65151.

Tel.: +62341569117 - Fax: +62341564755.

E-mail: imavike.fk@ub.ac.id

Key words: death anxiety; elderly; family support

Contributions: the authors contributed equally from the literature reviews, data collection process, statistical analysis to manuscript editing.

Conflict of interest: the author declares no potential conflict of interest.

Funding: this study was supported by Universitas Brawijaya

Ethics approval: ethical approval letter was issued by the Health Research Ethics Committee of the Faculty of Health Sciences, Brawijaya University no. 5851/UN10.F17.10/TU/2022.

Clinical trials: not applicable.

Conference presentation: part of this article was presented at the 4th International Nursing and Health Sciences Symposium, from 27th-28th of October 2023, Universitas Brawijaya, Malang, East Java, Indonesia.

Acknowledgment: the authors are grateful to the Nursing Department, Faculty of Health Sciences Universitas Brawijaya for the opportunity provided to conduct this study in the geriatrics.

Received: 3 November 2023. Accepted: 10 September 2024.

This work is licensed under a Creative Commons Attribution 4.0 License (by-nc 4.0).

°Copyright: the Author(s), 2024 Licensee PAGEPress, Italy Healthcare in Low-resource Settings 2024; 12(s1):13022 doi:10.4081/hls.2024.13022

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

