

Spiritual coping “tri hita karana” and depression prevention behavior among the elderly during the COVID-19 pandemic

I Wayan Suardana,¹ Ah Yusuf,² Rachmat Hargono,³ I Gede Juanamasta,⁴ Ketut Sudiantara,¹ I Ketut Gama,¹
I Wayan Mustika¹

¹Polytechnic of Health Denpasar, Bali; ²Faculty of Nursing, Universitas Airlangga, Surabaya, East Java; ³Faculty of Public Health, Universitas Airlangga, Surabaya, East Java; ⁴Nursing Program, STIKes Wira Medika Bali, Bali, Indonesia

Abstract

The elderly commonly struggle with mental health issues, especially depression. Spiritual coping is one of the factors that might prevent depression. The aim of this study is to investigate cultural involvement in spiritual coping to depression prevention among the Balinese elderly. From January to March 2021, this study was carried out on 273 elderly members of the Integrated Primary Public Health service in seven districts in Gianyar, Bali. The study used a cluster random sampling method to collect data and focused on quantitative measurements related to participant characteristics, depression prevention, and spiritual coping. The investigation into spiritual coping was based on the *tri hita karana*, a concept from Balinese philosophy. Bivariate analysis was used to assess the relationship between spiritual coping and depression prevention behaviour. The study comprised 146 male and 127 female participants aged between 65 and 79 years old. Among all participants, 190 (69.6%) were diagnosed with mild to moderate depression, while 83 (30.4%) with moderate to severe depression. However, for the majority, depression prevention was good 163 (59.7%), the rest was moderate 103 (37.7%), and poor 7 (2.6%). *Tri hita karana* spiritual coping for the majority was good 17 (53.8%). Spiritual coping has a strong correlation with depression prevention ($r=0.230$). Thus, spiritual coping has a strong correlation with depression prevention among the Balinese elderly. It can be concluded that the implementation of *tri hita karana* becomes a potent spiritual coping enhance depression prevention. Indigenous philosophy embedded in daily life supports social capital in dealing with stress.

Introduction

Nowadays aging population become an urgent issue globally. The phenomena shifted from high income country into low and middle income countries.¹ United Nation projected in 2050 1 in 6 people in the world will be 65 or over. The trend of rising elderly prevalence is fasten in South East Asia and Latin America.² However the impact of this issue into public health system should be highlighted to encourage active aging where elderly will be

empowered to responsible maintenance their physical and mental wellness.³

Strengthen the future health system driven into elderly wellbeing which emphasized two integral aspects⁴. Various studies had been declared both aspects in physical and mental or psychosocial.⁴ The outcome of physical aspects of elderly such as the fitness to move during 3-6 month dependently.⁵ While the mental health outcome aspects mostly studied were quality of sleep, depression, and anxiety.

Elderly as vulnerable population has higher risk of isolation away from family due to immobility and other physical disability. Since 2019, Covid 19 pandemic in makes this reality surging among old population. Elderly feel alone, insecure, helpless, and lack of social support make higher risk of relapse depression and anxiety.^{5,6} However, depression declared as pathological mental development of getting old. Unlike the younger population, the risk of depression is higher 80% among older adults.⁷ This exacerbated and relapsed classified as major depressive disorder, persistent depressive disorder, substance induced depressive disorder and depressive disorder due to a medical condition.⁸

Management of depression in elderly is essential because it significantly related to family burden, decreased quality of life, and induced idea of suicide.⁸ Comprehensive clinical guideline has been established.⁹ First is comprehensive assessment related depression symptom and severity. The symptom listed based on the etiology such as number of medication trigger depression and other underlying disease. Depression severity mostly measured by GDS, HAM-D, BDI, PHQ-9. For a short relive anti depressant was widely used as pharmacological treatment. Another choice were recommended such as psychotherapy CBT, problem solving, therapy could involved family and caregiver. For the worst case electroconvulsive therapy might be chosen.

However concept of healing recently moved into seeking inner peace that authorized the elderly him/herself together creating harmony with social belongings also go beyond with the Supreme Being/God. This approach determined as spiritual coping. The core of spirituality is transcendence with 4 dimensions personal, social, environmental, and religious ones.¹⁰ Coping is a form of resilience that is individual in nature as a challenge to successfully pass every event in old age.¹¹ In its development, the concept of coping that exists and develops so far is still very little

Significance for public health

Promoting active ageing is crucial, emphasizing the need to address physical and mental health. Older people are at significant risk of depression, exacerbated by factors such as social isolation and the challenges posed by the COVID-19 pandemic. It is essential to implement effective management strategies, including comprehensive assessment and treatment options, to alleviate the burden of depression in this demographic. Furthermore, spiritual coping mechanisms, such as the Balinese philosophy of tri hita karana, have demonstrated the potential to promote well-being and resilience among older individuals. This study sought to explore the correlation between spiritual coping, specifically tri hita karana, and depression prevention behavior among older people during the pandemic.

discussing culture. Even if the concept of culture already exists, most of the literature still focuses on changes in the mental outlook of cultures that adopt western values and behavior (Divale, 2001). Novelty of this study is accentuated Balinese philosophy of harmony known as *Tri Hita Karana*.¹²

Conceptually, *Tri Hita Karana* is defined three causes of happiness.¹³ Three elements of *Tri Hita Karana* were relation human to God (theocentric) known as *Parahyangan*; relation human to human (anthropocentric) known as *Pawongan*; and relation human to nature (cosmocentric) known as *Palemahan*.¹⁴ One study found that the imbalance of these three things causes disease.¹⁵ In looking at the etiology of mental illness, it is more than fully looking at the type of abstract (non-physical) disease.^{13,15} Thus, *Tri Hita Karana* might have important part in wellness and health. Despite many studies used *Tri Hita Karana* concept specifically related to economic, politic, social and anthropology research,¹⁶ lack number of study employ this concept in the health research. Therefore, the study aimed to examine the relationship between spiritual coping "*Tri Hita Karana*" and depression prevention behaviour among elderly during pandemic Covid-19.

Materials and Methods

Study design

Design of this study is an observational and cross-sectional. Study was conducted during January- Marc 2021 with elderly population in Gianyar Distric area Bali Province. Gianyar was chosen as the research site because this area is a pilot for elderly-friendly implementation based on Bali local regulations. This site also well-known as the heart of Balinese culture. This study involved 7 district primary care provider in rural and urban area Payangan, Ubud, Sukawati, Tegalalang, Blahbatuh, Tampaksiring and Gianyar. Participants of this study was the elderly who joined regular health promotion program for primary care in Indonesia. Inclusion criteria of the participants is fluent to communicate in Bahasa. Exclusion criteria of the sample were severe mental disorder, hearing disorder, refuse to continue join the research. We used estimated proportion formula to count the total sample.¹⁷ Based on that formula the assumption probability 0.05 and elderly population in Ginyar Distric 2019 39,637 people, elderly depression prevalence in Indonesia 23,4 % (Riskasdas, 2018). According the calculation total sample involved 273 people. Data collection was performed by enumerator interviewed the elderly using structured questionnaires.

Instruments

In this study, 2 variables were measured, namely depression prevention behavior and religious coping. Depression prevention behavior is measured by giving 12 questionnaires with a Likert scale of 0-2 developed from Llopis & Ggabilondo.^{18,19} The validity and reliability have been tested. The items-total score of Pearson's correlation was significant at .05 level and with Content Validity Index for Items (I-CVI) computation was 1.00, with Cronbach's alpha was .80. Depression prevention behavior classified in three level: good (score 17-24), moderate (score 8-16), less (score 0-7).

Spiritual coping was self created questionnaire constructed based one Balinese philosophy *Tri Hita Karana*. *Tri Hita karana* spiritual coping complied 3 items element. First prahyangan element (relation with the supreme god/transcends): 5 items sradha (activity creating offering and praying) and 5 items bhakti (activity to do self reflection to be surrender). Second pawongan element

(relation and interaction with human being): 5 items nyama (support system among family) and 5 items (braya support system by neighborhood). Third palemahan element: 3 item utama mandala (space to do praying activity), 4 items madya mandala (space to do human interaction), and 3 items palemahan (green space for plants and pets).²⁰ Self-created questions were tested using Pearson's Correlation Product Moment for validity with the returned values varying between .412 and .572 in the commonality analysis, and for reliability through Cronbach's alpha, with all of the questions demonstrating reliability with the value of .583 (higher than 0.5). Another characteristic variables also measured such as gender, age, family type, personality, stress and quality of geriatric health provider.

Data analysis

The data was managed using the Statistical Package for Social Sciences - SPSS version 23.0. We performed the descriptive statistics for the socio-demographic and mental health characteristics, the depression prevention and spiritual coping. In the bivariate analysis, Pearson's correlation coefficient was used between characteristic variables and depression prevention. The relationship magnitude and strength were determined by the following criteria: $r = 0.10 - 0.29$ (small or low relationship), $0.30 - 0.49 =$ medium relationship, and $0.50 - 1.0 =$ strong or high relationship.^{21,22} The value of significance (2-tailed) is less than 0.05 significant at a 95% confidence interval.

Ethical considerations

This study was reviewed by The Health Research Ethics Committee Faculty of Nursing Universitas Airlangga, with the ethical approval number 2303-KEPK. Informed consent was distributed to the participants before completing the main questionnaires. The informed consent has been applied approaching the principle of beneficial, no harm, confidential, justice, as well as voluntary participants. The provided information should be agreed to by participants before data collection. The participant could stop the survey anytime during the data completion. The informed consent had been reported during the Ethics Approval and Consent prior the study.

Results and Discussion

Table 1 shows that majority of respondents were 70-79 years old and more than half were male. Primary education was the highest number compare than another level of education. More than 70% was married with unstable income. Extended family type was the majority of the respondent, followed by nuclear family and living alone. For depression prevention, 59.7% participants had a good category, 37.7% moderate and the rest was poor. Almost half of them had poor spiritual coping (46.2%). Based on the analysis, depression behavior prevention had a significant correlation with age ($r = 0.592$), gender ($r = 0.492$), income ($r = 0.468$) and spiritual coping ($r = 0.230$; Table 2). Surprisingly, the majority category was good during pandemic situation. This elaborate our correlational finding *Tri Hita Karana* Spiritual Coping has significant correlation with depression prevention behavior.

The majority of participants age on range 60-69 years old (83,15%). The rest (16,85%) is 70 years old and over. The cut point 70 years old in this study decided based on the prior study of elderly definition according physical and mental health independency among Balinese population.²³ This study congruence with elderly classification visited emergency department defined in three group

: youngest-old, ages 65 to 74 years; middle-old, 75 to 84 years; and oldest-old, ≥ 85 years.²⁴ Based on gender, the proportion of male population is 53,5% and female 46,5%. A study elaborated this finding based on correlational study in China related Asian Ethnic measured by life expectancy at age 60 years (LE60) and quality-adjusted life expectancy at age 60 years (QALE60).²⁵ This study found that men have as shorter life expectancy but seems less suffering rather than women.

Recently depression on of the most common mental health issue among elderly especially since pandemic Covid 19 until 2022.^{25,26} Study among 457 Indonesian participants found 53,57% experienced depression.^{27,28} Another study explained predictors of depression in South Asia (India, Pakistan, Bangladesh, Nepal, Sri Lanka) such as : gender, chronic illness, financial and physical dependent with other, not able to work or unemployment, experienced stress full event, lack of social support, marital status and living arrangement, substance or elderly abuse, inadequate spiritual needs, tension in home, not engage in leisure activities or hobbies.²⁹ Especially Bali Province as setting of this researched suffered a poignant effect of plunged tourism aspect as main income. Another study in Denpasar as part of Bali in first semester of 2020 among 100 respondents experienced anxiety (53,1%); depression (51,2%); and stress (46,1%) since decreasing monthly income and termination of employment.³⁰ Coherent with this research findings all participants experienced depression: mild to moderate 69,59% and moderate to severe 30,41%. Compared with previous study related depression in Bali Old population, the trend is about increasing. Study involved 774 female and 719 male participants aged 60 to 100 using CES-D score of ≥ 10 as depression as assignment declared the prevalence of depression among older adults in Bali was high (42.3%).³¹

However, old population have been faced many life stress exposure and experienced make the survived and well adapted known as coping.³² The main coping strategies main coping strategies were: anticipatory mourning, the desire to die, isolation, submission, negotiation, acceptance, accommodation, seeking social support, seeking spiritual comfort and living in the moment.³³ Even during pandemic COVID-19, old population showed their natural resilient dealing with unexpected condition and isolation by positive motioned coping mechanism by generate good mindset, stay busy, and look for social support. Compared with young population, old population less likely used avoidant coping strategy which sign of depressive symptom.³⁴

Explaining the former shape of coping was seed of transactional theory developed by Lazarus and Folkman in 1984.³⁵ Transactional explanations emphasizing dynamic relation of individual ability to responds stress of environment by cognitive phenomenological processes. Theory of Coping by Lazarus triggered by stress which interaction environment and person. Later stress stimulate perception named appraisal.³⁶ Appraisal conduct adaptation known coping mechanism classified in two: problem focused coping and emotional focused coping.³⁷ Problem focused coping described by using several strategy to remove the stressor by solving the problem.³⁸ While emotional focused coping described by using emotional responds to deal with the stressor.³⁹

Recently, holistic health orientation involved spirituality aspect as coping mechanism.⁴⁰ Philosophical analysis of spiritual coping in nursing science enhanced physical, psychological, and social well-being, resilience, and self-transcendence.^{41,42} Research suggests that spiritual coping strategies, involving relationship with self, others, Ultimate other/God or nature were found to help individuals to cope with their ailments.⁴³ Congruence with Balinese culture the definition of spirituality known as *Tri Hita*

Karana. embodies universal values and represents harmonious and balanced human relationships with the spiritual, social and natural environment to achieve spiritual and physical wellbeing.⁴⁴

Table 1. Characteristic of participants.

Variable (N=273)	n	%
Age		
65-69 years old	96	35.2
70-79 years old	177	64.8
Gender		
Male	146	53.5
Female	127	46.5
Highest level of education		
None	68	24.9
Primary	84	30.8
Secondary	45	16.5
Tertiary	44	16.1
University	32	11.7
Marital status		
Married	200	73.3
Divorced	64	23.4
Unmarried	9	3.3
Income status		
Stable	67	24.5
Unstable	206	75.5
Family type		
Nuclear family	116	42.5
Living alone	8	2.9
Extended family	149	54.6
Depression Prevention		
Good	163	59.7
Moderate	103	37.7
Poor	7	2.6
Spiritual coping		
Good	147	53.8
Poor	126	46.2

Table 2. Correlation characteristic and depression prevention behavior.

Variables	Category	Poor	Moderate	Good	Correlation
Age	High risk	3	46	47	0.592**
	Lower risk	4	57	116	
Gender	Male	3	52	91	0.492**
	Female	4	51	72	
Education level	None	1	33	34	0.082
	Primary	4	32	48	
	Secondary	2	10	33	
	Tertiary	0	19	25	
	University	0	9	23	
Income Status	Stable	4	68	108	0.468**
	Non-stable	3	35	38	
Marital Status	Married	7	64	129	0.008
	Unmarried	0	36	28	
	Divorced	0	3	6	
Family Type	Nuclear family	6	51	59	0.010
	Living alone	0	5	3	
	Extended family	1	47	101	
Spiritual coping	Good	5	40	102	0.230**
	Poor	2	63	61	

**p<0.01.

Implementation of *Tri Hita Karana* embedded in Balinese people daily life. Everyday people in Bali create handmade offering to show gratitude of The Supreme God Blessing. That activity became indigenous culture infused by Hinduism belief (*Sradha*) and compassion to serve (*Bhakti*).⁴⁵ Another implementation of *Tri Hita Karana* as Spiritual Coping strategy during Covid 19 was implemented by reading Bhagawad Githa involved 100 participants in Bali.⁴⁶ As part of harmony with the Supreme God Balinese people tend to be more surrender and more likely do self-control and reflection known as *Mulat sarira*. Harmony among human as human being in Bali conceptualized *Menyama Braya*. *Menyama*, its semantic meaning refers to family ties by marriage and blood; while *Braya* refers to neighborhood. This concept during pandemic became social capital build coping and resilience during pandemic.⁴⁷ Many people give free food supplies to others, giving social loan, and personal protective equipment's such as mask, face shield, disinfectant. Creating harmony in nature physically reflected on Balinese concept house always provide green space for *palemahan*. Balinese house divided into 3 elements as *Tri Mandala*: sacred place or temple for praying, home for human integration, and space for garden / pets.⁴⁷ Even in dynamic in rural area such as Denpasar as Bali Capital is still maintained.⁴⁷ Gardening and connect with animal or pets became habitual activity among Balinese people which available in their green space during isolation period in pandemic. Gardening engage can facilitate stress reduction through interactions with natural elements, which bolster human health.⁴⁸ The implementation of *Tri Hita Karana* in Bali as spiritual coping facilitates coping goal in this contacts of research related depression prevention. Old population in Bali more likely implement spiritual coping rather than problem or emotional focused coping. Based on this observation, the majority of participants did not have role as decision maker and lack of their cognitive capability to solve their life problem. Participants also try to hide their emotional feeling to prevent their family burden. Balinese elderly selected other activity to keep them busy and calm down their mind and feeling. Implementation of *Tri Hita Karana* not only for coping strategy but also outcome itself as depression prevention. Implementation *Tri Hita Karana* among old people in Bali occupied their physical, social, natural bonding activity to regulate mental health.

Conclusions

Age, gender, income, and spiritual coping were positively correlated with depression behavior prevention. This study results can give input to the nurse to understand depression behavior prevention. Besides, the results can use to develop intervention through *Tri Hita Karana* to manage depression behavior prevention. Community nursing or public health nurse could use this concept to promote positive mental health to community. Besides, it might be related to health seeking behavior. Nurse needs to explore more about mental health, which can be an issue to explore for future studies. This study's findings might not represent Indonesia because data were gathered only in the one district. Therefore, further research is needed with a larger, multiethnic sample to understand spiritual coping "*Tri Hita Karana*" and depression behavior prevention in Indonesia. This finding might be limited to during pandemic situation, further studies with longitudinal methodology is highly recommended. In addition, the study findings did not present cause effects; thus, a causal model to confirm spiritual coping and depression prevention behavior is needed for future studies.

References

1. WHO. Ageing and health [Internet]. 2022 [cited 2024 June 19]. Available from: <https://www.who.int/news-room/fact-sheets/detail/ageing-and-health>.
2. United Nations, Department of Economic and Social Affairs, Population Division. World Population Ageing 2019: Highlights (ST/ESA/SER.A/430). New York: WHO; 2019.
3. WHO. Decade of Healthy Ageing. World Heal Organ. New York: WHO;2020.
4. Paralikas T, Maria M, Dimitrios T, et al. Physical and Mental Health Level of the Elderly Living in Central Greece. *Mater Socio Medica* 2021;33:16.
5. Kadariya S, Gautam R, Aro AR. Physical Activity, Mental Health, and Wellbeing among Older Adults in South and Southeast Asia: A Scoping Review. *Biomed Res Int* 2019;2019:6752182.
6. Mehra A, Rani S, Sahoo S, Parveen S, Singh AP, Chakrabarti S, Grover S. A crisis for elderly with mental disorders: Relapse of symptoms due to heightened anxiety due to COVID-19. *Asian J Psychiatr* 2020;51:102114.
7. Centers of Disease Control and Prevention. Depression is Not a Normal Part of Growing Older. Washington: CDC;2014.
8. National Institute on Aging. National Institute on Aging. 2021 [cited 2024 Jun 19]. Depression and Older Adults. Available from: <https://www.nia.nih.gov/health/mental-and-emotional-health/depression-and-older-adults>
9. . Avasthi A GS. Clinical Practice Guidelines for Management of Depression in Elderly. *Indian J Psychiatry* 2018;60:S341-62.
10. Charzyńska E. Multidimensional approach toward spiritual coping: construction and validation of the Spiritual Coping Questionnaire (SCQ). *J Relig Health* 2015;54:1629-46.
11. Stanhope M, Lancaster J. Public health nursing: Population-centered health care in the community. Philadelphia, USA: Elsevier Mosby; 2019.
12. Widhiastini W, Permatasari NPI. The Indonesian Journal of Social Studies Social Construction & Tri Hita Karana Concept : Wellbeing. 2020;3:100-8.
13. Satya PANIP, Lase FJ. The Peaceful means of Tri Hita Karana in Balinese culture. 2018 [cited 2024 Jun 19]; Available from: <https://repository.unpar.ac.id/handle/123456789/repository.unpar.ac.id/handle/123456789/7299>
14. Sukarma IW. Tri Hita Karana: Theoretical Basic of Moral Hindu. *Int J Linguist Lit Cult* 2016;2:84.
15. Putro BD. Mental disorders (Buduh) in Bali as a cultural phenomenon: A study of perceptions and behavior of Balinese mental disorder treatment choices. Yogyakarta: Universitas Gadjah Mada; 2004.
16. Kusuma IGAT, Landra N, Widnyana W. Construction of Balinese local wisdom based on social model in the tourism sector in creating happiness to enhance community satisfaction. *Asia Pacific Manag Bus Appl* 2019;8:53-64.
17. Charan J, Biswas T. How to calculate sample size for different study designs in medical research? *Indian J Psychol Med* 2013;35:121-6.
18. Jané-Llopis E, Gabilondo A. Mental Health in Older People. Luxembourg: European Communities; 2008. (Consensus paper).
19. Koenig HG. Research on Religion, Spirituality, and Mental Health: A Review. *Can J Psychiatry* 2009;54:283-91.
20. Wiana IK. *Tri Hita Karana According to Hindu Concepts*. Surabaya: Paramita; 2007.

21. Cohen J. Statistical power analysis for the behavioral sciences. New York: Routledge; 1988.
22. Kang K, Gholizadeh L, Inglis SC, Han HR. Validation of the Korean Version of the MacNew Heart Disease Health-Related Quality of Life Questionnaire. *J Nurs Res* 2019;27:e11.
23. Ouchi Y, Rakugi H, Arai H, et al. Redefining the elderly as aged 75 years and older: Proposal from the Joint Committee of Japan Gerontological Society and the Japan Geriatrics Society. *Geriatr Gerontol Int* 2017;17:1045–7.
24. Lee SB, Oh JH, Park JH, et al. Differences in youngest-old, middle-old, and oldest-old patients who visit the emergency department. *Clin Exp Emerg Med* 2018;5:249–55.
25. Hao L, Xu X, Dupre ME, et al. Adequate access to healthcare and added life expectancy among older adults in China. *BMC Geriatr* 2020;20:1–15.
26. Juanamasta IG, Wati NMN, Widana AAGO. Covid-19: A bali-nese viewpoint. *Belitung Nurs J* 2020;6:143–4.
27. Kurniawidjaja M, Susilowati IH, Erwandi D, et al. Identification of Depression Among Elderly During COVID-19. *J Prim Care Community Health* 2022;13:2150131 9221085380.
28. Aunguroch Y, Juanamasta IG, Gunawan J. Experiences of Patients with Coronavirus in the COVID-19 Pandemic Era in Indonesia. *Asian J Public Opin Res* 2020;8:377–92.
29. Assariparambil AR, Noronha JA, Kamath A, et al. Depression among older adults: a systematic review of South Asian countries. *Psychogeriatrics* 2021;21:201–19.
30. Hartawan IGNBRM, Sastrawan IGG, Parastan RH, Ani LS. Depression, anxiety, and stress levels in Denpasar community during the pandemic of Corona Virus Disease 2019 (COVID-19). *Qanun Medika* 2021;5:103.
31. Suriastini NW, Suryani LK, Sikoki B, et al. Depression among older people in bali. *Asian J Gerontol Geriatr* 2021;16:22–9.
32. Meeks S, Carstensen LL, Tamsky BF, et al. Age differences in coping: Does less mean worse? *Int J Aging Hum Dev* 1989;28:127–40.
33. Ribeiro M dos S. Coping strategies used by the elderly regarding aging and death: an integrative review. *Rev Bras Geriatr Gerontol* 2017;20:869–77.
34. Nieto M, Romero D, Ros L, et al. Differences in Coping Strategies Between Young and Older Adults: The Role of Executive Functions. *Int J Aging Hum Dev* 2020;90:28–49.
35. Odat JM. *This Brain Had a Mouth*. Amherst, MA: University of Massachusetts Press; 2021.
36. Stanislawski K. The coping circumplex model: An integrative model of the structure of coping with stress. *Front Psychol* 2019;10:1–23.
37. Lestari RF. Coping Strategies for Students Seen from the Big Five Personalities. *Acta Psychol* 2021;3:29–37.
38. Galiana L, Tomás JM, Fernández I, Oliver A. Predicting Well-Being Among the Elderly: The Role of Coping Strategies. *Front Psychol* 2020;11:1–8.
39. Hoyt MA. Emotional Approach Coping in Older Adults as Predictor of Physical and Mental Health. *Psychol Aging* 2020;35(4).
40. Southard ME. Spirituality: The Missing Link for Holistic Health Care. *J Holist Nurs* 2020;38:4–7.
41. Juanamasta IG, Aunguroch Y, Gunawan J, Fisher ML. Postgraduate and undergraduate student nurses' well-being: A scoping review. *J Prof Nurs* 2022;40:57–65.
42. Dunn KS, Robinson-Lane SG. A Philosophical Analysis of Spiritual Coping. *Adv Nurs Sci* 2020;43:239–50.
43. Baldacchino D, Draper P. Spiritual coping strategies: A review of the nursing research literature. *J Adv Nurs* 2001;34:833–41.
44. Huang H, Rockwell J. Nature and the Spirit: Tri Hita Karana, Sacred Artistic Practices, and Musical Ecology in Bali. *EnviroLab Asia* 2019;3:1–27.
45. Nerawati NGAA, Sudarsana IK. The Role of Pasraman Kilat in Raising Sradha and Bhakti of Teenagers. *J Penelit Agama Hindu* 2020;4:74.
46. Surpi NK, Istriyanti NLA, Avalokitesvari NNAN. Resilience and Coping Strategy of Bhagavad-Gītā Readers on the COVID-19 Pandemic in Bali. *Kurukshetra Univ Res J* 2021;55:26–33.
47. Arisanti NMD, Suderana IW. Handling The Covid-19 Pandemic: Collaboration Of District Government With KnpI Gianyar In Gianyar District, Bali. *SP* 2020;15:87.
48. Egerer M, Lin B, Kingsley J, et al. Gardening can relieve human stress and boost nature connection during the COVID-19 pandemic. *Urban For Urban Green* 2022;68:127483.

Correspondence: I Wayan Suardana, Nursing Department, Polytechnic of Health Denpasar, Bali, Indonesia, Jl. Sanitasi, No 1 Denpasar, Bali, Indonesia.
Tel.: +62361710447
E-mail: suardanawayan@yahoo.com

Key words: depression; elderly; spiritual coping; tri hita karana

Contributions: IWS, AY, and RH, made significant contributions to the study through their involvement in study design, data collection and analysis, manuscript writing, and data analysis; KS, IKG, and IWM, also played vital roles by contributing to study design, manuscript writing, and conducting reference searches.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: this study was financially supported by Polytechnic of Health Denpasar, Bali, Indonesia

Clinical trials: not applicable.

Ethics approval and consent to participate: this study was reviewed by The Health Research Ethics Committee Faculty of Nursing Universitas Airlangga, with the ethical approval number 2303-KEPK. Informed consent was distributed to the participants before completing the main questionnaires. The informed consent has been applied approaching the principle of beneficial, no harm, confidential, justice, as well as voluntary participants. The provided information should be agreed to by participants before data collection. The participant could stop the survey anytime during the data completion. The informed consent had been reported during the Ethics Approval and Consent prior the study.

Conference presentation: part of this article was presented at the 4th International Nursing and Health Sciences Symposium, from 27th-28th of October 2023, Universitas Brawijaya, Malang, East Java, Indonesia.

Acknowledgment: the authors acknowledge the Director of Polytechnic of Health Denpasar, Bali, Indonesia for generously funding this study.

Received: 3 November 2023.
Accepted: 8 June 2024.

This work is licensed under a Creative Commons Attribution 4.0 License (by-nc 4.0).

©Copyright: the Author(s), 2024
Licensee PAGEPress, Italy
Healthcare in Low-resource Settings 2024; 12(s1):13025
doi:10.4081/hls.2024.13025

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.