

# The effectiveness of hyperoxygenation in preventing oxygen desaturation in intubated infants treated with endotracheal suctioning

Sholihatul Amaliya,<sup>1</sup> Yeni Rustina,<sup>2</sup> Defi Efendi<sup>2</sup>

<sup>1</sup>Nurse Profession Program, Faculty of Health Sciences, Universitas Brawijaya, Malang, East Java; <sup>2</sup>Faculty of Nursing, Universitas Indonesia, Depok, West Java, Indonesia

## Abstract

Endotracheal suctioning is an effective nursing intervention for intubated infants undergoing invasive mechanical ventilation. However, this intervention has the potential to cause side effects, such as hypoxemia, which typically requires hyperoxygenation. Despite the widespread use of hyperoxygenation in clinical practice, there are limited reports on its effectiveness in infants. This study aims to determine the effect of hyperoxygenation on oxygen saturation of intubated infants. The study procedures were carried out using a quasi-experimental method with a cross-over approach. In addition, the sample population comprised 22 intubated infants who were treated at the perinatology unit of Dr. Cipto Mangunkusumo Hospital in Jakarta from March to April 2018. The participants were then divided into 2 groups based on their condition, and oxygen saturation level was assessed. The intervention comprised endotracheal suctioning with and without hyperoxygenation, with a 3-hour washing period. Data collection was performed at different measurement intervals, namely before, during, and after suctioning, followed by statistical analysis. The paired-sample t-tests and Wilcoxon test showed that there was a significant difference in oxygen saturation at certain measurement intervals ( $p=0.006$ ;  $p=0.010$ ;  $p=0.001$ ;  $p=0.001$ ;  $p=0.001$ ). In addition, Friedman test also showed the presence of a significant difference in the reduction of oxygen saturation between the control and intervention groups ( $p=0.001$ ). In conclusion, hyperoxygenation had the potential to prevent the reduction of oxygen saturation during endotracheal suctioning in intubated infants. However, individual assessment of the need for suctioning and hyperoxygenation was necessary for a safe and effective procedure.

## Introduction

Respiratory insufficiency in infants is a prevalent challenge among newborns, which typically necessitates intensive care at the Neonatal Intensive Care Unit (NICU).<sup>1-3</sup> Several studies have shown that the majority of infants treated at the NICU often require invasive mechanical ventilation (MV) and the use of endotracheal tube (ETT) to create a connection with the ventilator.<sup>4</sup>

However, ETT causes mild inflammation in the airway mucosa and suppresses mucociliary mechanisms, leading to an impairment in the ability to mobilize and expectorate secretions. The use of ETT also impairs cough reflexes, causing increased mucus production and accumulation in the airways.<sup>5,6</sup> This accumulation has the potential to disrupt the ventilation of infants, thereby requiring frequent endotracheal suctioning to maintain the patency of the airway.<sup>7-9</sup> According to previous studies, endotracheal suctioning is a routine nursing intervention for intubated patients to drain mucus secretion. In addition, it is often performed in patients who cannot remove secretions to maintain airway patency and provide optimal ventilation.<sup>8</sup> Several reports have shown that the use of proper endotracheal suctioning technique provides various benefits, namely improving respiratory sounds and gas exchange, decreasing airway resistance and PIP (peak inspiratory pressure) of the ventilator, improving dynamic lung compliance, as well as enhancing oxygen saturation ( $SpO_2$ ) and arterial blood gas.<sup>4</sup>

In infants, endotracheal suctioning is often performed using ETT with a smaller diameter, leading to challenges during the procedure and increasing the risk of complications. ETT with an internal diameter of  $\leq 4$ mm has been reported to have the potential to cause an immediate reduction in dynamic lung compliance and expired tidal volume (TV). This shows that ineffective endotracheal suctioning can cause ETT obstruction, atelectasis, as well as ineffective ventilation and oxygenation, thereby necessitating reintubation.<sup>4</sup> Although suctioning is essential for improving ventilation and oxygenation, the procedure is associated with various complications and risks. These include bleeding of mucosal airways, infection, atelectasis, hypoxia, hypoxemia, cardiovascular instability, and elevated intracranial pressure,<sup>5,10</sup> with hypoxia and hypoxemia being the most common and severe.<sup>5,9,11</sup> In addition, stress due to the procedure can influence the metabolic rate and increase oxygen demand,<sup>12</sup> showing the need for hyperoxygenation. According to previous studies, hyperoxygenation is the administration of oxygen fraction ( $FiO_2$ ) at a percentage greater than the baseline requirement before suctioning up to 100%.<sup>4,9</sup> Although this technique is widely used in clinical practice to avoid hypoxemia, there are limited reports on its effectiveness in intubated infants. Therefore, this study aims to determine the effectiveness of hyperoxygenation in endotracheal suctioning compared to suctioning without hyperoxygenation.

### Significance for public health

*Endotracheal suctioning is an essential nursing intervention for intubated infants treated with invasive mechanical ventilation. However, this procedure causes various side effects, including hypoxemia, and often requires hyperoxygenation. Despite the effectiveness of hyperoxygenation, there are limited reports on its use among infants. The study indicates a notable oxygen saturation variation among infants who underwent endotracheal suctioning with hyperoxygenation.*

## Materials and Methods

### Study Design

This was a quasi-experimental study with a cross-over approach, which was carried out to determine the immediate effects of hyperoxygenation in endotracheal suctioning compared to endotracheal suctioning without hyperoxygenation on oxygen saturation in 2 different groups of newborn infants. Each sample received 2 types of intervention, namely endotracheal suctioning with hyperoxygenation (for the intervention group) and endotracheal suctioning without hyperoxygenation (for the control group). In addition, this was a single-centered study, and trials were conducted based on standard protocols as well as good clinical practice and national ethical guidelines.

### Sample and setting

The sample population comprised 22 infants who were treated in Dr. Cipto Mangunkusumo Hospital, the National Referral Hospital located in Central Jakarta, Indonesia from March to April 2018. In addition, the participants were allocated based on their condition before suctioning procedure. Infants were eligible for admission when mechanically ventilated through ETT and had an indication for endotracheal suctioning. The indications included clinical history of decreased SpO<sub>2</sub> in previous suction, visible secretions in the ETT, audible secretions, changes in respiratory rate, and decreased SpO<sub>2</sub> immediately before suctioning. Exclusion criteria included ventilation using high-frequency oscillating (HFO) ventilation, the presence of conditions, requiring perfusion, such as hypotension, anemia, cold stress, congenital heart disease, and presence of skin lesions detected by oxygen saturation device based on blood saturation number, and chronological age of more than 28 days. The sample size estimation was performed based on a previous study by Walsh et. al (1987), which used 16 participants.<sup>13</sup> This study compared supplemental oxygenation or hyperoxygenation during tracheobronchial hygiene on the transcutaneous partial pressure of oxygen (TcPO<sub>2</sub>). The results showed that the standard deviation of change in TcPO<sub>2</sub> was 17.58 torr and the mean difference between groups considered significant (effect size) was 22.54 torr.<sup>13</sup> Based on these results, the calculated minimum sample size per group was 19, and this present report used a total of 22 infants. To minimize the between-subject variability, the participants used 2 types of ventilators with the same specifications. Suction procedure was carried out in the same position (supine) and was supported by nesting by the same nurse using the same standard intervention process.

### Intervention

The participants were allocated into 2 sequential enrollment schemes based on infants' condition before interventions (conditions that require suctioning), hence, the section was not carried out routinely.<sup>14</sup> The participants received 2 different interventions, namely endotracheal suctioning with and without hyperoxygenation. This showed when the participants had a significant decrease in SpO<sub>2</sub>  $\geq$  2 points (%) before suctioning procedure compared to the previous condition, hyperoxygenation was performed and the next suctioning was carried out without hyperoxygenation with infants being placed in group A (GA). In group B (GB), when there was a significant increase in SpO<sub>2</sub>  $\geq$  2 points (%), the reverse process was carried out. Endotracheal suctioning intervention was performed based on the hospital's protocol by 2 individuals (the researcher and research assistant) as recommended by professional organizations to ensure the safety of the participants.<sup>15,16</sup>

Intervention in group A comprised endotracheal suctioning procedure when infants showed indications. These indications included clinical history of decreased SpO<sub>2</sub> in previous suctioning, decreased SpO<sub>2</sub> before suctioning, visible secretions in the ETT, audible secretions, coarse and/or decreased breath sounds, and changes in respiratory rate and pattern. Before the first endotracheal suctioning was carried out, hyperoxygenation was performed by pressing the "suctioning support" button on the ventilator, leading to an additional 20% FiO<sub>2</sub> for 2 minutes. In this study, hyperoxygenation was performed 30 seconds before suctioning. Several reports showed that endotracheal suctioning must not exceed 10 seconds, with each suctioning comprising 3 probe insertions. Subsequently, the second suctioning was performed after infants' saturation returned to normal range after decreasing due to the first suctioning and the third suctioning. The procedures were performed in a supine position and the participants were supported by nesting. The use of nesting in premature infants facilitated relaxation and comfort to minimize pain.<sup>17</sup> Group B underwent the same intervention as group A, but without hyperoxygenation before endotracheal suctioning.

### Data collection and analysis

In this study, pre and post-suctioning oxygen saturation (SpO<sub>2</sub>) was obtained in 6 different measurements, namely 60 seconds before suctioning (T-1), immediately after hyperoxygenation (T0), after the first suctioning repetition (T1), after the second suctioning repetition (T2), after the third repetition (T3), and 3 minutes after suctioning (T4). Moreover, the socio-demographic characteristics, gestational age, and birth weight of the participants were also measured. Statistical analysis was performed using SPSS version 16, and the data normality was assessed using Shapiro Wilks test. The demographic data were evaluated by measuring distribution, frequency, and the central tendency for birth weight and gestational age. The differences across groups on infants' oxygen saturation (SpO<sub>2</sub>) were analyzed using paired-sample t-tests (normal data), and Wilcoxon (abnormal data). For data within groups, analysis was carried out using Friedman test, followed by Post-Hoc Wilcoxon test. Data were reported as mean difference (95% CI) and p-values < 0.05 were considered significantly different.

### Ethical considerations

This study was conducted after obtaining approval from the Ethics Committee of the Faculty of Nursing, University of Indonesia (No.139/UN2.F12.D/HKP.02.04/2018), hospital's approval where the procedures were conducted, and written consent from parents or guardians.

## Results and Discussion

### Participants' characteristics

The characteristics of the participants consisted of gestational age and birth weight, which were described in Table 1 and Figure 1. The results showed that the mean gestational age was 33.23 weeks and the majority of infants (36.4%) were late preterm (gestational age 34-36 weeks), with low birth weight (50%).

The majority of the participants were preterm infants, particularly late preterm, with gestational age ranging from 27 to 36 weeks and low birth weight of 1000 to 2499 grams. Preterm infants with low birth weight had a great risk for respiratory complications due to the late maturity of the respiratory system.<sup>18</sup>

Some of the problems affecting the respiratory system of

preterm infants and causing breathing difficulty were surfactant deficiency that led to respiratory distress syndrome (RDS), immaturity of respiratory control central causing apnea, and increased risk of airway obstruction due to smaller respiratory airways.

The most common respiratory complication in preterm infants with low birth weight was RDS caused by lung immaturity and lack of surfactant, which affected the poorly compliant lungs with poor gas exchange. This respiratory complication could cause respiratory failure that required invasive or non-invasive MV.

### The differences in oxygen saturation (SpO<sub>2</sub>) across groups

The results of the paired t-test in Table 2 and Figure 2 showed that there were significant differences in oxygen saturation at several measurements between the intervention and control groups. These differences were observed before the intervention (hyperoxygenation), in the first, second, and third suctioning repetition, and 3 minutes after mucus suctioning. Based on the results, the mean oxygen saturation of the intervention group was greater compared to the controls at T0, T1, T2, T3, and T4.

Figure 2 showed that the intervention group experienced an increase in oxygen saturation after hyperoxygenation. After the first, second, and third suctioning repetitions, oxygen saturation decreased but did not become a desaturation period where the percentage was less than 85%. These results were inconsistent with the control group, which experienced a decrease in oxygen saturation to desaturation after the third suctioning.

Oxygen saturation during endotracheal suctioning in this study was measured at 6 different periods, namely 60 seconds before suctioning, immediately after hyperoxygenation, after the first second, and third repetition, and three minutes after suctioning. The result showed that during endotracheal suctioning at first, second,




Figure 1. Recruitment diagram and respondent allocation.

Table 1. Respondents' characteristics based on the infants' gestational age and birth weight (n=22).

Variable	Mean	SD	Range (Min-Max)	95% CI
Gestational age (weeks)	33.23	3.380	27-38	31.73-34.73
Birth weight (grams)	1952.95	731.485	740-3520	1628.63-2277.28
Variable	f	%		
Gestational age (weeks)	Extremely Premature	1	4.5	
	Very Premature	6	27.3	
	Moderately Premature	2	9.1	
	Late Preterm	8	36.4	
	Aterm	5	22.7	
Birth weight (grams)	Extremely Low Birth Weight	2	9.1	
	Very Low Birth Weigh	4	18.2	
	Low Birth Weight	11	50	
	Normal birth weight	5	23	

Table 2. The differences in oxygen saturation (SpO<sub>2</sub>) across groups (n=22).

Variable	Suction with hyperoxygenation (mean+ sd)	95% CI	Suction without hyperoxygenation (mean+ sd)	95% CI	p value
T-1	87.32±7.52	83.98-90.65	91.91±3.82	90.22-93.60	0.006*
T0	93.68± 5.43	91.28-96.09	91.91±3.82	85.80-89.84	0.147
T1	91.68 (77-100)	89.32-94.05	87.82 (76-95)	85.80-89.94	0.014**
T2	91.82±5.82	89.24-94.40	85.91±4.59	83.88-87.94	0.001*
T3	90±5.54	87.54-92.45	83.77±5.30	81.42-86.12	0.001*
T4	96 (87-99)	93.61-96.12	90 (85-96)	88.81-91.46	0.001**

\* p < 0.05, there is a significant difference used paired t test. \*\* p < 0.05, there is a significant difference used Wilcoxon.


and third suctioning repetition, there was higher oxygen saturation in the intervention group compared to the controls. Based on the results of the paired t-test and Wilcoxon test, there was a significant difference in oxygen saturation after the first, second, and third suctioning repetitions between the intervention and control groups. This was because the intervention group was given hyperoxygenation by giving an additional 20% of FiO<sub>2</sub> for 2 minutes which could prevent and minimize the occurrence of hypoxemia in infants during suctioning.<sup>4,7,13</sup> Hyperoxygenation by adding FiO<sub>2</sub> of 10-20% was adequate in preventing hypoxemia when suctioning through ETT.<sup>7</sup>

### The differences of oxygen saturation (SpO<sub>2</sub>) within groups

Differences in oxygen saturation within the intervention and control groups were assessed using the Friedman test and continued with the post hoc Wilcoxon test (Table 3). The results of the intervention group showed that there were significant differences in several pairs of oxygen saturation measurement points, namely (T-1) - T<sub>0</sub>, T<sub>0</sub> - T<sub>1</sub>, T<sub>0</sub> - T<sub>2</sub>, and T<sub>0</sub> - T<sub>3</sub> but T<sub>0</sub> - T<sub>4</sub> did not show significant variation. This was different in the control group where there were differences in oxygen saturation with all measurement points (T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>, T<sub>4</sub>).

This study showed differences in SpO<sub>2</sub> before intervention after hyperoxygenation, showing that the administration of hyperoxygenation through the addition of 20% could increase SpO<sub>2</sub> in infants before suctioning. SpO<sub>2</sub> after hyperoxygenation was significantly different from the value in the first, second, and third suctioning repetitions in both groups. However, the decrease in SpO<sub>2</sub> in the control group was greater compared to the intervention group. This showed that endotracheal suctioning procedure was able to significantly reduce SpO<sub>2</sub> despite hyperoxygenation. The decrease in SpO<sub>2</sub> during suctioning procedure occurred due to the removal of air from the airways.<sup>5</sup> The administration of negative pressure during suctioning could also cause a collapse of distal airways and alveoli despite the use of appropriate pressure and technique, leading to a reduction of SpO<sub>2</sub>.<sup>19</sup> The use of the open suctioning method in this study comprised the temporary disconnection of MV from infants' ETT, thereby interrupting airflow from MV, and this could serve as another contributing factor.<sup>20</sup> Other studies comparing open and close suctioning in infants who underwent endotracheal suctioning showed that the participants were able to sustain SpO<sub>2</sub> during the procedure with no significant difference. This was caused by the participants' ability to perform spontaneous pulmonary ventilation, leading to an increase in RR immediately after ETT suctioning as a compensation mechanism to sustain their SpO<sub>2</sub>.<sup>21</sup> The result also showed that in the intervention group, there was no significant difference between SpO<sub>2</sub> after

hyperoxygenation (T<sub>1</sub>) compared to the value at 3 minutes after mucus suctioning (T<sub>4</sub>). Within 3 minutes after suctioning, the SpO<sub>2</sub> level in the intervention group could return to the previous value. This proved that the administration of hyperoxygenation for 2 minutes could restore the value of SpO<sub>2</sub> to the previous level and within the normal range (88%-93%). Different results were obtained in the control group where SpO<sub>2</sub> at T<sub>1</sub> significantly varied from T<sub>4</sub>. The results showed that within 3 minutes, SpO<sub>2</sub> had not been able to return to the previous value and also had experienced desaturation (SpO<sub>2</sub> less than 88%). These results were also consistent with previous studies where a group of preterm infants who received hyperoxygenation before suctioning experienced fewer episodes of hypoxemia compared to those who received suctioning without hyperoxygenation and needed less time to reach normal SpO<sub>2</sub>.<sup>22</sup> Exposure to non-physiological oxygen levels (hypoxia or hyperoxia) could cause ROP, and continuous or intermittent hypoxia typically reduced myelination and induced angiogenesis in preterm animals.<sup>23</sup>



**Figure 2.** The changes of oxygen saturation in endotracheal suctioning at control and intervention group.

**Table 3.** The differences of oxygen saturation (SpO<sub>2</sub>) within groups (n=22).

Variable	Suction with hyperoxygenation (mean+ sd)	Suction without hyperoxygenation (mean+ sd)
T-1	87.32±7.52	91.91±3.82
T0	95 (79-100)*	91.91±3.82
T1	91.86±5.41*	87.82±4.56*
T2	93.50 (76-99)*	85.91±4.59*
T3	91 (76-96)*	83.77±5.30*
T4	96 (87-99)	90.14±2.98*
p value	0.001	0.001

\*Significant difference at T<sub>0</sub> with some measurement points

## The comparison in oxygen saturation (SpO<sub>2</sub>) reduction across groups

In this study, the reduction in oxygen saturation (SpO<sub>2</sub>) was calculated as the difference between oxygen saturation after hyperoxygenation and the lowest SpO<sub>2</sub> recorded during endotracheal suctioning procedure. The result in Table 4 showed that there was a significant difference in SpO<sub>2</sub> reduction in the intervention group compared to the control group where the decrease in oxygen saturation in the control group was greater.

The decrease in SpO<sub>2</sub> in the control group was higher compared to the intervention group. This was because, in the intervention group, hyperoxygenation was performed before suctioning. Hyperoxygenation was performed on infants by adding 10-20% FiO<sub>2</sub> above the previous oxygen percentage. A 20% increase in FiO<sub>2</sub> before mucus suctioning was as effective as giving 100% oxygen.<sup>4</sup> Other studies also suggested hyperoxygenation in newborns by increasing FiO<sub>2</sub> by 10% before mucus suctioning.<sup>22</sup> However, the administration of hyperoxygenation must be considered carefully to prevent complications. Some experts stated that the procedure must not be performed routinely, but only when the neonate had an indication, namely a clinical history of decreased SpO<sub>2</sub> during previous suctioning and decreased SpO<sub>2</sub> due to accumulation of secretions before suctioning.<sup>4,22</sup>

## The correlation of oxygen saturation (SpO<sub>2</sub>) reduction and respondents' characteristics

The correlation between decreased oxygen saturation (SpO<sub>2</sub>) and the participants' characteristics was analyzed to control the confounding factors, such as birth weight and gestational age. The result of the Pearson correlation in Table 5 showed that there was no relationship between the decrease in oxygen saturation (SpO<sub>2</sub>) with birth weight and gestational age. Based on these results, the difference in the decrease in oxygen saturation that occurred in the intervention and control groups was caused by hyperoxygenation intervention.

The present study showed that infants' oxygen saturation during and after endotracheal suctioning across both groups was significantly different. In addition, there was a significant difference in the reduction of oxygen saturation across both groups, with the control group possessing a higher value. The implementation was based on infants' response to care and previous suctioning. Several studies showed that nurses must conduct individual assessments of

the need for suctioning and hyperoxygenation before performing endotracheal suctioning. When there are signs of infants requiring the procedure, namely visible secretions in the ETT, audible secretions, coarse and/or decreased breath sounds, changes in respiratory rate<sup>9</sup> and pattern and accompanied by a decrease in oxygen saturation below 85% or 90% or a history of decreased saturation on previous suctioning, it was necessary to perform hyperoxygenation before endotracheal suctioning.<sup>4</sup> In addition, nurses had the option to rely on their clinical judgment or experience to decide the frequency of repetitions and the appropriate adjustment of FiO<sub>2</sub>, with a maximum allowance of 3 times suctioning repetitions and a FiO<sub>2</sub> increase of up to 20%. Despite its potential, hyperoxygenation could also cause side effects, namely hyperoxemia, which was associated with serious long-term effects, such as Retinopathy Of Prematurity (ROP), and Chronic Lung Disease (CLD) due to oxygen free-radical damage<sup>9</sup>. Therefore, when nurses observed that suctioning with hyperoxygenation led to an increase in infants' oxygen saturation to 100%, this showed the possibility of reducing repetitions or decreasing FiO<sub>2</sub> addition during subsequent suctioning to mitigate the risk of hyperoxemia. According to previous reports, there was also a crucial need to consider suctioning, specifically for extremely preterm infants during the first days of life. This was because a retrospective study showed an inverse relationship between the incidence of IVH and the number of suctioning.<sup>6</sup>

## Conclusions

In conclusion, this study showed that an increase in FiO<sub>2</sub> of 20% from the baseline value before endotracheal suctioning could be sufficient, effective, and safe to prevent hypoxemia and quickly return infants' oxygen saturation to normal ranges. The results obtained could be used as evidence and the basis for nurses in carrying out hyperoxygenation before endotracheal suctioning. The limitation of this study was the failure to set a standard percentage of the definition of a "decrease in oxygen saturation" and the use of a reduction from previous conditions. Therefore, future studies were advised to set a standard for decreasing oxygen saturation with a certain number, such as <85% or <93%.<sup>24,25</sup> Randomization of participants was not carried out randomly and completely, because it was based on the patients' condition, hence, there was a

**Table 4.** The Differences in oxygen Saturation (SpO<sub>2</sub>) reduction across groups (n=22).

The decrease of SpO <sub>2</sub> in endotracheal suctioning	Mean	SD	95% CI	p value
Intervention control (hyperoxygenation)	4.27	3.18	2.86-5.68	0.001
Control group (With hyperoxygenation)	8.5	4.21	6.68-10.41	

Paired t test.

**Table 5.** The correlation of oxygen saturation (SpO<sub>2</sub>) reduction and respondents' characteristics.

	Gestational age	Birth weight
SpO <sub>2</sub> reduction	r = 0.232 p = 0.130	r = 0.050 p = 0.747

Pearson correlation

risk of bias. Although this study used a crossover approach, it was classified as a quasi-experiment. Moreover, the procedures did not monitor respiratory indicators, such as ventilation per minute, TV, and long-term survival of infants. Future studies were expected to observe oxygenation in vital organs, such as the brain during mucus suctioning procedure using NIRS. The desaturation period that occurred in mucus suctioning must also be monitored and compared in the two groups.

Correspondence: Sholihatul Amaliya, Faculty of Health Sciences, Universitas Brawijaya, Malang, Indonesia. Jl. Puncak Dieng, Kunci, Kalisongo, Kec. Dau, Malang, East Java, Indonesia, Postcode: 65151, Tel.: +62341569117, Fax: +62341564755, Email: liya.fk.psik@ub.ac.id

Key words: endotracheal suction; hyperoxygenation; intubated infant; oxygen saturation

Contributions: all authors participated in the acquisition, analysis, or interpretation of data. Concept and design by DE and SA. Drafting of the manuscript by SA and YR. Statistical analysis by SA. Administrative, technical, or material support by SA and YR. All authors have read and approved the final manuscript.

Conflict of interest: the author declares no potential conflict of interest.

Funding: the authors thank the Faculty of Health Sciences at Universitas Brawijaya for their generous support of this study.

Ethics approval: this study was conducted after obtaining approval from the Ethics Committee of the Faculty of Nursing, University of Indonesia (No.139/UN2.F12.D/HKP.02.04/2018), hospital's approval where the procedures were conducted, and written consent from parents or guardians.

Clinical trials: not applicable.

Conference presentation: part of this article was presented at the 4th International Nursing and Health Sciences Symposium, from 27th-28th of October 2023, Universitas Brawijaya, Malang, East Java, Indonesia.

Acknowledgment: the authors thank the hospitals for granting permission to conduct this study and the participants for their involvement. Special thanks are extended to Prof. Dr. Titin Andri Wihastuti, S.Kp., M.Kes. for their support during publication.

Received: 3 November 2023.

Accepted: 8 June 2024.

Early view: 10 September 2024.

This work is licensed under a Creative Commons Attribution 4.0 License (by-nc 4.0).

©Copyright: the Author(s), 2024

Licensee PAGEPress, Italy

Healthcare in Low-resource Settings 2024; 12(s1):13045

doi:10.4081/hls.2024.13045

*Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

## References

1. Reuter S, Moser C, Baack M. Respiratory distress in the newborn. *Pediatr Rev* 2014;35:417–28.
2. Oktaviani E, Rustina Y, Efendi D. Facilitated tucking effective to pain relief on the preterm infants in perinatal unit in Jakarta. *J Keperawatan Indones* 2018;21:9–16.
3. Babaei H, Dehghan M, Pirkashani LM. Study of Causes of neonatal mortality and its related factors in the neonatal intensive care unit of Imam Reza Hospital in Kermanshah, Iran during (2014-2016). *Int J Pediatr* 2018;6:7641–9.
4. Gonçalves RL, Tsuzuki LM, Carvalho MGS. Endotracheal suctioning in intubated newborns: An integrative literature review. *Rev Bras Ter Intensiva* 2015;27:284–92.
5. Kiraly JN, Tingai, G D, Mills FJ, Morley JC, Copnell B. Endotracheal suction. *Pediatr Res* 2008;26:527–9.
6. Bruschetini M, Zappettini S, Moja L, Calevo MG. Frequency of endotracheal suctioning for the prevention of respiratory morbidity in ventilated newborns. *Cochrane Database Syst Rev* 2015;2015.
7. Tavangar H, Javadi M, Sobhanian S, Jahromi FF. The effect of the duration of pre-oxygenation before endotracheal suction on hemodynamic symptoms. *Glob J Health Sci* 2016;9:127.
8. Kalyn A, Blatz S, Feuerstake S, et al. Closed suctioning of intubated neonates maintains better physiologic stability: A randomized trial. *J Perinatol* 2003;23:218–22.
9. Gardner DL, Shirland L. Evidence-based guideline for suctioning the intubated neonate and infant. *Neonatal Netw* 2009;28:281–302.
10. Trevisanuto D, Doglioni N, Zanardo V. The management of endotracheal tubes and nasal cannulae: The role of nurses. *Early Hum Dev* 2009;85:S85–7.
11. Tingay DG, Copnell B, Mills JF, et al. Effects of open endotracheal suction on lung volume in infants receiving HFOV. *Intensive Care Med* 2007;33:689–93.
12. Clifton-Koepfel R. Endotracheal tube suctioning in the newborn: a review of the literature. *Newborn Infant Nurs Rev* 2006;6:94–9.
13. Walsh CM, Bada H, Korones S, Carter M. Controlled supplemental Oxygenation during tracheobronchial hygiene.pdf. *Nurs Res* 1987;36:211–5.
14. Cavallin F, Res G, Monfredini C, et al. Time needed to intubate and suction a manikin prior to instituting positive pressure ventilation: a simulation trial. *Eur J Pediatr* 2021;180:247–52.
15. Altimier L, Brown B, Tedeschi L. NANN Guidelines for Neonatal Nursing Policies, Procedures, Competencies, and Clinical Pathways. Glenview: National Association of Neonatal Nurses; 2006. 2006 p.
16. Cone S, Pickler RH, Grap MJ, et al. Endotracheal Suctioning in Preterm Infants Using Four-Handed versus Routine Care. *JOGNN - J Obstet Gynecol Neonatal Nurs* 2013;42:92–104.
17. Efendi D, Sari D, Riyantini Y, et al. Pemberian Posisi (Positioning) Dan Nesting Pada Bayi Prematur: Evaluasi Implementasi Perawatan Di Neonatal Intensive Care Unit (Nicu). *J Keperawatan Indones* 2019;22:169–81.
18. Ricci SS. Essentials of Maternity, newborn, & women's health nursing [Internet]. Lippincott Williams & Wilkins; 2012. Available from: <https://books.google.co.id/books?id=NVcduwAACAAJ>.
19. Tan AM, Gomez JM, Mathews J, et al. Closed versus partially ventilated endotracheal suction in extremely preterm neonates: Physiologic consequences. *Intensive Crit Care Nurs*

- 2005;21:234–42.
- 20 Walsh BK, Hood K, Merritt G. Pediatric airway maintenance and clearance in the acute care setting: How to stay out of trouble. *Respir Care* 2011;56:1424–44.
- 21 Cardoso JM, Kusahara DM, Guinsburg R, Pedreira MLG. Randomized crossover trial of endotracheal tube suctioning systems use in newborns. *Nurs Crit Care* 2017;22:276–83.
- 22 Pritchard MA, Flenady V, Woodgate P. Systematic review of the role of pre-oxygenation for tracheal suctioning in ventilated newborn infants. *J Paediatr Child Health* 2003;39:163–5.
- 23 Morken TS, Nyman AKG, Sandvig I, et al. Brain development after neonatal intermittent hyperoxia-hypoxia in the rat studied by longitudinal mri and immunohistochemistry. *PLoS One* 2013;8:1–13.
- 24 Jain D, D’Ugard C, Bello J, et al. Hypoxemia episodes during day and night and their impact on oxygen saturation targeting in mechanically ventilated preterm infants. *Neonatology* 2017;113:69–74.
- 25 Castillo A, Sola A, Baquero H, et al. Pulse oxygen saturation levels and arterial oxygen tension values in newborns receiving oxygen therapy in the neonatal intensive care unit: Is 85% to 93% an acceptable range? *Pediatrics* 2008;121:882–9.

Non-commercial use only