

Frequency and seasonality of viable *Helicobacter pylori* in drinking water in Dhamar Governorate, Yemen

Dhary Alewy Almashhadany,¹ Sara Mohammed Mayas,² Jiyan Ali Omar,³ Thaera A. M. Muslat⁴

¹Department of Medical Laboratory Science, College of Science, Knowledge University, Erbil, Iraq; ²Microbiology Section, Department of Biology, Faculty of Applied Sciences, Thamar University, Yemen; ³Department of Medical Microbiology, College of Science, Knowledge University, Erbil, Iraq; ⁴Community Medicine Specialist, Independent Researcher, Baghdad, Iraq

Abstract

Helicobacter pylori is an important and common bacterial pathogen in humans. The accumulated evidence of the existence of *H. pylori* in water from different environmental sources suggests a water-borne transmission route. This study aimed to investigate the occurrence of *H. pylori* in different water sources used by human populations in Dhamar Governorate, Yemen. 250 samples were randomly collected from the municipal water supply network, wells, and springs. The samples were processed, plated onto modified campy-blood agar, and incubated under microaerobic conditions for 4-10 days. Bacterial identification was based on morphological properties and biochemical tests. Bacteriological analysis

showed that 9.6% and 13.2% of tap and surface water samples were contaminated with *H. pylori*, respectively. Despite a higher frequency in samples from rural areas, these were not significantly ($p=0.068$) more contaminated than the samples from urban areas. Regarding the seasonal variations of *H. pylori* detection, 85.71% of positive samples were detected in the late winter and spring seasons (February to May). To conclude, *H. pylori* transmission through water is likely to occur in Dhamar Governorate. Further prospective studies are highly recommended to provide further evidence and a clearer picture of *H. pylori* transmission.

Introduction

Helicobacter pylori has been recognized as a major pathogen in humans for nearly four decades. Although treating infected individuals and improving living standards in communities have reduced transmission of infection, *H. pylori* remains the most common human bacterial pathogen, infecting about half of the world's population (Hooi *et al.*, 2017). It is also associated with gastrointestinal diseases that range from gastritis to neoplastic diseases, such as mucosa-associated lymphoid tissue lymphomas and gastric cancer (Paradowska *et al.*, 2010; Abadi and Kusters, 2016).

A great deal has been learned about the epidemiology, infection, biology, pathogenesis, diagnosis, and treatment of *H. pylori*; however, major gaps in our knowledge remain to be filled. The precise mode of infection transmission remains unclear, even though many epidemiological studies have identified different potential risk factors (Kusters *et al.*, 2006; Chmiela and Kupcinkas, 2019). Although *H. pylori* infection occurs worldwide, its prevalence varies widely within and between countries. For instance, 70-90% of people in developing countries have *H. pylori*, while only 25-50% of those in developed countries are infected (Almashhadany and Mayass, 2018; Almashhadany *et al.*, 2022). Fortunately, *H. pylori* prevalence is decreasing as a result of improved sanitary conditions and treatment procedures (Azevedo *et al.*, 2007; Niknam *et al.*, 2014). Different bacterial pathogens, such as fast-growing atypical mycobacteria, *Burkholderia pseudomallei*, and *H. pylori*, might be transmitted directly by drinking water, especially in developing regions (Ashbolt, 2004). The main reservoir of *H. pylori* is humans, particularly the human stomach; however, environmental or animal reservoirs have also been investigated as sources of *H. pylori* infection (Almashhadany and Mayass, 2017). Food and water sources have been suggested as reservoirs outside the human gastrointestinal tract, and *H. pylori* or its DNA have been detected in these sources (Khalifa *et al.*, 2010; Momtaz *et al.*, 2014). It is worth mentioning that *H. pylori* can survive in water for prolonged periods (Enroth and Engstrand, 1995; Almashhadany, 2018, 2020).

Correspondence: Dhary Alewy Almashhadany, Department of Medical Laboratory Science, College of Science, Knowledge University, 44001 Erbil, Iraq.
E-mail: dhary.alewy@knu.edu.iq

Key words: Yemen, *Helicobacter pylori*, prevalence, water.

Contributions: all authors contributed equally.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: none.

Availability of data and materials: data and materials are available from the corresponding author upon request.

Acknowledgments: the authors would like to thank Baynoun Laboratories in Dhamar Governorate (Yemen) and Dr. Hassan Aljabry for providing access to the resources needed for this work.

Received: 9 September 2022.

Accepted: 29 August 2023.

Early view: 6 September 2023.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

©Copyright: the Author(s), 2023

Licensee PAGEPress, Italy

Italian Journal of Food Safety 2023; 12:10855

doi:10.4081/ijfs.2023.10855

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

The fact that *H. pylori* may be transmitted through water and contaminated food is posing a major problem in public health (Almashhadany, 2018; Almashhadany *et al.*, 2018; Vesga *et al.*, 2018). Epidemiological studies in Peru and other developing countries strongly support the transmission of *H. pylori* via water, as evidenced by its detection through polymerase chain reaction (PCR), suggesting that water can be an important vehicle for *H. pylori* infection (Hulten *et al.*, 1996; Boehnke *et al.*, 2018). The source of *H. pylori* in drinking water has been attributed to fecal contamination (Talaei *et al.*, 2015).

Various findings that support the hypothesis that *H. pylori* is a water-borne pathogen have been reported. It has been mentioned that individuals who consume raw vegetables are more likely to acquire *H. pylori* (Chen *et al.*, 2005; Yahaghi *et al.*, 2014). The association of the infection with the consumption of raw vegetables is additional indirect evidence of the presence of *H. pylori* in the water used for the irrigation of these vegetables (Abdel-Latif and Abouzied, 2016). Studies from developing countries with low socioeconomic status and poor management of drinking water suggest that environmental factors play important roles in *H. pylori* epidemiology (Aziz *et al.*, 2015). Indeed, an association between the prevalence of *H. pylori* in Peruvian children and the source of drinking water was shown more than three decades ago (Klein *et al.*, 1991).

In Yemen, *H. pylori* infection is primarily acquired in early childhood (Almashhadany and Mayass, 2017). Its transmission routes are debated; therefore, this work aimed to study the occurrence of *H. pylori* in drinking water in Dhamar Governorate (Yemen) and to determine its occurrence in drinking water for months during the study period.

Materials and Methods

Sampling sites and study design

Dhamar Governorate is located in the middle west of Yemen at 2400-2500 m above sea level and has a small main city. Tap water samples were collected from the governmental municipal water network in the city, while surface water and groundwater samples were taken from small natural springs/streams and wells in the villages around the city. These natural sources are used as drinking water for villagers. The samples (1 L each) were collected randomly between February and October 2021 to assess the frequency of detecting culturable *H. pylori* in different seasons.

Sample collection and processing

A total of 250 drinking water samples (136 tap water and 114 surface water) were collected from different places in Dhamar Governorate, Yemen. The samples were collected in sterile bottles and transported to the laboratory inside an icebox containing ice on the internal walls (Almashhadany and Mayass, 2017). Samples were filtered through a 42-mm-diameter 0.45- μ m-pore-sized nylon membrane. The membrane was transferred onto a sterile 140-mm Petri dish containing 25 mL of phosphate-buffered saline and the organisms were resuspended using a cell scraper to rub the surface of the membrane and elute the depositions off the surface. A volume of 10 mL was transferred into a sterile container and stored at 4°C until culture within half an hour (Whale *et al.*, 2003).

Detection of *Helicobacter pylori*

The stored volume (10 mL) was centrifuged at 4000 rpm for 30 minutes, and the upper 5 mL of the supernatant was discarded (Whale *et al.*, 2003). Then 20 μ L of the sample was spread onto modified campy-blood agar with Dent's supplement and incubated under microaerobic conditions (O₂ 5%, CO₂ 5%, H₂ 2%, N₂ 88%) for 4-10 days (Almashhadany and Mayass, 2017; Almashhadany *et al.*, 2022). The identification of *H. pylori* isolates started with colony morphology on the agar plates and was confirmed by biochemical tests according to a published protocol (Whale *et al.*, 2003). Suspected small, round, and translucent colonies of *H. pylori* were purified by subculture and then identified by the mentioned biochemical tests. The biochemical tests included catalase test, oxidase test, urease test, indole test, growth in 1% glycine, growth in 3.5% sodium chloride, triple sugar iron agar reactions, hydrogen sulfide production, cephalothin/nalidixic acid sensitivity/resistance, and hippurate hydrolysis test (Table 1).

Statistical analysis

Data were analyzed using SPSS software version 25 (IBM SPSS Inc, Chicago, IL, USA). Confidence intervals of prevalence were calculated using the "exact" Clopper-Pearson method at an a level of 0.05. The chi-square test was applied to test the difference between groups.

Results

Out of 250 samples of different types of drinking water, 28 (11.2%) gave a positive result for the isolation of *H. pylori*. This result includes 13/136 (9.6%) positive samples from tap water and 15/114 (13.2%) positive samples from surface water (Table 2). Statistical analysis using the chi-square test showed no significant differences between sources in terms of the occurrence of *H. pylori* among tap water and surface water samples ($p=0.370$, $\chi^2=0.80$). Additionally, it is estimated that a maximum of 15.78% of drinking water may be contaminated with *H. pylori*.

Table 1. Biochemical tests confirming the identity of *Helicobacter pylori*.

Biochemical tests	Results
Catalase	Positive
Oxidase	Positive
Urease	Positive
Indole	Negative
Growth in 1% glycine	Negative
Growth in 3.5% NaCl	Negative
H ₂ S production in TSI agar	Negative
TSI agar with lead acetate paper	Positive
Nalidixic acid	Resistance
Cephalothin	Sensitive
Hippurate hydrolysis	Negative

NaCl, salt; H₂S, hydrogen sulfide; TSI, triple sugar iron.

Occurrence of *Helicobacter pylori* in water according to collection sites

In rural drinking water, *H. pylori* was found at a higher frequency in comparison to drinking water from urban areas. However, that difference is not statistically significant ($p=0.068$, $\chi^2=3.33$). Table 3 shows the details of *H. pylori* prevalence in drinking water according to collection sites.

Isolation of *Helicobacter pylori* from water according to the type of water sources

Fourteen of 136 tap water samples (10.3%), 6 out of 80 groundwater samples from wells (7.5%), and 8 of 34 samples from spring water (23.5%) were found to be contaminated with *H. pylori* (Figure 1).

Monthly variations in the rate of *Helicobacter pylori* isolation

The rate of positive samples detected during the study period varied widely, with 85.71% of positive samples being detected in the late winter and spring (February to May). The highest rate of isolation from drinking water was in April (33.3%), with an observed decrease in the isolation rate in earlier and later months, and no isolation could be obtained from late summer to late autumn (August to October). There is a significant difference between seasons in terms of the isolation of *H. pylori* from water ($p<0.01$) (Figure 2).

Discussion

Humans are the principal source of *H. pylori* transmission, and control measures that can be applied to protect drinking water supplies include preventing contamination through human waste and providing adequate disinfection since *H. pylori* is sensitive to oxidizing disinfectants (Castillo-Rojas *et al.*, 2004). As the transmission of *H. pylori* through water has become increasingly suspected, the World Health Organization (WHO) now refers to this microorganism as a water contaminant (WHO, 2022). *H. pylori* acquisition is linked to the quality of the drinking water, particularly the consumption of untreated well water (Bellack *et al.*, 2006; Doyle,




Figure 1. Collected samples and *Helicobacter pylori*-positive samples. Percentages of collected samples from different sources along with the culture-positive samples.




Figure 2. Relationship between months and the prevalence of *Helicobacter pylori* in water during the period of the study. Error bars represent 95% confidence intervals.

Table 2. Isolation of *Helicobacter pylori* from drinking water types in Dhamar Governorate.

Water type	No. examined	Positive samples		95% confidence interval
		n	%	
Tap water	136	13	9.6	5.19-15.79
Surface and ground water	114	15	13.2	7.56-20.77
Total	250	28	11.2	7.57-15.78

Table 3. Isolation of *Helicobacter pylori* in drinking water according to collection sites.

Collection site	No. examined	Positive samples		95% confidence interval
		n	%	
Urban	130	10	7.7	3.75-13.69
Rural	120	18	15.0	9.14-22.67
Total	250	28	11.2	7.57-15.78

2012; Plonka *et al.*, 2014). The detection of *H. pylori* in water varies between studies that used different detection tests. The current findings are consistent with the study results previously obtained in Dhamar Governorate (Almashhadany and Mayass, 2017), where the total occurrence of *H. pylori* in water samples was 11.61%. However, we found fewer positive samples than those reported by Hegarty *et al.* (1999), who found actively respiring *H. pylori* in 60% of the surface water samples (n=62) and 65% of the shallow groundwater samples. The identity of *H. pylori* isolates was confirmed by 16S rRNA PCR (Hegarty *et al.*, 1999). On the other hand, our findings are higher than those found in similar studies from Iraq (n=471) (Al-Sulami *et al.*, 2012), Pakistan (n=50) (Khan *et al.*, 2012), and Iran (n=400) (Ranjbar *et al.*, 2016), where the detection rate ranged between 1.77% and 3.63% only.

Such variations are expected to be linked to differences in geographical locations, human activities near water sources, and detection methodologies. Several studies have been performed to assess the viability of *H. pylori* in water since it has been suggested that the coccoid form of *H. pylori* is responsible for transmission in the environment (Kayali *et al.*, 2018). Moreover, *H. pylori* could survive disinfection practices normally used in drinking water treatment in a viable but non-culturable form (Moreno *et al.*, 2007), which would allow the bacterium to go undetected by culture methods.

Regarding water types, our results are lower than those reported by Ahmed *et al.* (2007) in South India, who found that the prevalence of *H. pylori* infection among people who drank water from wells was 92% compared with 74.8% of those who drank tap water (p<0.001). *H. pylori* infection prevalence was found to be higher in people with a low clean water index (CWI) (88.2%) than in those with a higher CWI (33.3%) (p<0.001).

The seasonality of *H. pylori* is not well understood. Our seasonality results are not in agreement with the study by Ranjbar *et al.* (2016) performed in Iran, who found that the samples collected in the summer season had the highest prevalence of *H. pylori* (4.54%). A significant statistical difference was seen in the prevalence of *H. pylori* between the warm and cold seasons of the year (p<0.05). In another study carried out in Venezuela, the findings showed that *H. pylori* infection frequency was significantly higher during the rainy season (96%) than during the dry season (Domínguez-Bello *et al.*, 2002).

Conclusions

Water is an important vehicle for the transmission of numerous pathogens, including *H. pylori*. In Dhamar Governorate, transmission of *H. pylori* through water is very likely to occur, as viable cells of this pathogen have been recovered through traditional bacteriological analysis. Surface water was more frequently contaminated than municipal water supplies. Regarding the seasonality of *H. pylori*, the highest detection rates occurred during winter and spring, while no positive samples were detected during autumn. Further prospective studies are highly recommended to provide additional evidence and a clearer picture of *H. pylori* transmission.

References

- Abadi ATB, Kusters JG, 2016. Management of Helicobacter pylori infections. BMC Gastroenterol 16:94.
- Abdel-Latif MMM, Abouzied MM, 2016. Molecular mechanisms of natural honey against H. pylori infection via suppression of NF-κB and AP-1 activation in gastric epithelial cells. Arch Med Res 47:340-8.
- Ahmed KS, Khan AA, Ahmed I, Tiwari SK, Habeeb A, Ahi JD, Abid Z, Ahmed N, Habibullah CM, 2007. Impact of household hygiene and water source on the prevalence and transmission of Helicobacter pylori: a South Indian perspective. Singapore Med J 48:543-9.
- Almashhadany DA, 2018. Application of stool antigen test for monitoring Helicobacter pylori among human in Erbil Governorate, Kurdistan Region, Iraq. Int J Pharm Pharm Sci 10:49-53.
- Almashhadany DA, 2020. Epidemiology of Helicobacter pylori among human at Erbil Governorate/Kurdistan Region/Iraq. World J Pharm Pharm Sci 9:435-47.
- Almashhadany DA, Ismael LQ, Zaki AM, 2018. Seroprevalence of Helicobacter pylori among human in Erbil governorate, Kurdistan region, Iraq. Res J Life Sci Bioinform Pharm Chem Sci 4:268-80.
- Almashhadany DA, Mayass SM, 2017. Incidence of Helicobacter pylori in food and water in Dhamar governorate/Yemen. Int J Curr Res 9:45320-6.
- Almashhadany DA, Mayass SM, 2018. Prevalence of Helicobacter pylori in human in Dhamar Governorate/Yemen. J Med Pharm Sci 2. DOI: 10.26389/AJSRPS101217.
- Almashhadany DA, Mayas SM, Ali NL, 2022. Isolation and identification of Helicobacter pylori from raw chicken meat in Dhamar Governorate, Yemen. Ital J Food Saf 11:10220.
- Almashhadany DA, Zefenkey ZF, Zaki AM, 2022. Dental risk factors associated with oral Helicobacter pylori infection: a cross-sectional study based on saliva antigen test. J Infect Dev Ctries 16:516-21.
- Al-Sulami AA, Al-Edani TAA, Al-Abdula AA, 2012. Culture method and PCR for the detection of Helicobacter pylori in drinking water in Basrah Governorate Iraq. Gastroenterol Res Pract 2012:245167.
- Ashbolt NJ, 2004. Microbial contamination of drinking water and disease outcomes in developing regions. Toxicology 198:229-38.
- Azevedo NF, Guimaraes N, Figueiredo C, Keevil CW, Vieira MJ, 2007. A new model for the transmission of Helicobacter pylori: role of environmental reservoirs as gene pools to increase strain diversity. Crit Rev Microbiol 33:157-69.
- Aziz RK, Khalifa MM, Sharaf RR, 2015. Contaminated water as a source of Helicobacter pylori infection: a review. J Adv Res 6:539-47.
- Bellack NR, Koehoorn MW, MacNab YC, Morshed MG, 2006. A conceptual model of water's role as a reservoir in Helicobacter pylori transmission: a review of the evidence. Epidemiol Infect 134:439-49.
- Boehnke KF, Brewster RK, Sánchez BN, Valdivieso M, Bussalleu A, Guevara M, Saenz CG, Alva SO, Gil E, Xi C, 2018. An assessment of drinking water contamination with Helicobacter pylori in Lima, Peru. Helicobacter 23:e12462.
- Castillo-Rojas G, Mazari-Hiriart M, López-Vidal Y, 2004. Helicobacter pylori: focus on CagA and VacA major virulence factors. Salud Publica Mex 46:538-48.
- Chen MC, Hu CT, Wang LY, Lin HH, Sheu BS, Kao AW, Cheng HC, Hunag SF, Chen TW, Lu CC, Wu JJ, 2005. The efficacy of Helicobacter pylori eradication related to CYP2C19 metabolism. Aliment Pharmacol Ther 22:274-5.
- Chmiela M, Kupcinkas J, 2019. Review: pathogenesis of Helicobacter pylori infection. Helicobacter 24:e12638.
- Domínguez-Bello M, Beker B, Guelrud M, 2002. Short report:

- socioeconomic and seasonal variations of *Helicobacter pylori* infection in patients in Venezuela. *Am J Trop Med Hyg* 66:49-51.
- Doyle ME, 2012. *Helicobacter* spp. Food or Waterborne Pathogens? Available from: https://fri.wisc.edu/files/Briefs_File/FRI_Brief_Helicobacter_Feb2012.pdf.
- Enroth H, Engstrand L, 1995. Immunomagnetic separation and PCR for detection of *Helicobacter pylori* in water and stool specimens. *J Clin Microbiol* 33:2162-5.
- Hegarty JP, Dowd MT, Baker KH, 1999. Occurrence of *Helicobacter pylori* in surface water in the United States. *J Appl Microbiol* 87:697-701.
- Hooi JKY, Lai WY, Ng WK, Suen MMY, Underwood FE, Tanyingoh D, Malfertheiner P, Graham DY, Wong VWS, Wu JCY, Chan FKL, Sung JJY, Kaplan GG, Ng SC, 2017. Global prevalence of *Helicobacter pylori* infection: systematic review and meta-analysis. *Gastroenterology* 153:420-9.
- Hulten K, Han SW, Enroth H, Klein PD, Opekun AR, Gilman RH, Evans DG, Engstrand L, Graham DY, El-Zaatari FAK, 1996. *Helicobacter pylori* in the drinking water in Peru. *Gastroenterology* 110:1031-5.
- Kayali S, Manfredi M, Gaiani F, Bianchi L, Bizzarri B, Leandro G, Mario F Di, De'angelis GL, 2018. *Helicobacter pylori*, transmission routes and recurrence of infection: state of the art. *Acta Biomed* 89:72-6.
- Khalifa MM, Sharaf RR, Aziz RK, 2010. *Helicobacter pylori*: a poor man's gut pathogen? *Gut Pathog* 2:2.
- Khan A, Farooqui A, Kazmi SU, 2012. Presence of *Helicobacter pylori* in drinking water of Karachi, Pakistan. *J Infect Dev Ctries* 6:251-5.
- Klein PD, Graham DY, Gaillour A, Opekun AR, Smith EO, 1991. Water source as risk factor for *Helicobacter pylori* infection in Peruvian children. Gastrointestinal physiology working group. *Lancet* 337:1503-6.
- Kusters JG, Vliet AHM Van, Kuipers EJ, 2006. Pathogenesis of *Helicobacter pylori* infection. *Clin Microbiol Rev* 19:449-90.
- Momtaz H, Dabiri H, Souod N, Gholami M, 2014. Study of *Helicobacter pylori* genotype status in cows, sheep, goats and human beings. *BMC Gastroenterol* 14:61.
- Moreno YP, JL A, González AM, 2007. Survival and viability of *Helicobacter pylori* after inoculation into chlorinated drinking water. *Water Res* 41:3490-6.
- Niknam R, Seddigh M, Fattahi MR, Dehghanian A, Mahmoudi L, 2014. Prevalence of *Helicobacter pylori* in patients with dyspepsia. *Jundishapur J Microbiol* 7:e12676.
- Paradowska A, Sieja A, Braksator M, 2010. *Helicobacter pylori* infection and oral cavity. *Gastroenterologia Polska* 17:423-6.
- Plonka M, Targosz A, Brzozowski T, 2014. Can drinking water serve as a potential reservoir of *Helicobacter pylori*? Evidence for water contamination by *Helicobacter pylori*. In: Roesler BM. *Trends Helicobacter pylori Infection*. London, UK: IntechOpen.
- Ranjbar R, Khamesipour F, Jonaidi-Jafari N, Rahimi E, 2016. *Helicobacter pylori* isolated from Iranian drinking water: *vacA*, *cagA*, *iceA*, *oipA* and *babA2* genotype status and antimicrobial resistance properties. *FEBS Open Bio* 6:433-41.
- Talaei R, Souod N, Momtaz H, Dabiri H, 2015. Milk of livestock as a possible transmission route of *Helicobacter pylori* infection. *Gastroenterol Hepatol Bed Bench* 8:30:S30-6.
- Vesga FJ, Moreno Y, Ferrús MA, Campos C, Trespalacios AA, 2018. Detection of *Helicobacter pylori* in drinking water treatment plants in Bogotá, Colombia, using cultural and molecular techniques. *Int J Hyg Environ Health* 221:595-601.
- Whale A, Said B, Owen B, Watson C, Drobniewski F, Nichols G, Lee J V, Weldon L, Edwards R, Lai S, 2003. Further studies on the Incidence of Mycobacterium avium Complex (MAC) in drinking water supplies (including the detection of *Helicobacter pylori* in water and biofilm samples). Available from: https://dwi-content.s3.eu-west-2.amazonaws.com/wp-content/uploads/2020/10/27110801/DWI70-2-146_mac1.pdf.
- WHO, 2022. Drinking-water: key facts. Available from: <https://www.who.int/news-room/fact-sheets/detail/drinking-water>.
- Yahaghi E, Khamesipour F, Mashayekhi F, Dehkordi FS, Sakhaei MH, Masoudimanesh M, Khameneie MK, 2014. *Helicobacter pylori* in vegetables and salads: genotyping and antimicrobial resistance properties. *Biomed Res Int* 2014:757941.