



Italian Journal of Food Safety

<https://www.pagepressjournals.org/index.php/ijfs/index>

eISSN 2239-7132

Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. The Early Access service lets users access peer-reviewed articles well before print/regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

The **Italian Journal of Food Safety** is, therefore, E-publishing PDF files of an early version of manuscripts that have undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination, and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear in a regular issue of the journal.

The E-publishing of this PDF file has been approved by the authors.

Please cite this article as:

Di Giacomo L, Napoleoni M, Angellotti A, et al. **Monophasic variant of *Salmonella* Typhimurium 4,[5],12:i:- outbreak: an investigation by the Competent Authority.** *Ital J Food Saf* doi:10.4081/ijfs.2024.12522

Submitted: 28-03-2024

Accepted: 20-05-2024

 © the Author(s), 2024
Licensee PAGEPress, Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

**Monophasic variant of *Salmonella* Typhimurium 4,[5],12:i:- outbreak:
an investigation by the Competent Authority**

Loredana Di Giacomo,^{1*} Maira Napoleoni,^{2*} Antonio Angellotti,¹ Ezio Ferretti,¹ Valentina Gentili,¹
Marta Grifi,¹ Francesco Livini,¹ Monika Tardella,¹ Vanessa Travanti,¹ Simonetta Ruggeri¹

¹Animal Based Food Hygiene Service (ABFHS), Department of Prevention, Local Health Unit, Fermo; ²Regional Reference Center for Enteric Pathogens for the Marche Region, Istituto Zooprofilattico Sperimentale dell'Umbria e delle Marche "Togo Rosati", Tolentino, Italy

*These authors contributed equally.

Correspondence: Loredana Di Giacomo, Animal Based Food Hygiene Service (ABFHS), Department of Prevention, Local Health Unit, Fermo, Italy.
Tel.: +(39) 3393070198.
E-mail: loredana.digiaco@sanita.marche.it

Key words: official control, foodborne disease outbreaks, monophasic variant of *Salmonella* Typhimurium, porchetta.

Contributions: all the authors made a substantial intellectual contribution, read and approved the final version of the manuscript, and agreed to be accountable for all aspects of the work.

Conflict of interest: the authors declare no potential conflict of interest.

Ethics approval and consent to participate: not applicable.

Funding: none.

Availability of data and materials: data and materials are available from the corresponding author upon request.

Conference presentation: this paper was presented at the XXXII National Conference of the Italian Association of Veterinary Food Hygienists (AIVI), September 13-15, 2023, Maierato, Italy.

Abstract

Foodborne diseases represent a significant public health issue, regarding both collective health and the economy, with implications for healthcare costs and agribusiness. This paper shows the description and results of the investigation of a *Salmonella enterica* subsp. *enterica* foodborne outbreak that occurred in the Marche Region (Italy) in 2022, linked to the consumption of a roasted pork product (*porchetta*).

As part of the outbreak investigations discussed in this article, molecular analysis to evaluate the genomic correlation between clinical, food, and environmental origin strains was carried out. All *Salmonella* strains of different origins were serotyped as monophasic variants of *Salmonella* Typhimurium with an antibiotic-resistance pattern and an allelic profile such as to define the "cluster strain" allowing the correlation between clinical and food/environmental strains, definitively confirmed by whole genome sequencing analysis.

Following the laboratory evidence, corrective measures at the *porchetta* processing plant and at the retail stores involved were carried out by the Local Animal Based Food Hygiene Service.

The results of this study show that effective intervention is only possible if efficient data exchange, standardized procedures, and staff training are guaranteed. The latter aspect also concerns the food business operator, who must take appropriate measures to minimize the risk.

Introduction

Foodborne diseases are a serious public health concern and still one of the most important causes of economic losses and social costs (Hoffmann and Ahn, 2021).

In many cases, outbreaks are not recognized, reported, and/or epidemiologically investigated. In addition, the food source or the responsible pathogen remains unknown, so traceback procedures that could quickly stop the consumption of the contaminated product are not activated. This means that many outbreaks go undetected, not allowing for prevention and control measures. According to the European Union's One Health 2022 Zoonoses Report (EFSA and ECDC, 2023), in 70 (40%) outbreaks out of 175 total reported in Italy in 2022, the causative pathogen remained unknown, a figure in line with the European situation (46.1%). As far as the Italian situation is concerned, salmonellosis remains the most frequently reported zoonosis in humans, followed by campylobacteriosis, in contrast to the European trend.

In 2022, Italy notified the European Food Safety Authority of a total of 175 foodborne epidemic outbreaks involving 1604 human cases and resulting in 303 hospitalizations and 15 deaths.

Salmonella was the pathogen responsible for most of the epidemic outbreaks (52 out of 175) and also caused a high number of human cases (212 out of 1604). In 12 outbreaks (37.5 %) out of a total of 32 (18.3%), the food source involved was meat and meat products, mainly of pork origin.

At the European Union level, the European Community established procedures for the surveillance and collection of information on zoonoses as early as 1999 with Directive 2003/99/EC (European Parliament and Council of the European Union, 2003), adopted in Italy by Legislative Decree No. 191/2006, which provides for the obligation to investigate foodborne outbreaks [Italian Republic, 2006 - Article 7(2)].

As a result, an epidemiological surveillance network with updated infectious agents for surveillance was created, and a new information system for reporting infectious diseases called "PREMAL" was set up in 2017 with the decree of March 3 (Italian Republic, 2017). The Marche Region has also actively addressed the management of foodborne disease outbreaks and included it in the strategic macro-objective M06-9 of the 2020-2025 prevention plan: "ensure the identification of the food responsible for a foodborne outbreak so that the alert system can be activated to withdraw/recall the dangerous or potentially dangerous food from the market" (Marche Region, 2021).

The predefined program PP09, "Environment, Climate and Health", defines measures on various aspects: i) legal acts for the establishment of the regional reference laboratory; ii) the development and implementation of a guide for foodborne diseases; iii) the publication of information material for

citizens; iv) the training/information of official control officers; v) food business operators (FBOs) and primary production; vi) health personnel and others involved in notification.

In this article, we present the description and the results of the investigation of a foodborne outbreak of *Salmonella enterica* subsp. *enterica* that occurred in the Marche Region (Italy) in 2022, as part of several foodborne illnesses linked to the consumption of a roasted pork product (*porchetta*).

Materials and Methods

Between July and September 2022, several notifications for suspected food-related infections were shared by the Hygiene and Public Health Service (HPHS) in Fermo and neighboring provinces with the Animal Based Food Hygiene Service (ABFHS, hereafter Service) of the Fermo Local Health Unit (LHU). All the cases had in common enteric symptoms and feces confirmed positive for *Salmonella*; some of them also declared the consumption of roasted pork product (*porchetta*) bought and/or consumed in different places and times but produced at the same food-processing plant located in the Fermo area (site A) and recognized under EC Regulation No. 853/2004 (European Parliament and Council of the European Union, 2004a).

ABFHS inspected site A and also the two retail stores (sites B and C) recognized under EC Regulation No. 852/2004 (European Parliament and Council of the European Union 2004b). A total of 3 samples of *porchetta* and 23 environmental samples (using the sponge swab method) were collected for *Salmonella* detection. As *porchetta* does not fall within the scope of Regulation (EC) No. 2073/2005 (Commission of the European Communities, 2005 - Annex 1, Chapter 1, paragraph 8), the sampling strategy was changed. Annex 7 - Table 20 of the Deliberation No. 1508/2016 in which the "guide values" for the matrices that do not fall within the scope of the European standard are defined, was used as the legal basis (Marche Region, 2016). In this particular case, *porchetta* was considered a ready-to-eat product.

From the combination of the provisions of the regional regulation and Legislative Decree no. 27/2021 (Italian Republic, 2021 - Annex 1, point 2), a legal single aliquot and a single sample unit of *porchetta* were collected for a single non-repeatable analysis. Positivity for *Salmonella* spp. was considered as a food safety criterion.

For surface sampling, 7.5×3.5 cm cellulose sponges were used, moistened with 10 mL buffered peptone water in the case of unsanitized surfaces or with neutralizing Dey-Engley broth in the case of sanitized surfaces. The choice of sponges rather than swabs was necessary to sample a larger area and to ensure more effective removal of bacterial cells. A single, non-repeatable analysis was performed on the surface samples to also ensure the operator's right of defense.

In addition to the sampling, an assessment of the hygienic conditions of the premises and equipment was carried out in all sites (A, B and C) and an assessment of the procedures implemented by the FBO based on hazard analysis critical control point (HACCP) principles was carried out at site A.

The environmental and food samples collected at sites A, B, and C were analyzed for *Salmonella* at the food safety laboratory of the *Istituto Zooprofilattico Sperimentale dell'Umbria e delle Marche* (IZSUM) according to the UNI EN ISO 6579-1:2020 standard (ISO, 2020).

Salmonella spp. strains isolated from *porchetta*, environmental swabs, and human clinical samples were serotyped at the Regional Reference Center for Enteric Pathogens for the Marche Region (CRRPE) of IZSUM to ISO/TR 6579-3:2014 (ISO, 2014), and subsequently confirmed by multiplex polymerase chain reaction (PCR) for *fljB* (flagellar antigen H:2) gene (Rahn *et al.*, 1992).

Further sub-characterization analysis of a monophasic variant of *Salmonella* Typhimurium (MVST) strains was performed. In particular, antimicrobial susceptibility testing was performed by disk diffusion method (Kirby-Bauer test) according to the guidelines of the Clinical and Laboratory Standards Institute (2023).

Molecular typing by multiple-locus variable-number tandem repeat analysis (MLVA) according to the European Center for Disease Prevention and Control standard laboratory operating procedure for MLVA of *Salmonella enterica* serotype Typhimurium (ECDC, 2011), was carried out by the Department of Infectious Diseases of *Istituto Superiore di Sanità* and the National Reference Center

for Salmonellosis of *Istituto Zooprofilattico Sperimentale delle Venezie* for human and for food and environmental samples, respectively.

Whole genome sequencing was performed on a representative selection of 30 strains, among the total collected clinical, food and environmental strains, based on the MLVA results and epidemiological information (for further details, see Napoleoni *et al.*, 2023).

Results

Positivity for *Salmonella* spp. was detected both from environmental swabs (4 out of 23 total) and from *porchetta* (1 out of 3 total) (Table 1). The positive swabs regarded unsanitized surfaces in contact with the roasted pork: transporting board for cooked *porchetta* (site A), teflon chopping board for supporting and cutting *porchetta* (site B), wooden chopping board for *porchetta* and *porchetta* knife (site C). *Porchetta* sample, which tested positive for *Salmonella*, was collected at site C.

Salmonella spp. strains isolated from food *porchetta* (n=1), environmental swabs (4 out of 5 total positives for *Salmonella* spp.), and human clinical cases involved (n=64) were serotyped and confirmed by multiplex PCR as MVST 4,[5],12:i:-.

57 clinical strains of MVST out of 64 showed resistance to the following antibiotics: ampicillin (A), chloramphenicol (C), streptomycin (S), sulfisoxazole (Su), gentamicin (Gm), trimethoprim (Tmp), and trimethoprim-sulfamethoxazole (Sxt) (ACSSu+Gm+Tmp+Sxt profile, *i.e.*, “cluster strain” antibiotic resistance type). The remaining 7 clinical strains showed different partial types of antibiotic resistance with respect to the type of cluster strain: ACSSu+Tmp+Sxt (n=3), ACSSu+Gm (n=1), ACSSu+Amc+Gm+Tmp+Sx (n=1), CSSu+Gm+Tmp+Sxt (n=1).

63 clinical strains of MVST out of 64 (57 with the cluster strain antibiotic resistance type and 7 with antimicrobial resistance types close to the cluster-strain type), presented 3-14-10-na-211 allelic profile while the remaining strain, a closely related allelic profile, *i.e.*, 3-13-10-na-211. Therefore, a total of 57 isolates from cases of human salmonellosis were identified as belonging to the same cluster of the MVST (antibiotic resistance type ACSSu+Gm+Tmp+Sxt and MLVA 3-14-10-na-211 or 3-13-10-na-211). Additionally, seven isolates from human cases were defined as being closely related to this cluster of the MVST.

All 5 food/environmental strains of the MVST (4 from environmental swabs and 1 from *porchetta*) had the antibiotic resistance type ACSSu+Gm+Tmp+Sxt and the MLVA profile 3-14-10-na-211, corresponding to those of the human cluster strains.

The core genome multilocus sequence typing (cgMLST) analysis showed that no significant allelic distance was highlighted between the clinical (n=24), food-related (n=1), and environmental (n=4) strains tested, except for one clinical strain that was shown to be 49 alleles distant from the main group and was therefore considered a different clone (for further details about the obtained results, see Napoleoni *et al.*, 2023).

An inspection conducted at the food-processing plant revealed structural, sanitation, and documentary deficiencies regarding the procedures for good manufacturing and hygiene practices and those based on HACCP principles. In particular, the full production package did not correctly identify critical control points (CCPs) in the *porchetta* processing steps by managing cooking and subsequent chilling as good manufacturing practices (GMPs). No structural and sanitation nonconformities were found at site B. The inspection at site C found a lack of hygiene and non-compliance with product storage procedures.

Following the results of the inspections and sampling, the service took the following actions:

i) *Porchetta* production was suspended according to Article 138 of EU Regulation No. 625/2017 (European Parliament and Council of the European Union, 2017); violations of Articles 4 and 5 of EC Regulation No. 852/2004 for hygiene and sanitation deficiencies and non-application of procedures resulted in the imposition of two administrative sanctions pursuant to Article 6, paragraphs 5 and 6, of Legislative Decree No. 193/2007 (European Parliament and Council of the European Union, 2004b; Italian Republic, 2007);

- ii) Site C was informed of a minor non-compliance rule with corresponding compliance deadlines in relation to structural deficiencies and the implementation of self-checking procedures, resulting in a measure under Article 138 of EU Regulation No. 625/2017 with compliance deadlines and the imposition of two administrative sanctions for non-compliance with official control requirements and non-application of the procedure under Article 6(5) and (7) of Legislative Decree No. 193/2007 (Italian Republic, 2007; European Parliament and Council of the European Union, 2017);
- iii) Pursuant to Article 347 of the Code of Criminal Procedure (Italian Republic, 1988), a criminal complaint was issued against the legal representative of the production plant for violation of Articles 444 "trafficking in substances intended for food and harmful to public health" and 452 "offenses against public health" of the Criminal Code (Codice Penale, 1930); these measures were ordered following the isolation of the same serotype in all samples and clinical cases;
- iv) in accordance with Article 50 of Regulation (EC) No 178/2002 (European Parliament and Council of the European Union, 2002), an information notification was sent by the competent authority to the Rapid Alert System for Food and Feed, followed by a validation without triggering an alert, as the product was no longer on the market;
- v) Case management report to the Veterinary and Food Safety Functional Unit of the Marche Region.

Discussion

The first interface in the food-related infections investigations is the HPHS, which evaluates warnings from various sources (Competent Authorities, Hospitals, other LHUs, *etc.*). In the case described, the segnalation sources were both the emergency room of Fermo Hospital and the Departmental Services of neighboring provinces. The next phase of the investigation involved interconnected operational steps: epidemiological investigation, environmental and food samplings, laboratory analysis, and preparation of the final report. During the period of the outbreak, the regional foodborne diseases investigation guidelines (Marche Region, 2022a) that would have been available in 2022 with the following standardization of the operative flows (Figures 1 and 2), had not yet been published; however, the activity carried out did not deviate from what the document would have later provided. The management of epidemiological investigation and its subsequent sharing by HPHS with ABFHS was one of the critical issues highlighted in the case. The development of shared procedures and guidelines as previously reported, will surely improve this critical point.

In general, another critical point related to the management of foodborne infection cases is the human samples sent to several laboratories, not allowing complete epidemiological investigations of the clinical cases.

In the Marche Region, the presence of a single reference laboratory, *i.e.*, the CRRPE, where strains of *Salmonella* spp. of clinical, veterinary, food and environmental origin are sent by the regional hospital analysis laboratories and the human/food/environmental private analysis laboratories, allowed for a rapid understanding of the extent of the outbreak at a regional level. The Enteric Regional Reference Center, established by the Marche Regional Department of Health following a note of 12 September 2001 under the express request of the *Istituto Superiore di Sanità*, was subsequently formalized and further identified as the Regional Reference Laboratory for Foodborne Diseases in Humans in 2022 by Executive Decree No. 149 of October 7, 2022 (Marche Region, 2022b).

In the case described, the CRRPE also received clinical strains isolated at hospital and private laboratories of the region; that allowed detecting an increase in cases of salmonellosis in a given period (July-September 2022).

The territorial surveillance about the circulation of enteric pathogens is provided by the Enter-Net and Enter-Vet surveillance networks. The former, coordinated at national level by the Department of Infectious Diseases of the *Istituto Superiore di Sanità*, deals with European surveillance of infections caused by bacteria such as *Salmonella* spp., *Campylobacter* spp., *Shigella* spp., *Yersinia* spp., *Aeromonas* spp. and *Vibrio* spp. and has been incorporated into the European surveillance system for "Food and Waterborne Diseases" coordinated by the European Center for Disease Prevention and

Control. The latter, coordinated at national level by the National Reference Center and World Organisation for Animal Health Reference Laboratory for Salmonellosis of the *Istituto Zooprofilattico Sperimentale delle Venezie*, collects data on *Salmonella* spp. isolates from animals, food and environment.

In the Marche Region both networks, Enter-Net and Enter-Vet, are powered with data by the Regional Reference Center in collaboration with the peripheral laboratories of human, veterinary, food and environmental microbiology which send to it, the strains.

In cases of outbreaks or international alerts, but also in the management of sporadic cases, the CRRPE, supported by the National Reference Centers, is in charge of analyzing and comparing bacterial isolates of different origins providing real-time surveillance data on the circulation of enteric pathogens in the regional territory.

The above aspects show the strategic importance of the simultaneous management of animal, food, environmental and human origin strains according to a "One Health" approach.

Regarding the type and methodology of sampling used in the case study described, supporting the epidemiological investigation with environmental sampling made it possible to correlate the source of infection (*porchetta*) with the clinical cases. In addition, the positivity for *Salmonella* obtained from environmental samples carried out in a single, unrepeatable analysis allowed the ABFHS to apply the measures provided by the legislation.

The result of the inspection revealed a wrong application of HACCP principles by the FBO. As laid down in Article 5 of Regulation (EC) No 853/2004 (European Parliament and Council of the European Union, 2004b), the operator must develop, implement, and maintain one or more permanent procedures based on the principles of HACCP; one of the most important steps is to identify the CCPs at the level(s) where control proves essential to prevent or eliminate the risk or reduce it to an acceptable level. In the *porchetta* production HACCP plan, the only CCP highlighted by the FBO was cold storage, while cooking and subsequent refrigeration were treated as GMP. Cooking at the appropriate time and temperature is considered the only phase that can confer adequate sanitary and hygienic properties to the final product due to the inactivation of all non-spore-forming pathogens; subsequent blast chilling and storage at refrigeration temperature do not allow the proliferation of heat-resistant bacteria. Therefore, the simple application of good cooking and blast chilling practices does not provide a sufficient guarantee against microbiological hazards such as *Salmonella* spp.

Conclusions

Over the last three decades, a new *Salmonella* serotype has emerged worldwide, namely the MVST with the antigenic formula 4, [5],12:i:-. This serotype was first identified in chicken carcasses in Portugal in 1980 and gradually spread through the swine chain, especially in Europe and the United States (Machado *et al.*, 1990).

The description of this outbreak shows how important the collaboration between the various institutions involved (medical and veterinary services, regional authorities, *Istituto Zooprofilattico Sperimentale*) is to carry out effective official control to identify the source of foodborne cases and then to take effective and appropriate measures. Therefore, the multidisciplinary approach is important for a successful epidemiological investigation but requires trained and specialized personnel. The training activity also concerns the FBOs, as the case described has shown that, although more than 20 years have passed since the mandatory introduction of procedures based on HACCP principles, FBOs are still not fully aware of the importance of proper implementation of hygiene and health risk prevention.

In conclusion, rapid intervention is essential to act effectively, both for the accurate and timely collection of relevant information and for the subsequent conduct of inspections and sampling. In addition to intensifying data exchange networks, it is necessary to create appropriate standardized procedures for conducting epidemiological investigations, collecting food, environmental and biological samples and archiving the data in a common way allowing an appropriate evaluation by the staff involved. At the local level, a joint procedure is currently being prepared between the services

of the Prevention Department of the LHU of Fermo with the involvement of the microbiology laboratory of the Regional Hospital and the CRRPE of IZSUM. The next planned step is the involvement of other professionals (*e.g.*, general practitioners, pediatricians, private laboratories) to increase the percentage of foodborne outbreaks where the food source of infection can be identified, as defined in the Marche Region Prevention Plan.

References

- Clinical and Laboratory Standards Institute, 2023. Performance standards for antimicrobial susceptibility testing, 33rd ed. CLSI, Institute: Wayne, PA, USA.
- Codice Penale, 1930. Approvazione del testo definitivo del codice penale e s.m.i. Regio Decreto 19 ottobre 1930, n. 1398 [Decree in Italian].
- Commission of the European Communities, 2005. Regulation (EC) No 2073/2005 of 15 November 2005 on microbiological criteria of foodstuffs. In: Official Journal, L 338/1, 22/12/2005.
- ECDC, 2011. Laboratory standard operating procedure for MLVA of *Salmonella enterica* serotype Typhimurium. Available from: https://www.ecdc.europa.eu/sites/default/files/media/en/publications/Publications/1109_SOP_Salmonella_Typhimurium_MLVA.pdf.
- EFSA, ECDC, 2023. The European Union one health 2022 zoonoses report. EFSA J 21:e8442.
- European Parliament, Council of the European Union, 2002. Regulation (EC) No. 178/2002 of the European Parliament and of the Council of 28 January 2002 laying down the general principles and requirements of food law, establishing the European Food Safety Authority and laying down procedures in matters of food safety. In: Official Journal, L 31/1, 1/02/2002.
- European Parliament, Council of the European Union, 2003. Directive 2003/99/EC of the European Parliament and of the Council of 17 November 2003 on the monitoring of zoonoses and zoonotic agents, amending Council Decision 90/424/EEC and repealing Council Directive 92/117/EEC. In: Official Journal, L 325, 17/11/2003.
- European Parliament, Council of the European Union, 2004a. Regulation (EC) No 853/2004 of the European Parliament and of the Council of 29 April 2004 laying down specific hygiene rules for food of animal origin. In: Official Journal, L 139/55, 30/04/2004.
- European Parliament, Council of the European Union, 2004b. Regulation (EC) No 852/2004 of the European Parliament and of the Council of 29 April 2004 on the hygiene of foodstuffs. In: Official Journal, L 139/1, 30/04/2004.
- European Parliament, Council of the European Union, 2017. Regulation (EU) 2017/625 of the European Parliament and of the Council of 15 March 2017 on official controls and other official activities performed to ensure the application of food and feed law, rules on animal health and welfare, plant health and plant protection products, amending Regulations (EC) No 999/2001, (EC) No 396/2005, (EC) No 1069/2009, (EC) No 1107/2009, (EU) No 1151/2012, (EU) No 652/2014, (EU) 2016/429 and (EU) 2016/2031 of the European Parliament and of the Council, Council Regulations (EC) No 1/2005 and (EC) No 1099/2009 and Council Directives 98/58/EC, 1999/74/EC, 2007/43/EC, 2008/119/EC and 2008/120/EC, and repealing Regulations (EC) No 854/2004 and (EC) No 882/2004 of the European Parliament and of the Council, Council Directives 89/608/EEC, 89/662/EEC, 90/425/EEC, 91/496/EEC, 96/23/EC, 96/93/EC and 97/78/EC and Council Decision 92/438/EEC (Official Controls Regulation). In: Official Journal, L 95/1, 7/04/2017.
- Hoffmann S, Ahn JW, 2021. Economic cost of major foodborne illnesses increased \$2 billion from 2013 to 2018. Available from: <https://www.ers.usda.gov/amber-waves/2021/april/economic-cost-of-major-foodborne-illnesses-increased-2-billion-from-2013-to-2018/>.
- ISO, 2014. Microbiology of the food chain—horizontal method for the detection, enumeration and serotyping of *Salmonella*—part 3: guidelines for serotyping of *Salmonella* spp. ISO Norm 6579-3:2014. International Standardization Organization ed., Geneva, Switzerland.

- ISO, 2020. Microbiology of the food chain—horizontal method for the detection, enumeration and serotyping of *Salmonella*—part 1: detection of *Salmonella* spp.—amendment 1: broader range of incubation temperatures, amendment to the status of annex D, and correction of the composition of MSRV and SC. ISO Norm 6579-1:2017/Amd 1:2020. International Standardization Organization ed., Geneva, Switzerland.
- Italian Republic, 1988. Decreto Presidente della Repubblica 22 settembre 1988, n. 447 e smi. Approvazione del codice di procedura penale. [Regulation in Italian].
- Italian Republic, 2006. Decreto Legislativo n. 191 del 4 aprile 2006. Attuazione della direttiva 2003/99/CE sulle misure di sorveglianza delle zoonosi e degli agenti zoonotici. In: Official Journal, 119, 24-05-2006). [Decree in Italian].
- Italian Republic, 2007. Decreto Legislativo n. 193 del 06 novembre 2007. Attuazione della direttiva 2004/41/CE relativa ai controlli in materia di sicurezza alimentare e applicazione dei regolamenti comunitari nel medesimo settore. In: Official Journal, 261, 9/11/2007. [Decree in Italian].
- Italian Republic, 2017. Decreto Presidente Consiglio dei Ministri 3 marzo 2017. Identificazione dei sistemi di sorveglianza e dei registri di mortalità, di tumori e di altre patologie. [Decree in Italian]
- Italian Republic, 2021. Decreto Legislativo n. 27 del 2 febbraio 2021. Disposizioni per l'adeguamento della normativa nazionale alle disposizioni del regolamento (UE) 2017/625 ai sensi dell'articolo 12, lettera a), b), c), d) ed e) della Legge 4 ottobre 2019, n. 117. In: Official Journal, 60, 11/03/2021. [Decree in Italian].
- Machado J, Bernardo F, 1990. Prevalence of *Salmonella* in chicken carcasses in Portugal. J Appl Bacteriol 69:477-80.
- Marche Region, 2016. Deliberazione della Giunta Regionale n. 1508 del 05 dicembre 2016. Recepimento dell'intesa 212/CSR del 10 novembre 2016 ai sensi dell'articolo 8, comma 6 della legge 5 giugno 2003, n. 131, tra il governo, le regioni e le provincie autonome di Trento e Bolzano sul documento recante "Linee guida per il controllo ufficiale ai sensi dei Regolamenti CE 882/2004 e 854/2004". [Deliberation in Italian]
- Marche Region, 2021. Deliberazione della Giunta Regionale n. 1640 del 28/12/2021. Approvazione del Piano Regionale della Prevenzione 2020-2025.
- Marche Region, 2022a. Decreto del Direttore della Agenzia Regionale Sanitaria, n.98 del 22 dicembre 2022; PNP 2020-2025: approvazione della Guida Operativa per la gestione delle malattie a trasmissione alimentare nella Regione Marche. [Decree in Italian].
- Marche Region, 2022b. Decreto Dirigenziale n. 149 del 07 ottobre 2022 per l'istituzione presso la sede di Tolentino dell'Istituto Zooprofilattico Sperimentale Umbria e Marche "Togo Rosati", del Centro di Riferimento Regionale per i Patogeni Enterici e la sua individuazione come Laboratorio di Riferimento Regionale per le malattie trasmesse da alimenti in ambito umano. [Decree in Italian].
- Napoleoni M, Villa L, Barco L, Lucarelli C, Tiengo A, Baggio G, Dionisi AM, Angellotti A, Ferretti E, Ruggeri S, Staffolani M, Rocchegiani E, Silenzi V, Morandi B, Blasi G, 2024. Monophasic variant of *Salmonella* Typhimurium 4,[5],12:i:- (ACSSuGmTnpSxt type) outbreak in central Italy linked to the consumption of a roasted pork product (porchetta). Microorganisms 11:2567
- Rahn K, De Grandis SA, Clarke RC, McEwen SA, Galán JE, Ginocchio C, Curtiss R, Gyles CL, 1992. Amplification of an *invA* gene sequence of *Salmonella* Typhimurium by polymerase chain reaction as a specific method of detection of *Salmonella*. Mol Cell Probes 6:271-9.

Table 1. Results of the sampling.

Site	Environmental swabs			Roasted pork product (<i>porchetta</i>)		
	Negative	Positive	Total	Negative	Positive	Total
A	7	1	8	1	0	1
B	9	1	10	1	0	1
C	3	2	5	0	1	1
	19	4	23	2	1	3




Figure 1. Flowchart of single case and outbreak investigation as proposed by the regional foodborne diseases investigation guidelines. Source: Marche Region, 2022a (Decree No. 98/2022). FHNS, Food Hygiene and Nutrition Service; ABFHS, Animal Based Food Hygiene Service; LHU, Local Health Unit; HPHS, Hygiene and Public Health Service; FCD, foodborne disease; FBO, food business operators.




Figure 2. Flowchart of the food safety measures adopted by the competent authority as proposed by the regional foodborne diseases investigation guidelines. Source: Marche Region, 2022a (Decree No. 98/2022). FHNS, Food Hygiene and Nutrition Service; ABFHS, Animal Based Food Hygiene Service.