

Quality assessment of *Zeus faber* (Peter's fish) ovaries regularly commercialized for human consumption

Filippo Giarratana, Graziella Ziino, Valerio D'Andrea, Antonio Panebianco, Alessandro Giuffrida

Department of Veterinary Science, University of Messina, Italy

Abstract

In the last few years, the consumption of fish eggs has increased rapidly, finding widespread use also in mass catering. This increase has involved also those of the Peter's fish (*Zeus faber*). Females of this species, by their reproductive characteristics, have highly developed gonads in different periods of the year, making the raw material easy to find. The aim of the present study was to perform a quality assessment of *Zeus faber* ovaries regularly commercialized for human consumption. A total number of 34 samples, divided in fresh (11) and frozen (23), were processed for microbiological characterization, parasitological and histological evaluations. Fresh and frozen samples have significant ($P < 0.01$) differences in total bacterial charge, with values of 4.75 ± 0.5 Log CFU/g and 3.65 ± 0.7 Log CFU/g respectively. The mean value of *Enterobacteriaceae* was 2.58 ± 0.7 Log CFU/g in fresh products, while 52.17% (12) of frozen samples reported loads of < 1 Log CFU/g. No *Salmonella* spp. and *Listeria monocytogenes* were found. *Aeromonas* spp. was detected in two frozen sample (with loads of 2.2 and < 1 Log CFU/g) and in 5 fresh ovaries with value ranged from 1.70 to 3.48 Log CFU/g. *Vibrio* spp. was found in 4 (36.36%) and 3 (13.04%) of fresh and frozen samples respectively, with loads always < 1 Log CFU/g. All 31 *Vibrio* strains isolated, were identified as *Vibrio alginolyticus*, and 61.29% (19) of them was positive for the ToxRS factor and 6.45% (2) for ToxR. The 47.06% (16) of total samples showed infestations by larvae of *Anisakis* Type 1 in the serous and inside the ovary. In this last case, histologically it was found to be free larvae. This study attested satisfactory hygiene conditions for *Zeus faber* ovaries currently marked for human consumption. The presence of potentially pathogenic strains of *V. alginolyticus* and *Aeromonas* spp., but above all the frequent infestation by *Anisakis* larvae, represent a potentially hazard for the consumer.

Introduction

The use and conservation of fish gonads as foodstuff dates back more than three thousand years ago in Sardinia as gift of the Phoenicians. Today, they are considered a valuable material and sometimes, as regards of some fishes, even difficult to find. Both female and male gonads are usually considered edible foods. Male gonads are constituted by fish testes and are commonly known as *latti* or *lattumi* (Palese and Palese, 1992). Female gonads have a wider distribution and are commercialized as fresh products but also salted, dried, smoked and marinated. Depending on the source, the local traditions and the preservation technologies, gonads are processed and then commercialized like *ovarian sack* (i.e. *Bottarga*) or as *eggs*, without the serous coating (caviar, salmon eggs, lumpfish roes, etc.) (Giuffrida and Panebianco, 2008). In the last few years, the consumption of fish eggs has increased rapidly, finding widespread use also in mass catering (EUROSTAT, 2017).

This increase has involved also those of the Peter's fish (*Zeus faber*). Female of this species, thanks to their reproductive characteristics, show highly developed gonads in different periods of the year (Fulton, 1898). Moreover, this species is widespread all over Mediterranean areas, making the raw material easy to find. Several studies reported microbiological characterization of different fish roe product, revealing the presence, in some occasion, of food borne pathogens (Altug and Bayrak 2003; Boiko *et al.*, 2004; Oeleker *et al.*, 2015; Razavilar and Rezvani, 2004; Voidarou *et al.*, 2011). No information on Peter's fish roe microbiology are, instead, available. Furthermore, *Zeus faber* is often subjected to *Anisakidae* colonization that can be found frequently in ovaries (Pekmezci *et al.*, 2014; Yardimci *et al.*, 2014). For all these reasons, the aim of the present study was to perform a quality assessment of *Zeus faber* ovaries regularly commercialized for human consumption.

Materials and Methods

Sample collection and macroscopic observations

A total number of 34 *Zeus faber* ovaries (11 fresh and 23 frozen samples) were collected from local market of Mazara del Vallo in a period from October to May. Fishes were caught in FAO area 37.1.2 and 37.2.2, by the trawl fleet of Mazara del Vallo (Sicily, Italy). Mazara del Vallo is widely considered to be the most important

Correspondence: Filippo Giarratana, Department of Veterinary Science, University of Messina, Polo Universitario dell'Annunziata, 98168 Messina, Italy. Tel.: +39.090.3503768. E-mail: fgjarratana@unime.it

Key words: *Zeus faber*; fish eggs; *Anisakis*; microbiology; histology.

Contributions: the authors contributed equally.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: none.

Received for publication: 16 August 2017.

Revision received: 6 November 2017.

Accepted for publication: 7 November 2017.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

©Copyright F. Giarratana *et al.*, 2018

Licensee PAGEPress, Italy

Italian Journal of Food Safety 2018; 7:6997

doi:10.4081/ijfs.2018.6997

fishing center in Italy and contributes for more than 3/4 to the production and turnover of the national trawl fleet (ISTAT data).

Fresh ovaries were transported to our laboratory under refrigerated conditions (4°C) and processed within 24h from their arrival. Frozen samples were, instead, primarily thawed at 4°C for 24h. Each sample was carefully examined for the presumptive presence of parasites. All parasites were examined under stereoscopic microscope (Leica M 205C), and the belonging to the *Anisakis* genus was made according to guidelines proposed by Murata *et al.* (2011).

Microscopic analysis

Portions of each sample were fixed in buffered formalin (10%), embedded in paraffin, and sliced into 5- μ m sections. Sections were stained using hematoxylin-eosin (HE) and trichrome Masson. Stained sections were examined using light microscopy (Leica DM 4000B).

Microbiological analysis

Each sample was processed for the count of: i) Aerobic mesophilic bacteria (AMB) according to UNI EN ISO 4833:2004; ii) *Enterobacteriaceae* according to UNI EN ISO 21528-2:2004; iii) *Vibrio* spp. on Thiosulphate Citrate Bile Salt Sucrose Agar (TCBS Oxoid, Italy), with 3% NaCl, incubated at 37°C for 24h;

iv) *Aeromonas* spp. in Glutamate Starch Phenol Red Agar (GSP) with Ampicillin Selective Supplement (Merk, Italy), incubated at 30°C for 24h.

The detection of the following parameter was also conducted: i) *Vibrio* spp. with a preliminary enrichment on Phosphate Buffer Saline (PBS) at 30°C for 24h, then spread on TCBS, with 3% NaCl and incubated at 37°C for 24h; ii) *Aeromonas* spp. with a preliminary enrichment on PBS at 30°C for 24h, then spread on in GSP with Ampicillin and incubated at 30°C for 24h; iii) *Listeria monocytogenes* detection according to UNI EN ISO 11290-1: 2005; iv) *Salmonella* spp. detection according to UNI EN ISO 6579: 2002.

Vibrio identification: biochemical protocol

Suspected alophilic vibrios were confirmed to genus level to according to biochemical protocol suggested by Ottaviani *et al.* (2003). Isolated colonies growth on TCBS, were tested for Gram stains, Ossidase test, vibriostatic factor O/129 (150 mg, Oxoid) and growth on Kligler Iron Agar (KIA, Biolife, Italy). All the strains that resulted belonging to *Vibrio* spp. were, then, confirmed by molecular methods.

Vibrio identification: multiplex PCR assay

For DNA extraction 1 mL of broth culture was centrifuged at 12.000 rpm for 5 min; the pellet was re-suspended in 1 mL of sterile distilled water, boiled for 5 min, and centrifuged again. The supernatant was stored at 20°C until use. DNA was quantified by means of a spectrophotometer (SmartSpec Plus, Bio-Rad, Milan, Italy). For *Vibrio* genus identification specific gene *rpoA* primers were employed, according to La Neve *et al.* (2006) and Dalmaso *et al.* (2009). Confirmed strains belonging to *Vibrio* genus were selected for further identification at species level. For *V. alginolyticus*, and *V. parahaemolyticus* identification, were employed specific primers encoding a collagenase gene portion, as suggested by Di Pinto *et al.* (2005). Primers for the screening of pathogenic factors *toxR* and *toxRS* were also added to the PCR mix as reported by Xie *et al.* (2005).

Multiplex PCR mix consisted in a total volume of 50 mL containing 5 mL of Buffer 10X, 200 mmol of each dNTPs, 1.5 mmol of MgCl₂, 100 nmol of each primer, 1 mL of DNA template and 1U of Platinum Taq DNA Polymerase (Invitrogen).

pH determination

The pH of each sample was measured at room temperature with a pH meter HI90023CW (Hanna Instruments, Italy)

equipped with a Mettler Toledo electrode (InLab 427) after homogenizing 3 g of sample in 27 mL distilled water for 10 s at 1300 rpm with an Ultra-Turrax T25 macevator (Janke & Kunkel, Staufen, Germany). Each value was the mean of three replicates.

Data analysis

A t-Test was performed in order to evaluate significative differences in our results among the samples and between fresh and frozen products (XLSTAT, Microsoft Excel, Addinsoft). Significance level was assumed as P<0.01. Microbiological charges were converted in Log CFU/g for a better expression of results.

Results and Discussion

Microbiological results

Table 1 shows mean values for each microbiological parameter. The mean values for AMC and *Enterobacteriaceae* were 3.94±0.5 Log CFU/g and 1.30±1.2 Log CFU/g respectively. Fresh and frozen samples have significant (P<0.01) differences in AMC, with mean values of 4.75±0.5 log CFU/g and 3.65±0.7 log CFU/g respectively. In fresh ovaries, AMC oscillated from 3.65 to 5.17 Log CFU/g, while, in frozen samples ranged from 1.74 to 4.60 Log CFU/g. Also for *Enterobacteriaceae* loads, significant differences (P<0.01) were observed between fresh and frozen products. In fresh products *Enterobacteriaceae* charges ranged from 1.17 to 3.43 Log CFU/g, while 52.17% (12) of frozen samples reported *Enterobacteriaceae* loads <1 Log CFU/g.

Considering bacteriological hygiene indicators (AMC and *Enterobacteriaceae*), *Zeus faber* ovaries presented an overall satisfactory condition even as fresh product stored at refrigeration temperatures (Altug and Bayrak 2003; Boiko *et al.*, 2004; Oeleker *et al.*, 2015; Razavilar and Rezvani, 2004; Voidarou *et al.*, 2011).

The only microbiological concern may be related to pathogen detection. No *Salmonella* spp. and *Listeria monocytogenes* were found.

Otherwise, *Aeromonas* spp. and *Vibrio* spp. were detected in several samples of *Zeus faber* ovaries. *Aeromonas* strains have a significative incidence in marine environment all over Europe and were frequently detected in fishery product, including fish eggs (Davies *et al.*, 2001; Hänninen *et al.*, 1997; Hansen and Olafsen, 1989; Muscolino *et al.*; 2014). Bacteria belonging to *Aeromonas* genus may be, also, responsible of gastro-enteric diseases outbreaks in humans, thanks to the production of enterotoxins, cytotoxins and haemolysins (Deodhar *et al.*, 1991; Kirov, 1993). For all these reasons, they could represent a microbiological hazard for the consumer.

In our study, *Aeromonas* spp. was found in seven samples: two in frozen and five in fresh samples. The two positive frozen samples, were characterized by loads of 2.2 Log CFU/g and <100 CFU/g respectively. In fresh ovaries, instead, *Aeromonas* spp was reported in 45.45% (5) of samples with loads ranging from 2.0 to 3.0 Log CFU/g. However, *Aeromonas* spp. mean loads revealed no significative difference (P>0.01) between fresh and frozen samples (Table 1). Considering its psychrophilic behavior, its tolerance and survival at frozen temperatures as well as its stress resistant variants, this study confirms *Aeromonas* spp. as potential microbiological hazard in *Zeus faber* ovaries intended for consumption (Castro-Escarpulli *et al.*, 2003; Giuffrida *et al.*, 2010).

Vibrio spp. was detected in the 36.36% (4) and 13.04% (3) of fresh and frozen samples respectively, but always with loads <1 Log CFU/g.

A total number of 31 suspected *Vibrio* spp. strains (14 from fresh and 17 from frozen samples) were isolated from TCBS with 3% NaCl. All suspected strains were confirmed as belonging to *Vibrio* spp., resulted positive for biochemical test and for molecular identification at genus level (*rpoA* +). Their presence in frozen samples, confirmed the remarkable resistance of halophilic *Vibrio* at freezing temperatures (Bang and Drake, 2002; Johnston and Brown, 2002). Multiplex PCR analysis

Table 1. Mean values ± standard deviation expressed in Log CFU/g, of microbiological loads.

	AMB	<i>Enterobacteriaceae</i>	<i>Aeromonas</i> spp.
Fresh	4.75±0.5 ^a (11/11)	2.58±0.7 ^a (11/11)	1.09±1.3 ^a (5/11)
Frozen	3.65±0.7 ^b (23/23)	0.58±0.7 ^b (11/23)	0.09±0.4 ^a (1/23)
Total	3.94±0.8 (34/34)	1.30±1.2 (22/34)	0.42±0.9 (6/34)

^{a,b}Different letters represent significative differences (P<0.01), (positive samples/total samples).

revealed that all 31 strains isolated were *V. alginolyticus*.

This bacteria is often found in Italian marine coastal environments and sea products (Dumontet *et al.*, 2000; Panebianco *et al.*, 2011; Narracci *et al.*, 2013; Ziino *et al.*, 2010; Ziino *et al.*, 2014). It is also, widely spread all over the world, and usually has the highest prevalence in *Vibrio* communities (Schets *et al.*, 2010; Jones *et al.*, 2013). *V. alginolyticus* pathogenic potential in humans is well recognized, causing various infections and inflammation patterns (Schmidt *et al.*, 1979). It raised great importance as emergent foodborne pathogen, responsible for several cases of gastroenteritis linked to the consumption of fish products (Mustapha *et al.*, 2013). Moreover, in immune-compromised subjects, serious extra-intestinal infections, such as septic shock, have been reported as consequence of consumption of seafood contaminated by *V. alginolyticus* (Dong-Young *et al.*, 2008). ToxRS and ToxR pathogenic factors encode trans-membrane proteins that mediate the regulation of the virulence gene expression in several pathogenic *Vibrio* strains, including *V. alginolyticus* (Das *et al.*, 2016). These factors are frequently investigated for a better characterization of bacterial pathogenic potential (Fu *et al.*, 2016). Moreover, ToxR is widely proposed for the specie identification and phylogenetic analyses (Montieri *et al.*, 2010). Considering all the 31 *V. alginolyticus* strains, 61.5% carried the ToxRS factor and in lower percentage (6.5%) the *ToxR* gene. In particular, among the 14 strains isolated from fresh samples, 12 (85.7%) were ToxR positive and 2 (14.3%) carried the ToxRS pathogenic factor. Otherwise, among the 17 strains isolated from frozen samples, 7 (41.2%) were ToxR positive but no ToxRS positive strains were found. It was asserted that ToxR may assist *V. alginolyticus* in host target cells adhesion, which is crucial step for the initiation of the infection (Chang *et al.*, 2002). As reported by previous studies, fish roe represent an ideal setting for *Vibrio* colonization (Voidarou *et al.*, 2011). Our results highlight the role of *Zeus faber* ovaries as source of *V. alginolyticus* pathogenic strains, which could be potentially implicated in foodborne disease episodes.

pH results

The pH mean values was 6.32 ± 0.2 for fresh ovaries and 6.22 ± 0.3 for frozen products with no significant differences ($P > 0.01$) between the two kind of samples.

Macroscopic and microscopic analysis results

The 47.05% (16) of total samples

revealed parasitic infestation by presumptive nematode, confirmed by stereomicroscopic observation as *Anisakis* larvae type I. Specifically, larvae were found in 16 ovaries: 45.45% (5) and (11) 47.82% of fresh and frozen samples respectively. Various levels of infestation were observed: from isolated larvae to a massive infestation, in which nematodes were bunch-shaped aggregated (Figure 1).

The histological investigation showed the presence of various patterns, ascribable to different gonadic development stages (Abou-Seedo *et al.*, 2003; Macrì *et al.*, 2011). Specifically, in some analyzed ovaries, a primary stage of oocyte growth (Previtellogenic stage) was observed (Figure 2), while in other samples a second-

ary stage of oocyte maturation (Vitellogenic stage) was predominant (Figure 3), with no relation to season or area of fishing.

In samples infested by *Anisakis* larvae, parasites were found encysted on the serosa, and were not observed tissue and cellular reactions (Figure 4). Only in two cases, however, it was possible to observe parasites inside the gonads, not surrounded by any inflammatory reaction (Figure 5). Our results confirm the parasitological risk due to the consumption of *Zeus faber* ovaries. *Anisakis* larvae detection deserve great attention, as the nematode have been also observed in the depth of the gonads. The lack of connective capsule and inflammatory response around the parasites, are consequences of larvae tissue mobilization,




Figure 1. Massive infestation of *Anisakis* larvae type I in fresh sample of *Zeus faber* ovaries.




Figure 2. Previtellogenic stage: a primary stage of oocyte growth in fresh sample of *Zeus faber* ovaries.




Figure 3. Vitellogenic stage: a secondary stage of oocyte maturation in fresh sample of *Zeus faber* ovaries.




Figure 4. Presence of *Anisakis* larvae type I encysted on the serosa, without tissue and cellular reactions, in fresh sample of *Zeus faber* ovaries.




Figure 5. Presence of *Anisakis* larvae type I inside the gonads, not surrounded by any inflammatory reaction, in frozen sample of *Zeus faber* ovaries. Aspect of structural damage in frozen samples: oocytes coerced, with irregular contours, fragmented and well-spaced. The cytoplasm lost its characteristic granular appearance and are evident intracytoplasmic vacuoles.

searching for favorable conditions after the fish death. In these cases, *Anisakis* could be unnoticed to a superficial inspection, especially if a moderate infestation occur. However, it would, still, represent a risk for the consumer as thermal shock is not sufficient to avoid possible allergic reaction in sensitive subjects, caused by allergens resistant to high temperatures and freezing conditions (Audicana and Kennedy, 2008; Speciale *et al.*, 2017).

In frozen samples, other microscopic findings are the damage related to the presence of intra and extra-cytoplasmic ice crystals. During thawing process, ice particles damaged the oocyte cell wall, causing the loss of cellular content. In particular, oocytes appeared coerced, with irregular contours, fragmented and well spaced. The cytoplasm lost its characteristic granular appearance, becoming uniformly acidophilous and, sometimes, voluminous intracytoplasmic vacuoles were observed (Figure 5). As reported in previous study microscopic analysis resulted a valid means to discriminate fresh products from those thawed (Meistro *et al.*, 2016; Muscolino *et al.*, 2012).

Conclusions

In conclusion, *Zeus faber* fish ovaries showed more than satisfactory hygiene conditions. The only microbiological concern was related to the presence of potentially pathogenic strains of *V. alginolyticus* and *Aeromonas* spp.. However, microbiological hazard can be reduced by adequate cooking and the prevention of cross-contamination. In relation to the presence of *Anisakis* larvae, only an appropriated consumer information seems the most important measure to prevent allergic manifestations, beside an efficient thermal treatment or similar strategies in order to inactivate nematodes (Anastasio *et al.*, 2015; Giarratana *et al.*, 2012; Giarratana *et al.*, 2014; Giarratana *et al.*, 2015; Giarratana *et al.*, 2017a; Giarratana *et al.*, 2017b; Valero *et al.*, 2015).

References

- Abou-Seedo F, Dadzie S, Al-Kanaan KA, 2003. Histology of ovarian development and maturity stages in the yellowfin seabream *Acanthopagrus latus* (Teleostei: Sparidae) (Hottuyn, 1782)

- reared in cages. *Kuwait J Sci Engin* 30.
- Altug G, Bayrak Y. 2003. Microbiological analysis of caviar from Russia and Iran. *Food Microbiol* 20:83-6.
- Anastasio A, Smaldone G, Cacace D, Marrone R, Lo Voi A, Santoro M, Cringoli G, Pozio E, 2015. Inactivation of *Anisakis pegreffii* larvae in anchovies (*Engraulis encrasicolus*) by salting and quality assessment of finished product. *Food Control* 64:115-9.
- Audicana MT, Kennedy MW, 2008. *Anisakis simplex*: from obscure infectious worm to inducer of immune hypersensitivity. *Clin Microbiol Rev* 21:360-79.
- Bang W, Drake MA, 2002. Resistance of cold-and starvation-stressed *Vibrio vulnificus* to heat and freeze-thaw exposure. *J Food Protect* 65:975-80.
- Boiko AV, Pogorelova NP, Zhuravleva LA, Lartseva LV, 1993. Microbial colonization of the caviar of the sturgeon fishes. *Gigiena i sanitaria* 11:30-1.
- Castro-Escarpulli G, Figueras MJ, Aguilera-Arreola G, Soler L, Fernandez-Rendon E, Aparicio GO, Chacon MR, 2003. Characterisation of *Aeromonas* spp. isolated from frozen fish intended for human consumption in Mexico. *Int J Food Microbiol* 84:41-9.
- Chang C, Qing-bai W, Zhu-Hong L, Jing-jing Z, Xiao J, Hong-yan S, Chao-qun H, 2012. Characterization of role of the *toxR* gene in the physiology and pathogenicity of *Vibrio alginolyticus*. *Antonie Van Leeuwenhoek* 101:281-8.
- Dalmaso A, La Neve F, Suffredini E, Croci L, Serracca L, Bottero MT, Civera T, 2009. Development of a PCR Assay Targeting the *rpoA* Gene for the Screening of *Vibrio* Genus. *Food Analytical Methods* 2:317-24.
- Das SC, Kumar A, Kaushik P, Kumari S, 2016. Pathogenic and pandemic *Vibrio parahaemolyticus* detection in fish and shellfish isolates. *Indian J Geo-Marine Sci* 45:1195-8.
- Davies AR, Capell C, Jehanno D, Nychas GJ, Kirby RM, 2001. Incidence of food-borne pathogens on European fish. *Food Control* 12:67-71.
- Deodhar LP, Saraswathi K, Varudkar A, 1991. *Aeromonas* spp. and their association with human diarrheal disease. *J Clin Microbiol* 29:853-6.
- Di Pinto A, Ciccarese G, Tantillo G, Catalano D, Forte VT, 2005. A collagenase-targeted multiplex PCR assay for identification of *Vibrio alginolyticus*, *Vibrio cholerae*, and *Vibrio parahaemolyticus*. *J Food Protect* 68:150-3.
- Dong-Young L, Soo-Youn M, Sang-Oh L, Hee-Young Y, Hee-Joo L, Mi Suk L, 2008. Septic Shock due to *Vibrio alginolyticus* in a Cirrhotic Patient: The First Case in Korea. *Yonsei Med J* 49:329-32.
- Dumontet S, Krovacek K, Svenson SB, Pasquale V, Baloda SB, Figliuolo G, 2000. Prevalence and diversity of *Aeromonas* and *Vibrio* spp. in coastal waters of Southern Italy. *Comparative Immunol Microbiol Infect Dis* 23:53-72.
- EUROSTAT 2017. Production of fish eggs for human consumption from aquaculture (from 2008 onwards). EU open data portal. Available at: <https://data.europa.eu/euodp/data/dataset/IexxNEN7Uhv2yXTYlaog>
- Fu K, Li J, Wang Y, Liu J, Yan H, Shi L, Zhou L, 2016. An Innovative Method for Rapid Identification and Detection of *Vibrio alginolyticus* in Different Infection Models. *Front Microbiol* 7:651.
- Fulton TW, 1898. The ovaries and ovarian eggs of the angler or frog-fish (*Lophius piscatorius*) and of the John Dory (*Zeus faber*). *Ann Rep Fish Board Scotland* 16:125-34.
- Giarratana F, Giuffrida A, Gallo F, Ziino G, Panebianco A, 2012. Study of the Resistance Variability of *Anisakis* Larvae to Some Technological Stressors. In *Veterinary Science*, Springer Berlin Heidelberg, pp. 155-159.
- Giarratana F, Muscolino D, Beninati C, Giuffrida A, Panebianco A, 2014. Activity of *Thymus vulgaris* essential oil against *Anisakis* larvae. *Experiment Parasitol* 142: 7-10.
- Giarratana F, Muscolino D, Panebianco F, Patania A, Benianti C, Ziino G, Giuffrida A, 2015. Activity of R (+) limonene against *Anisakis* larvae. *Ital J Food Saf* 4:5499.
- Giarratana F, Muscolino D, Ziino G, Giuffrida A, Marotta SM, Lo Presti V, Chiofalo V, Panebianco A, 2017a. Activity of *Tagetes minuta* Linnaeus (*Asteraceae*) essential oil against L3 *Anisakis* larvae type I. *Asian Pacific J Trop Medicine* 10:461-5.
- Giarratana F, Muscolino D, Ziino G, Lo Presti V, Rao R, Chiofalo V, Giuffrida A, Panebianco A, 2017a. Activity of Catmint (*Nepeta cataria*) essential oil against *Anisakis* larvae. *Trop Biomed* 34:22-31.
- Giarratana F, Panebianco F, Muscolino D, Beninati C, Ziino G, Giuffrida A, 2015. Effect of allyl isothiocyanate against *Anisakis* larvae during the anchovy marinating process. *J Food Protect* 78:767-71.
- Giuffrida A, Panebianco A, 2008. Igiene e tecnologie dei prodotti della pesca freschi e trasformati., In: Colavita G, ed. *Igiene e tecnologia dei prodotti di origine animale*. Le Point Vétérinaire Italie, Milano, Italy, pp 274-276.
- Giuffrida A, Ziino G, Valenti D, Donato G, Panebianco A, 2007. Application of an interspecific competition model to predict the growth of *Aeromonas hydrophila* on fish surfaces during refrigerated storage. *Archiv Lebensmittelhyg* 58:136-41.
- Hänninen ML, Oivanen P, Hirvelä-Koski V, 1997. *Aeromonas* species in fish, fish-eggs, shrimp and freshwater. *Int J Food Microbiol* 34:17-26.
- Hansen G H, Olafsen JA, 1989. Bacterial colonization of cod (*Gadus morhua* L.) and halibut (*Hippoglossus hippoglossus*) eggs in marine aquaculture. *App Environment Microbiol* 55:1435-46.
- ISO 11290-1:1996. Microbiology of food and animal feeding stuffs. Horizontal method for the detection and enumeration of *Listeria monocytogenes*. Part 1: Detection method. International Standardization Organization Geneva, Switzerland.
- ISO 21528-2:2004. Microbiology of the food chain — Horizontal methods for the detection and enumeration of Enterobacteriaceae Part 2: Colony-count method. International Standardization Organization Geneva, Switzerland.
- ISO 4833-1:2013. Microbiology of the food chain. Horizontal method for the enumeration of microorganisms. Part 1: Colony count at 30 degrees C by the pour plate technique. International Standardization Organization Geneva, Switzerland.
- ISO 6579:2002. Microbiology of food and animal feeding stuffs. Horizontal method for the detection of *Salmonella* spp. International Standardization Organization Geneva, Switzerland.
- Johnston MD, Brown MH, 2002. An investigation into the changed physiological state of *Vibrio* bacteria as a survival mechanism in response to cold temperatures and studies on their sensitivity to heating and freezing. *J App Microbiol* 92:1066-77.
- Jones EH, Feldman KA, Palmer A, Butler E, Blythe D, Mitchell CS, 2013. *Vibrio* infections and surveillance in Maryland, 2002-2008. *Public Health Rep* 128:537-45.
- Kirov SM, 1993. The public health significance of *Aeromonas* spp. in foods. *Int J Food Microbiol* 20:179-98.
- La Neve F, Pedonese F, Nuvoloni R,

- D'Ascenzi C, Dalmaso A, Civera T, 2006. Identificazione di Vibrioni di interesse sanitario in orate di allevamento mediante metodiche biomolecolari. LX Convegno Nazionale della Società Italiana delle Scienze Veterinarie, 432.
- Macri F, Rapisarda G, Marino G, De Majo M, Aiudi G, 2011. Use of Laparoscopy for the Evaluation of the Reproductive Status of Tench (*Tinca tinca*). *Reprod Domestic Anim* 46:130-3.
- Meistro S, Pezzolato M, Muscolino D, Giarratana F, Baioni E, Panebianco A, Bozzetta E, 2016. Histology as a valid tool to differentiate fresh from frozen-thawed marinated fish. *J Food Protect* 79:1457-9.
- Montieri S, Suffredini E, Ciccozzi M, Croci L, 2010. Phylogenetic and evolutionary analysis of *Vibrio parahaemolyticus* and *Vibrio alginolyticus* isolates based on *toxR* gene sequence. *New Microbiol* 33:359-72.
- Murata R, Suzuki J, Sadamasu K, Kai A, 2011. Morphological and molecular characterization of *Anisakis* larvae (Nematoda: Anisakidae) in *Beryx splendens* from Japanese waters. *Parasitol Int* 60:193-8.
- Muscolino D, Giarratana F, Beninati C, Tornambene A, Panebianco A, Ziino G, 2014. Hygienic-sanitary evaluation of sushi and sashimi sold in Messina and Catania, Italy. *Ital J Food Saf* 3:134-6.
- Muscolino D, Giarratana F, Giuffrida A, Panebianco A, 2012. Inspective investigation on swordfish (*Xiphias gladius*) frozen slices of commerce: anatomohistopathological findings. *Czech J Food Sci* 30:206-10.
- Mustapha S, Mustapha EM, Nozha C, 2013. *Vibrio alginolyticus*: an emerging pathogen of food borne diseases. *Int J Sci Tech* 2:302-9.
- Narracci M, Acquaviva MI, Cavallo RA, 2013. Mar Piccolo of Taranto: *Vibrio* biodiversity in ecotoxicology approach. *Environ Sci Pollut Res Int* 21:2378-85.
- Oeleker K, Alter T, Kleer J, Pund RP, Gözl G, Hildebrandt G, Huehn S, 2015. Microbiological and chemical investigation of caviar at retail. *J Verbraucherschutz Lebensmittelsicherheit* 10:35-7.
- Ottaviani D, Masini L, Bacchiocchi S, 2003. A biochemical protocol for the isolation and identification of current species of *Vibrio* in seafood. *J App Bacteriol* 95:1277-84.
- Palese L, Palese A, 1992. Il controllo sanitaria e qualitative dei prodotti alimentari della pesca, vol. II. Ed. Piccin, Padova.
- Panebianco A, Giuffrida A, Tasca D, Atanassova V, Ziino G, 2011. Shelf-life and microbiological properties of refrigerated sea urchin (*Paracentrotus lividus*) roe. *Archiv Lebensmittelhygiene* 62:11-5.
- Pekmezci GZ, Onuk EE, Bolukbas CS, Yardimci B, Gurler AT, Acici M, Umur S, 2014. Molecular identification of *Anisakis* species (Nematoda: Anisakidae) from marine fishes collected in Turkish waters. *Vet Parasitol* 201:82-94.
- Razavilar V, Rezvani S, 2004. Microbial risk assessment of Persian caviar during processing and cold storage. *Dev Food Sci* 42:441-6.
- Schets FM, Van Den Berg HH, Rutjes SA, De Roda Husman AM, 2010. Pathogenic *Vibrio* species in dutch shellfish destined for direct human consumption. *J Food Prot* 73:734-8.
- Schmidt U, Chmel H, Cobbs C, 1979. *Vibrio alginolyticus* infections in humans. *J Clin Microbiol* 10:666-8.
- Speciale A, Trombetta D, Saija A, Panebianco A, Giarratana F, Ziino G, Minciullo PL, Cimino F, Gangemi S, 2017. Exposure to *Anisakis* extracts can induce inflammation on in vitro cultured human colonic cells. *Parasitol Res* 116:2471-7.
- Valero A, Romero MC, Gómez-Mateos M, Hierro I, Navarro MC, 2015. Natural products: Perspectives in the pharmacological treatment of gastrointestinal anisakiasis. *Asian Pacific J Trop Medicine* 8:612-7.
- Voidarou C, Alexopoulos A, Plessas S, Noussias H, Stavropoulou E, Fotou K, Demertzi-Akrida K, 2011. Microbiological quality of grey-mullet roe. *Anaerobe* 17:273-5.
- Xie ZY, Hu CQ, Chen C, Zhang LP, Ren CH, 2005. Investigation of seven *Vibrio* virulence genes among *Vibrio alginolyticus* and *Vibrio parahaemolyticus* strains from the coastal mariculture systems in Guangdong, China. *Lett App Microbiol* 41:202-7.
- Yardimci B, Pekmezci GZ, Onuk EE, 2014. Pathology and molecular identification of *Anisakis pegreffii* (Nematoda: Anisakidae) infection in the John Dory, *Zeus faber* (Linnaeus, 1758) caught in Mediterranean Sea. *Ankara Universitesi Veteriner Fakultesi Dergisi* 61:233-6.
- Ziino G, Donato G, Giarratana F, Panebianco A, 2014. Bacteriological analysis to *Arca noae* (Linné, 1758) in the Sud of Italy: the first record of halophilic *Vibrio*. *Cahiers Biol Marine* 55:389-97.
- Ziino G, Nibali V, Panebianco A, 2010. Bacteriological investigation on "Mauro" sold in Catania. *Vet Res Comm* 34:157-61.