

Olive mill wastewater phenolic concentrate as natural antioxidant against lipid-protein oxidative deterioration in chicken meat during storage

Rossana Roila,¹ Andrea Valiani,²
Dino Miraglia,¹ David Ranucci,¹
Claudio Forte,²

Massimo Trabalza-Marinucci,¹
Maurizio Servili,³ Michela Codini,⁴
Raffaella Branciarì¹

¹Dipartimento di Medicina Veterinaria,
Università degli Studi di Perugia;

²Istituto Zooprofilattico Sperimentale
dell'Umbria e delle Marche "Togo

Rosati", Perugia; ³Dipartimento di
Scienze Agrarie, Alimentari ed

Ambientali, Università degli Studi di
Perugia; ⁴Dipartimento di Scienze

Farmaceutiche, Università degli Studi di
Perugia, Italy

Abstract

Considering that many plant-derived substances show antioxidant and antimicrobial properties, natural antioxidant administered through feed in livestock animals could increase the shelf life of meat and meat products. The aim of this work was to study the effect of olive oil by-products on chicken meat lipid and protein oxidation and oxidative stability during storage. Two hundred and ninety-seven 22-day-old fast growing (Ross 308) female chicks were randomly assigned to three experimental grower-finisher diets: i) a basal control diet (CTR), ii) CTR diet supplemented with a low dosage (4.8%) of olive mill wastewater extract (L-OW) and iii) CTR diet supplemented with a high dosage (9.9%) of olive mill wastewater extract (H-OW). Breast meat of animals belonging to each experimental group was sliced, overwrapped with oxygen-permeable packaging and analysed at three different storage times (zero, three and seven days). At the three sampling times considered, for all samples, colour coordinates (a^*), saturation index, Hue angle, peroxide value, thiobarbituric reactive substance, carbonyl assay and the oxygen radical absorbance capacity determinations were performed. No differences in colour were detected among the groups in all the sampling times considered. In conclusion, the supplementation of chicken diet with olive mill wastewater extract (OW) affected oxidation of meat, retarding lipid and protein

oxidation and improving antioxidant activity during storage.

Introduction

Oxidation is one of the major causes of quality deterioration limiting meat acceptability. Oxidative deterioration, in any type of meat, is responsible for discoloration, development of off-flavour, formation of toxic compounds and short shelf-life. Meat becomes susceptible to oxidative deterioration due to high concentrations of unsaturated lipids, heme pigments in specific species, metal catalysts and a range of oxidizing agents in the muscle tissue (Branciarì *et al.*, 2015a).

Oxidation in lipid and protein fractions of meat has been demonstrated as the main non-microbial cause of quality deterioration during processing. This happens because lipids and proteins in meat are easily susceptible to oxidative damages due to the rapid depletion of endogenous antioxidants after slaughter (Xiao *et al.*, 2013). However, the susceptibility of meat to oxidation has also been found to be influenced by animal breed and specie, muscle type and anatomical location (Min *et al.*, 2008). Moreover, the type of diet consumed by animals during the last part of their life has a great influence on the susceptibility of meat to oxidation.

In recent years special attention has been paid to several natural antioxidants (Insani *et al.*, 2008; Ranucci *et al.*, 2013; Roila *et al.*, 2016) that can be applied directly in food as technological strategies or through animal diet to reduce or prevent oxidative process in muscle food (Dal Bosco *et al.*, 2012; Novelli *et al.*, 2014). Technological strategies involve the application of antioxidants directly into the meat and meat products or by coating packaging materials with plant extracts to improve the oxidative stability of the products. In dietary manipulations antioxidants are introduced into the muscle via feed.

Olive oil by-products are amongst those plant derived natural compounds that can be used as potential sources of antioxidants for muscle food preservation and nutritional quality improvement (Branciarì *et al.*, 2017) because of their high content of phenolic substances (Novelli *et al.*, 2014). In particular, olive mill wastewater is characterised by a high content in hydroxytyrosol (3,4-DHPEA), tyrosol (*p*-HPEA), secoiridoids derivatives, in particular the dialdehydic form of decarboxymethyl elenolic acid linked to 3,4-DHPEA or *p*-HPEA (3,4-DHPEA-EDA or *p*-HPEA-EDA) and verbascoside (Branciarì

Correspondence: Raffaella Branciarì,
Dipartimento di Medicina Veterinaria,
Università degli Studi di Perugia, Via San
Costanzo 4, 06126 Perugia (PG), Italy.
Tel.: +39.075.5857932 - Fax: +39.075.5857932.
E-mail: raffaella.branciarì@unipg.it

Key words: Poultry, TBARS, Carbonyls,
Oxidative Stability

Conflict of interest: the authors declare no
potential conflict of interest.

Funding: this work is part of a research project
funded by the Italian Ministry of Health (RC
007/2012 "Valutazione dell'effetto di una
dieta integrata con polifenoli derivanti da
reflui dell'industria elaiotecnica sulle caratte-
ristiche igienico-sanitarie, chimico-fisiche,
reologiche e organolettiche della carne fresca
e delle preparazioni di carne: filiera del pollo
da carne").

Received for publication: 13 February 2018.

Revision received: 12 June 2018.

Accepted for publication: 16 July 2018.

This work is licensed under a Creative
Commons Attribution-NonCommercial 4.0
International License (CC BY-NC 4.0).

©Copyright R. Roila *et al.*, 2018

Licensee PAGEPress, Italy

Italian Journal of Food Safety 2018; 7:7342

doi:10.4081/ijfs.2018.7342

et al., 2017). No data are currently available on the effects of dietary supplementation with olive mill wastewater on chicken meat oxidation during storage.

The aim of this work was to evaluate the effect of dietary treatment with olive mill wastewater on lipid and protein oxidative stability during storage.

Materials and Methods

Two hundred ninety-seven 22-days-old female Ross 308 chicks, previously fed with the same starter diet from day 1 to day 21, were allocated for a total of 20 days in an experimental farm (Umbria, Italy) under environmental condition simulating those present in conventional intensive systems. The subjects were randomly assigned to one of the three experimental dietary groups and then divided in three replicates of 33 birds each (experimental units). Groups were fed three different diets: a basal grower-finisher control diet (CTR); CTR feed added with low dosage (4.8%) of olive mill wastewater (L-OW); CTR feed added with high dosage (9.9%) of olive mill wastewater (H-OW). The diets resulted isoenergetic and

isonitrogenous and were formulated to meet National Research Council standards for broiler chickens (NRC, 1994). Olive mill waste water polyphenol content of the different feeds was analyzed resulting 263.2 mg/kg for the L-OW diet and 556.5 mg/kg for the H-OW feed. Olive mill wastewater supplement was obtained as reported by Branciarì *et al.* (2016) from processing of Italian cultivar *Moraiolo* of *Olea europaea* using a filtration system based on progressive permeability membranes.

At the end of the trial, at 41 days of age, the animals were transported and slaughtered at a local slaughterhouse. Immediately after slaughter the carcasses were promptly chilled to 4°C. After 24 h the *Pectoralis major* was removed from carcasses, sliced, randomly assigned to retail packs overwrapped with oxygen-permeable packaging and stored at 4±1°C for 7 days. Packed meat samples were analysed during storage for colour, peroxide value (POV), thiobarbituric reactive substances (TBARS), carbonil assay and ORAC_{FL}.

Lipid oxidation

For the evaluation of primary meat lipid oxidation, the POV was determined, during storage, as reported by Branciarì *et al.* (2015b). Secondary meat lipid oxidation was assessed using the TBARS according to Tarladgis *et al.* (1960); the values were expressed as mg malondialdehyde (MDA)/kg.

Protein oxidation

Protein oxidation, measured by total carbonyl content, was evaluated by derivatization with DNP_H according to the method described by Reznick and Packer (1994) with slight modifications: 2 g of meat

were thawed, minced and then homogenized 1:10 (w/v) in phosphate buffer (pH 7.4), consisting of 50 mM NaH₂PO₄ and 1mM EDTA, using an ultraturax homogenizer for 30 s. The homogenates were divided in two equal aliquots of 0.1 mL. Afterwards proteins were precipitated in both aliquots by adding 1 mL of 10% TCA and centrifuged for 5 min at 5000 rpm. Finally, the supernatants were removed, one pellet was treated with 1 mL 2 N HCl (for quantifying protein concentration) and the other one with an equal volume of 0.2% (w/v) DNP_H in 2 N HCl (for carbonyl concentration measurement). Both samples were incubated for 1 h at room temperature (shaken every 15 min). Afterwards the samples were precipitated with 1 mL of 10% TCA and washed twice with 1 mL of 1:1 ethanol/ethyl acetate (v/v), shaken and centrifuged for 5 min at 10000 rpm. The pellets were then dissolved in 1.5 mL of 20 mM sodium phosphate buffer (pH 6.5) containing 6 M guanidine hydrochloride, stirred and centrifuged for 2 min at 5000 rpm to remove insoluble fragments. Protein concentration was calculated from absorption at 280 nm using bovine serum albumin as standard.

The number of carbonyls was measured at 370 nm and expressed as nmol of carbonyl per mg of protein using the adsorption coefficient for the protein hydrazones (21.0 mM⁻¹ cm⁻¹).

Colour measurement

The colour measurement was performed, during each of the sampling intervals, after a 30 minutes bloom period at refrigeration temperature using a CR400 Minolta Chromameter (Minolta, Osaka, Japan – light source of D65 calibrated against a standard

white tile). The results were expressed as redness (a*), hue value (tan-1 b*/a*) and saturation index, or chroma ((a*² + b*²)/2), (CIE L*a*b* colour system, 1976).

Antioxidant capacity of meat

The antioxidant capacity of meat (ten samples for each treatment in triplicate) was determined using the oxygen radical absorbance capacity method (ORAC_{FL}) based on the fluorescence decay rate of a probe in the presence of a radical oxygen species (ROO) compared with that of a reference standard, Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid, Sigma-Aldrich, Steinheim, Germany). The extraction was performed on 2 g of meat sample according to Branciarì *et al.* (2015b).

The ORAC_{FL} assays were carried out on a FLUOstar OPTIMA microplate fluorescence reader (BMG LABTECH, Offenburg, Germany) at an excitation wavelength of 485 nm and an emission wavelength of 520 nm. The procedure was based on the method of Branciarì *et al.* (2015a). Briefly 2,20-azobis (2-methylpropionamide) dihydrochloride (AAPH; Sigma-Aldrich) was used as a peroxy radical generator, Trolox was used as a reference antioxidant standard, and fluorescein was used as fluorescent probe. A 100 µL volume of diluted sample, blank or Trolox calibration solution (10-80 µmol) was mixed with 1 mL of fluorescein (80 nM), then 200 µL of each mixture was placed in a well of the microplate. The microplate was placed in the reader and preincubated for 15 min at 37°C. To each well, 60 µL of AAPH was automatically added to initiate the reaction. The fluorescence (FL) was measured every 1.9

Table 1. Effect of storage time on peroxide value (POV meq O₂kg⁻¹ fat), thiobarbituric acid-reactive substances (TBARS mg MDA kg⁻¹) and carbonyls (nmol/mg protein) in breast meat stored at 4°C

Days of storage	Sample	POV	TBARS	Carbonyls
0	CTR	1.39 ^{bx}	0.56 ^{bx}	1.77 ^{bx}
	L-OW	0.89 ^{ax}	0.38 ^{abx}	1.35 ^{ax}
	H-OW	0.77 ^{ax}	0.31 ^{ax}	1.23 ^{ax}
3	CTR	2.35 ^{by}	0.88 ^{by}	1.93 ^{by}
	L-OW	1.63 ^{ay}	0.48 ^{ay}	1.49 ^{ax}
	H-OW	1.57 ^{ay}	0.41 ^{ay}	1.33 ^{ay}
7	CTR	0.31 ^{bz}	1.60 ^{bz}	2.15 ^{by}
	L-OW	0.53 ^{az}	0.76 ^{az}	1.60 ^{ax}
	H-OW	0.61 ^{ax}	0.69 ^{az}	1.55 ^{ay}
SEM	-	0.09	0.132	0.078
P-value	T	<0.001	<0.05	<0.001
	D	<0.001	<0.05	<0.001
	TXD	<0.001	0.057	0.897

CTR, meat from chicken fed a standard diet; L-OW, meat from chicken fed enriched diet with low dosage of olive mill wastewater; H-OW, meat from chicken fed enriched diet with high dosage of olive mill wastewater. a,b Within each day of storage, different superscripts indicate differences between dietary treatments (P<0.05). ^{x,y,z} Within each dietary treatment, different superscripts indicate differences between days of storage (P<0.05); T, time effect; D, diet effect.

min. All the reaction mixtures were prepared in duplicate, and at least three independent assays were performed for each sample. The final ORAC_{FL} values were calculated by using a linear regression equation ($Y = a + bX$) to describe the relationship between the Trolox concentration (Y) and the net area under the FL decay curve (X). Linear regression was used in the range of 10-80 IM Trolox. The data are expressed as micromoles of Trolox equivalents (TE) per gram of sample ($\mu\text{mol TE g}^{-1}$) by applying the following formula:

$$\text{Orac}(\mu\text{molTE}) = \frac{\text{Trolox}(AUC_{\text{Sample}} - AUC_{\text{Blank}})k}{(AUC_{\text{Trolox}} - AUC_{\text{Blank}})}$$

where *Ctrolox* is the concentration of Trolox, *k* is the sample dilution factor, and AUC is the area below the fluorescence decay curve of the sample, the blank and Trolox, respectively, calculated by applying the following formula (Ou *et al.*, 2001) in a Microsoft Excel spreadsheet (Microsoft, Washington, DC, USA)

$$AUC = 0.5 + f_1/f_0 + \dots + f_i/f_0$$

where f_1 is the initial fluorescence reading at $t = 0$ min and f_i is the fluorescence reading at time *i*. The net AUC for each sample was obtained by subtracting the AUC of the corresponding blank from that of the sample.

Statistical analysis

The data were analysed using the GLM procedure of SAS (2001). ANOVA model was used with diet (CTR, L-OW and H-OW) and time (day 0, day 3 and day 7) as fixed variables. The differences of the means were analysed using Tukey test and were considered to be significant when $P < 0.05$.

Results

The POV, TBARS values and the amount of carbonyl compounds are reported in Table 1. POVs showed an increase at 3 days of storage than decreased at 7 days. The POV's value was higher in CTR sample at 0 and 3 days of storage. At 7 days of storage higher value of POV was registered for L-OW and H-OW than CTR sample, because lipid peroxides were cleaved to yield secondary oxidation products. The analysis of TBARS showed an increase during storage. Higher values for CTR sample than L-OW and H-OW were recorded at 0 and 3 and 7 days of storage.

For the colour parameters considered no difference during storage, neither among samples, were detected for redness (a^*), Hue angle and saturation index. The effect of time and diet was evident for all lipid and protein oxidation parameters but no for the color (Figures 1-3).

The results obtained for the ORAC_{FL} determinations in meat samples are reported in Figure 4. Differences were recorded for CTR versus L-OW and H-OW groups showing higher values in the antioxidant activity compared to control in all time of storage.

Discussions

The results of the present study show that supplementing chicken diet with

phenolic compounds through olive oil by-products protects meat against lipid and protein oxidation. Chemical deterioration, and in particular lipid oxidation, is one of the main factors able to limit the shelf-life of meat. The inclusion of natural antioxidants in animal diets has been reported by various authors as a way to slow down meat lipid oxidation during storage, therefore improving the quality and the conservation of products (Ranucci *et al.*, 2013; Luciano *et al.*, 2013).

The POV value increased and thereafter decreased with the time of storage, moreover the decomposition of hydroperoxides into secondary products appears to increase at a higher rate in the CTR compared with the treated samples. These data are in accordance with those of Hwang *et al.*




Figure 1. Effect of storage time on CIE a^* value of breast meat stored at 4°C.




Figure 2. Effect of storage time on Hue angle of breast meat muscle stored at 4°C.

(2013), who reported that POV values of cooked pork patties added with natural polyphenols, increased and thereafter decreased with the time of storage and highlighted that the decomposition of hydroperoxides into secondary products resulted delayed in treated samples.

Meat from animals fed dietary OW supplementation showed lower TBARS values during storage when compared to the control group. These results agree with other authors who found a reduction of lipid oxidation measured through TBARS during storage in lamb and beef muscles after olive oil by-products dietary supplementation (Luciano *et al.*, 2013; Branciari *et al.*, 2015a). Tufarelli *et al.* (2016) found an improved antioxidant defense system and a reduced TBARS level in chicken liver following dietary supplementation with extra virgin olive oil. Other authors have reported the beneficial effects of olive polyphenols on the oxidative status of meat measured through TBARS (Dal Bosco *et al.*, 2012; Branciari *et al.*, 2017), demonstrating their antioxidant effect in meat of animals fed olive phenolic compounds.

The extent of protein oxidation was evaluated by the formation of protein carbonyls. There are many reports on the biochemical changes in different meats during storage, most of which have focused on lipid oxidation (Abdel-Kader, 1996; Pikul *et al.*, 1984). Little work has been done on protein oxidation of meat obtained from animal fed natural antioxidant. Zhang *et al.* (2011) reported an increase in lipid and protein oxidation in the breast muscles of birds that had been fed an oxidized oil diet compared to antioxidant-supplemented and control diets. Jongberg *et al.* (2011) demonstrated the positive effect of grape extract on protein oxidation in chill stored beef patties. A significant decrease in carbonyl formation in meat added 3% (w/w) plant fruit extracts compared to control burger patties was observed in the study by Ganhão *et al.* (2010). Protein oxidation has been shown to induce a number of changes in proteins, such as modification of amino acid side chains, formation of protein polymers, loss of solubility, increase in carbonyl groups, change in amino acid composition and increase in proteolytic susceptibility (Levine *et al.*, 1990; Xiong, 2000). This modification of muscle proteins, because of denaturation and proteolysis, induces changes in meat quality including texture traits, colour, aroma, flavor and water-holding capacity.

Considered the foregoing it is clear that lipid and protein oxidations are closely associated with deteriorative processes able to affect the overall quality of meat and meat

products.

Meat colour was not affected by the phenolic compounds, contrary to what was observed in meat of other species (Branciari *et al.*, 2015a). This result might be due to the characteristics of *P. major* muscle which is mainly composed by α -white fibres that contain a lower amount of iron compared to the α -red or β -red type fibres and for this reason are less susceptible to colour modification (Branciari *et al.*, 2015b).

The results obtained from ORAC_{FL} assay showed the increased antioxidant activity of treated groups compared to control.

The effect of dietary OW on meat antioxidant activity might be related to

several factors, including the presence of bioactive molecules in tissues (muscle and liver) of animal fed with olive oil polyphenols as reported by Branciari *et al.* (2017). Indeed, there is evidence that polyphenols supplementation can exert a protective effect on α -tocopherols, a defensive barrier against lipid oxidation from oxidative decay (Bars-Cortina *et al.*, 2018). In addition, polyphenols were shown to affect the expression of genes involved in lambs' stress response and in primary homeostasis pathways, in both liver and muscle, increasing the expression of MYLK genes and improving the meat quality in female lambs (Sabino *et al.*, 2018).




Figure 3. Effect of storage time on Saturation index of breast meat stored at 4°C.




Figure 4. Effect of storage time on ORAC_{FL} of breast meat stored at 4°C.

Conclusions

The results of the present experiment demonstrated that olive polyphenols used in chicken diets could delay meat lipid and protein oxidation without affecting its colour stability. Further studies are needed in order to better clarify the role of olive mill wastewater bioactive compounds in the shelf-life of chicken meat, packaged with different systems. Moreover, additional data is required to define the most suitable concentration of polyphenol compounds to be used in animal diets able to exert at best their activity.

References

- Abdel-Kader ZM, 1996. Lipid oxidation in chicken as affected by cooking and frozen storage. *Mol Nut Food Res* 40:21-4.
- Bars-Cortina D, de las Hazas MCL, Benavent-Vallés A, Motilva MJ, 2018. Impact of dietary supplementation with olive and thyme phenols on alpha-tocopherol concentration in the muscle and liver of adult Wistar rats. *Food Funct* 9:1433-43.
- Branciari R, Ranucci D, Orteni R, Roila R, Trabalza-Marinucci M, Servili M, Papa P, Galarini R, Valiani A, 2016. Dietary administration of olive mill wastewater extract reduces campylobacter spp. Prevalence in broiler chickens. *Sustainability* 8:837.
- Branciari R; Ranucci D, Miraglia D, Urbani S, Esposto S, Servili M, 2015a. Effect of dietary treatment with olive oil by-product (olive cake) on physicochemical, sensory and microbial characteristics of beef during storage. *Ital J Food Saf* 4:225-9.
- Branciari R, Ranucci D, Trabalza-Marinucci M, Codini M, Orru M, Orteni R, Forte C, Ceccarini MR, Valiani A, 2015b. Evaluation of the antioxidant properties and oxidative stability of Pecorino cheese made from the raw milk of ewes fed *Rosmarinus officinalis* L. leaves. *Int J Food Sci Tech* 50:558-65.
- Branciari R, Galarini R, Giusepponi D, Trabalza-Marinucci M, Forte C, Roila R, Miraglia D, Servili M, Acuti G, Valiani A, 2017. Oxidative status and presence of bioactive compounds in meat from chickens fed polyphenols extracted from olive oil industry waste. *Sustainability* 9:1566.
- Dal Bosco A, Mourvaki E, Cardinali R, Servili M, Sebastiani B, Ruggeri S, Mattioli S, Taticchi A, Esposto S, Castellini C, 2012. Effect of dietary supplementation with olive pomaces on the performance and meat quality of growing rabbits. *Meat Sci* 92:783-8.
- Ganhão R, Morcuende D, Estévez M, 2010. Protein oxidation in emulsified cooked burger patties with added fruit extracts: Influence on colour and texture deterioration during chill storage. *Meat Sci* 85:402-9.
- Hwang KE, Choi YS, Choi SM, Kim HW, Choi JH, Lee MA, Kim CJ, 2013. Antioxidant action of ganghwayakssuk (*Artemisia princeps* Pamp.) in combination with ascorbic acid to increase the shelf life in raw and deep fried chicken nuggets. *Meat Science* 95: 593-602.
- Insani EM, Eyherabide A, Grigioni G, Sancho AM, Pensel NA, Descalzo AM, 2008. Oxidative stability and its relationship with natural antioxidants during refrigerated retail display of beef produced in Argentina. *Meat Sci* 79:444-52.
- Jongberg S, Skov SH, Tørrngren MA, Skibsted LH, Lund MN, 2011. Effect of white grape extract and modified atmosphere packaging on lipid and protein oxidation in chill stored beef patties. *Food Chem* 128:276-83.
- Levine RL, Garland D, Oliver CN, Amici A, Climent I, Lenz AG, Ahan BW, Shaltied S, Stadtman E R, 1990. Determination of carbonyl content in oxidatively modified proteins. *Method Enzymol* 186:464-78.
- Luciano G, Pauselli M, Servili M, Mourvaki E, Serra A, Monahan FJ, Lanza M, Priolo A, Zinna A, Mele M, 2013. Dietary olive cake reduces the oxidation of lipids, including cholesterol, in lamb meat enriched in polyunsaturated fatty acids. *Meat Sci* 93:703-14.
- Min Á, Nam KC, Cordray J, Ahn DU, 2008. Endogenous factors affecting oxidative stability of beef loin, pork loin, and chicken breast and thigh meats. *J Food Sci* 73:C439-46
- National Research Council, 1994. Nutrient requirements of poultry, Q19 9th rev. ed. Natl. Acad. Press, Washington, DC.
- Novelli E, Fasolato L, Cardazzo B, Carraro L, Taticchi A, Balzan S, 2014. Addition of phenols compounds to meat dough intended for salami manufacture and its antioxidant effect. *Ital J Food Saf* 3:154-6.
- Ou B, Hampsch-Woodill M, Prior RL, 2001. Development and validation of an improved oxygen radical absorbance capacity assay using fluorescein as the fluorescent probe. *J Agr Food Chem* 49:4619-26.
- Pikul J, Lesczynski DE, Bechtel PJ, Kummerow FA, 1984. Effects of frozen storage and cooking on lipid oxidation in chicken meat. *J Food Sci* 49:838-43.
- Ranucci D, Branciari R, Acuti G, Della Casa G, Trabalza-Marinucci M, Miraglia D, 2013. Quality traits of Ciauscolo salami from meat of pigs fed rosemary extract enriched diet. *Ital J Food Saf* 2:16.
- Reznick AZ & Packer L, 1994. Oxidative damage to proteins: Spectrophotometric method for carbonyl assay. *Method Enzymol* 233:357-63.
- Roila R, Branciari R, Ranucci D, Orteni R, Urbani S, Servili M, Valiani A, 2016. Antimicrobial activity of olive mill wastewater extract against *Pseudomonas fluorescens* isolated from mozzarella cheese. *Ital J Food Saf* 2:111.
- Sabino M, Carmelo VAO, Mazzoni G, Cappelli K, Capomaccio S, Ajmone-Marsan P, Verini-Supplizi A, Trabalza-Marinucci M, Kadarmideen HN 2018. Gene co-expression networks in liver and muscle transcriptome reveal sex-specific gene expression in lambs fed with a mix of essential oils. *BMC Genom* 19:236.
- Tarladgis BG, Watts BM, Yonathan M, 1960. Distillation method for the determination of malonaldehyde in rancid foods. *J Am Oil Chem Soc* 37:44-8.
- Tufarelli V, Laudadio V, Casalino E, 2016. An extra-virgin olive oil rich in polyphenolic compounds has antioxidant effects in meat-type broiler chickens. *Environ Sci Pollut Res Int* 23: 6197-204.
- Xiao S, Zhang WG, Lee EJ, Ahn DU, 2013. Effects of diet, packaging and irradiation on protein oxidation, lipid oxidation of raw broiler thigh meat. *Anim Industry Rep* 659:12.
- Xiong YL, 2000. Protein oxidation and implications for muscle food quality. *Antioxidants in Muscle Foods: Nutritional Strategies to Improve Quality*. Wiley and Sons, New York, NY, pp 85-90.
- Zhang W, Xiao S, Lee EJ, Ahn DU, 2011. Consumption of oxidized oil increases oxidative stress in broilers and affects the quality of breast meat. *J Agri Food Chem* 59:969-74.