



eISSN 2284-0230 - pISSN 1826-883

<https://www.pagepressjournals.org/index.php/jbr/index>

**Publisher's Disclaimer.** E-publishing ahead of print is increasingly important for the rapid dissemination of science. The **Early Access** service lets users access peer-reviewed articles well before print / regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

The **Journal of Biological Research** is, therefore, e-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear on a regular issue of the journal.

E-publishing of this PDF file has been approved by the authors.

J Biol Res 2024 [Online ahead of print]

*To cite this Article:*

Halimi E, Qirinxhi X, Paparisto A, Subashaj G. **Contribution to knowledge of the true bugs species (Hemiptera: Heteroptera) in the habitats of the Berati area in Albania.** *J Biol Res* doi: 10.4081/jbr.2024.12479

 ©The Author(s), 2024

Licensee [PAGEPress](#), Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

# **Contribution to knowledge of the true bugs species (Hemiptera: Heteroptera) in the habitats of the Berati area in Albania**

Eltjon Halimi,<sup>1</sup> Xhuliana Qirinxhi,<sup>2</sup> Anila Papparisto,<sup>1</sup> Gerta Subashaj<sup>3</sup>

<sup>1</sup>Faculty of Natural Sciences, University of Tirana, Tirana; <sup>2</sup>Faculty of Natural Sciences and Social Sciences, University “Fan.S. Noli” Korçë, Korçë; <sup>3</sup>Faculty of Medical Technical Sciences, Reald University College, Vlora, Albania

**Correspondence:** Eltjon Halimi, Faculty of Natural Sciences, University of Tirana, Address Tirana, Albania.

E-mail: [eltjon.halimi@fshn.edu.al](mailto:eltjon.halimi@fshn.edu.al)

**Key words:** heteroptera, taxonomy, ecosystems, ecology, Albania.

**Contributions:** EH, manuscript writing, material collection, taxonomic determination, ecological analysis; conclusion; XhQ, literature and references analysis, taxonomic analysis; AP, taxonomic and ecological analysis, manuscript review; recommendations; GS, systematic data analysis

**Conflict of interest:** the authors declare no conflict of interest.

**Availability of data and materials:** all data generated or analyzed during this study are included in this published article.

## **Abstract**

The present study provides taxonomic information pertaining to the distribution patterns of “true bugs” insects (Hemiptera: Heteroptera), throughout the ecological systems of the Berat region in Albania. Specimens were collected from May to September spanning the years 2018 to 2020.

The data were collected from stations that exhibited ideal habitats for this specific order, encompassing regions with ample flora, cultivated fields with crops, elevated hills, and hilly terrains. In the Berat region, our inquiry has successfully identified a total of 38 species belonging to the Heteroptera suborder (Hemiptera order). Furthermore, we have furnished citations for every Heteroptera species that have been identified inside the Berat region. Before our investigation, there was a total of 47 recorded species of “true bugs” in Berat. Significantly, our research uncovers a total of 24 species that have not been previously recognized in the Berat region. As a result, our research significantly expands the documented count of Heteroptera species in Berat to 71.

## **Introduction**

The Suborder Heteroptera (Hemiptera Order) exhibits a remarkable level of diversity, as seen by the current global list of around 42,300 species, and many more await description.<sup>1</sup> The recent Catalogue of the Heteroptera of the Palaearctic Region lists approximately 3000 species for Europe.<sup>2</sup> According to McGavin, it is the most extensive group of hemimetabolous insects.<sup>3</sup> The dimensions range from around 1mm to almost 11cm. The taxonomic identification of these organisms is facilitated by the systematically relevant traits of their external morphology and coloration.<sup>4</sup> The “true bugs” possess antennae consisting of four, less frequently three or five segments. True bugs live in virtually all terrestrial and aquatic ecosystems, including rainforest canopies, the open surface of the ocean, torrential and stagnant rivers, ephemeral rain pools and phytotelmata, large lakes, and aphotic caves and man-made buildings.<sup>5,6</sup> True bugs feed on many different resources (e.g., haemolymph of insects, phloem-, xylem-, or parenchym-sap of mosses, ferns, monocotyledons, mostly dicotyledons, the endosperm of seeds, plant pollen). The mouthparts, which evolved as sucking stylets for the uptake of liquid food and the injection of secretions from the salivary gland, are among their characteristic features; they commonly observe restricted diets.<sup>7</sup> Most species are phytophagous; some feed exclusively on particular plant species, genera, or families, whereas others are polyphagous, feeding on dozens to hundreds of different host plants. Many species are predatory, while others serve as biocontrol agents against agricultural pests.<sup>8</sup> The taxonomic characterization of defensive secretory glands

located in the metathorax holds significant importance.<sup>9</sup> According to Derjanschi and Pericart, the front part of mesothoracic wings is reinforced and separated into multiple sections.<sup>10</sup> Taxonomic determination is facilitated by male genitalia, wherein the ninth segment is frequently employed due to its distinct location throughout several families.<sup>11</sup> In the majority of instances, it is feasible to differentiate between females and males due to the presence of a vertical fissure in certain abdomen segments.<sup>10</sup>

In this study, we present the true bugs species (Heteroptera) found in the area of Berat, analyzing all references pertaining to these species within our study area. The relief of this area encompasses plains intertwined with hills, southern mountain ranges, and valleys, while the climate exhibits characteristics of a hill-mountain climate, particularly evident in the variation of climatic elements with altitude. Winter temperatures range from 7°C to 3.2°C, while in summer, they fluctuate between 23.5°C and 21°C. The predominant land cover consists of brown and dark brown forested areas.

## **Materials and Methods**

Heteropteran insects were collected between 2018 and 2020 during the warm period from May to September. This period corresponds with the development of hemipteran species after overwintering and with their main flight activity.<sup>12</sup> The collection of heteropterans was carried out in the district of Berat, at stations characterized by favorable habitats for these insects, including fields, hills, and vegetation of agricultural crops.<sup>13</sup> We have collected data at Poshnja station, located at coordinates 40° 46' 49" N and 19° 50' 39" E, with altitudes ranging from 20 to 30 meters above sea level; Lapardha station, located at coordinates 40° 23' 12" N and 19° 33' 29" E, with altitudes ranging from 120 to 180 meters above sea level; Uznova station, located at coordinates 40° 41' 50" N and 19° 58' 49"E, with altitudes ranging from 50 to 85 meters above sea level; Vodica station, located at coordinates 40° 40' 36"N and 20° 0' 31"E, with altitudes ranging from 30 to 60 meters above sea level; Peshtan station, located at coordinates 40° 14' 5"N and 20° 32' 11"E, has altitudes ranging from 300 to 450 meters above sea level and Lybesh station, located at coordinates 40° 39' 49"N and 20° 6' 57"E, has altitudes ranging from 700 to 8,000 meters above sea level. Surveys were carried out by visiting each station three times, each lasting 4-5 days, under sunny weather conditions from 10<sup>00</sup> to 14<sup>00</sup> during the daytime, and samplings were performed randomly in the stations habitats. An entomological net was used for

heteropterans, considering the life cycle of heteropterans.<sup>14</sup> Plastic bottles, labeled with the date and collection site, were employed for insect storage. The insects were stored in bottles containing a solution of 70% ethanol, acetic acid, and distilled water (in 80:5:20 ratio), along with a few drops of ether.<sup>14</sup> Genitalia were prepared for observation by maceration in 10% potassium-hydroxide (KOH).<sup>15</sup> The identification of specimens was facilitated using ZEISS stereomicroscope (Uzi Seria 240/3 Model 50240003), keys specific to heteropterans,<sup>16-21</sup> as well as reference to previous collections housed at the Museum of Natural Science in Tirana. All publications with references to true bugs insects (Hemiptera: Heteroptera) in the areas of Albania were analyzed. From these references, only those publications presenting references to Heteroptera species specifically for our study area, Berat in Albania, were considered valid for our study. Based on these references, as well as the data from our previous publication on this order,<sup>10</sup> in this region, we further developed our analysis and systematic data for this order of insects.

## **Results**

We report 38 species of true bugs insects (Hemiptera: Heteroptera) for the Berat region, along with the collection station, expedition date, and whether it is referred only by us or by other reference sources. We have grouped the species according to the taxonomic classification of families and the Infraorder of these families (Supplementary Materials, Table 1).

The analysis of references related to "true bugs" (Hemiptera: Heteroptera) in Albania has been conducted.<sup>22-32</sup> In each of these publications, we have analyzed the reference stations for these insects to determine the publications that refer to these insects from our study area. Only publications that report "true bugs" (Hemiptera: Heteroptera) species for our study region, have been selected from these places. Out of all the prior papers on Albania, only one publication specifically mentions the presence of "true bugs" (Hemiptera: Heteroptera) species in our research area in Berat.<sup>26</sup> The reference has been included in the aforementioned list. In this study, we have identified 14 species that have been previously referred by Josifov,<sup>26</sup> while we have identified for the first time 24 species that were not noticed previously for their presence in this region.

## **Discussion**

The analysis conducted in this study is based on the publication authored by Josifov,<sup>26</sup> which serves as an extensive and authoritative resource for studying “true bugs” (Hemiptera: Heteroptera) species in the Berat region. This publication documents 47 Heteroptera species in the Berat area (Supplementary Materials). We observed that out of the 38 species we encountered, 14 were previously documented by Josifov,<sup>26</sup> while 24 species represent new records for this area. Out of the 38 species encountered, the family Pentatomidae, with 12 species, and the infraorder Pentatomorpha, with 28 species, exhibit the highest biodiversity; based on this, we can conclude that the habitats in the Berat area provide the best ecological conditions for these species. The habitats of the considered stations in the Berat area consisted of either agricultural farms or covered in crops, weeds, or grasses fields. There are fields planted with plantations of apples, peaches, berries, peppers, beans, and pecans, or fields planted with sorghum and cotton, which are considered field crops. After incorporating species references from our research, the cumulative number of “true bugs” (Hemiptera: Heteroptera) species recorded in the Berat region of Albania has reached 71 species.

Meanwhile, 34 Heteroptera species referenced by Josifov<sup>26</sup> for the Berat area were not encountered during our expeditions to stations in this region. We attribute this difference in species composition to several factors, including the temporal gap between the expeditions conducted by the German Entomological Institute in the 1960s and our study. Additionally, socio-economic changes in the area have likely played a role. The decline in rural populations and shifts in land use patterns, particularly following the collapse of communism in Albania in the 1990s, may have resulted in habitat degradation and loss, impacting the distribution of Heteroptera species.<sup>28</sup> Our study increases the knowledge about species diversity of “true bugs” (Hemiptera: Heteroptera) in the Berat area, giving valuable information not only for this region but also for Albania as a whole. Further research is needed to assess the impact of ongoing environmental changes on insect populations and to update our knowledge of species distributions in the region.

## **Conclusions**

Our investigation in the Berat area has revealed the presence of 38 species of “true bugs” (Hemiptera: Heteroptera). Our study contributes to the knowledge of 24 “true bugs” species for the first time in the Berat area. Consequently, following our study the total count of “true bugs”

(Hemiptera: Heteroptera) species for this region increases to 71. However, it is worth noting that we did not encounter 34 species listed in the literature references. This disparity underscores the need for further research to comprehensively document the “true bugs” (Hemiptera: Heteroptera) fauna in the Berat area.

Based on our findings, we strongly advocate for the implementation of measures and action plans aimed at preserving the natural habitats in the Berat area. This is crucial to prevent further loss of biodiversity among Heteroptera species. Additionally, we emphasize the importance of exercising caution in new construction projects in the region, as they pose a significant threat to habitat integrity and biodiversity. In conclusion, our study highlights the importance of proactive conservation efforts to safeguard the “true bugs” (Hemiptera: Heteroptera) diversity in the Berat area. By prioritizing habitat protection and minimizing anthropogenic impacts, we can work towards ensuring the long-term survival of “true bugs” (Hemiptera: Heteroptera) species in this ecologically significant region.

## References

1. Schuh RT, Weirauch C. True bugs of the world (Hemiptera: Heteroptera) classification and natural history. 2<sup>nd</sup> ed. (Mongraph Series Volume 8). London: Siri Scientific Press; 2020.
2. Aukema B, Rieger Chr. Catalogue of the Heteroptera of the palaeartic region. Vol. 5. Pentatomomorpha II. Amsterdam: The Netherlands Entomological Society; 2006.
3. McGavin CG. Bugs of the world. London: Blandford Pr; 1999.
4. Ribes J. Un nouvel emesinae troglophile indo-malais (Heteroptera, Reduviidae). [A new indo-malayan thread-legged bug troglophile (Heteroptera, Reduviidae)]. Rev Suisse Zool 1987;94:251-6.
5. Henry TJ. Biodiversity of Heteroptera. Insect biodiversity: science and society. Chichester: Blackwell Publishing Ltd; 2009. pp: 23–63.
6. Schuh RT, Slater J. True bugs of the world (Hemiptera: Heteroptera) classification and natural history. New York, NY: Cornell University Press; 1995.

7. Rabitsch W. True bugs (Hemiptera, Heteroptera) chapter 9.1. Alien terrestrial arthropods of Europe. *BioRisk* 2010;4:407–33.
8. Schaefer CW, Panizzi AR. Heteroptera of economic importance. Florida: CRC Press; 2000.
9. Caprile JM. Systematic catalogue of the Reduviidae of the world (Insecta: Heteroptera). Mayagüez: University of Puerto Rico; 1990.
10. Derjanschi V, Pericart J. Hemipteres Pentatomoidea euro-mediterraneens. [Euro-Mediterranean Hemiptera: Pentatomoidea]. Vol 1 Faune de France 90. Paris: Fédération Française des Sociétés de Sciences Naturelles; 2005.
11. Grazia J, Panizzi A.R, Greve C, et al. True bugs (Heteroptera) of the neotropics. Berlin: Springer; 2015.
12. Miller NCE. The biology of the Heteroptera 2<sup>nd</sup> ed. London: E.W.Classey Ltd; 1971. pp 116-32.
13. Halimi E, Papparisto A, Alameti E. Taxonomic and ecological data on the Hemiptera fauna from Berati region habitats in Albania. *J Biol Res* 2023;96:11105.
14. Colas G. Guide del'entomologiste. [Entomologist's guide]. Paris: Editions N. Boubée; 2000.
15. Tsagkarakis A, Thanou Z, Chaldeou A, et al. New records and updated checklist of the Pentatomoidea (Hemiptera: Heteroptera) of Greece. *Insects* 2022;13:749.
16. Forero D. The systematics of the Hemiptera. *Rev Colomb Entomol* 2008;34:1–21.
17. Chinery M. Insects identify bugs, beetles, bees, and much more. London: Collins Publishers; 2012.
18. Aukema B, Rieger Ch. Catalogue of the Heteroptera of the palearctic region. Vol. 4. Pentatomomorpha I. Amsterdam: The Netherlands Entomological Society; 2001.
19. Aukema B, Rieger Ch, Rabitsch W. Catalogue of the Heteroptera of the palaeartic region. Volume 6. Supplement. Amsterdam: The Netherlands Entomological Society; 2013.



20. Péricart J. Hémiptères Pentatomoidea Euro-Méditerranéens; Faune de France. [Euro-Mediterranean Hemiptera Pentatomoidea; Fauna of France]. Paris: Fédération Française des Sociétés de Sciences Naturelles; 2010.
21. Ramsay AJ. Pentatomoidea (Hemiptera: Heteroptera) of Greece - An annotated checklist. *Monogr Up Sil Mus* 2019;10:9–27.
22. Misja K. Të dhëna paraprake mbi inventarizimin e Hemipterëve (Hemiptera). [Preliminary data on the inventory of hemiptera (Hemiptera)]. *Bul. Shkenc. Nat. U.SH.T* 1979;4:7-16.
23. Ciski E. Csiki Erno állattani kutatásai Albániában. - explorationes zoologicae ab E. Csiki in Albania peractae. Viii. Levéldarazsak-Tenthredinoidea. [Erno Csiki's zoological research in Albania. - explorationes zoologicae ab E. Csiki in Albania completed. Viii Sawflies – Tenthredinoidea] *J MTA Kvt Kézirtár Kat* 1923;1:103-8.
24. Horváth G. Albania Hemiptera - faunaja. [Hemiptera fauna of Albania]. *Ann Hist Nat Mus Hung* 1916;14:1-16.
25. Frisse G. Ergebnisse der Albanian-expedition 1961 des deutschen entomogischen institutes. Verzeichnis albanischer fundort. [Results of the 1961 Albania expedition of the German entomological institute. Directory of Albanian localities]. *Beitr Entomol Berlin* 1967;3/4:405-34.
26. Josifov M. Ergebnisse der Albanien-Expedition 1961 des Deutschen Entomologischen Institutes. [Results of the 1961 Albania expedition of the German Entomological Institute]. *Beit Entomol Berl* 1970;7/8:825-956.
27. Schumacher F. Hemipteren aus Albanien und Epirus. [Hemiptera from Albania and Epirus]. *Sber Ges naturf Freunde Berl* 1914;15:116-27.
28. McKinney ML. Urbanization, biodiversity, and conservation. *BioScience* 2002;52:883-90.
29. Štys P, Kerzhner I. The rank and nomenclature of higher taxa in recent Heteroptera. *Acta Entomol Bohemoslov* 1975;72:65-79.

30. Henry T.J. 1997. Phylogenetic analysis of family groups within the infraorder Pentatomomorpha (Hemiptera: Heteroptera), with emphasis on the Lygaeoidea. *Ann Entomol Soc Am* 1997;90:275–301.
31. Faúndez E. Los Coreoidea Leach, 1815 (Hemiptera: Heteroptera) de la región de Magallanes: Lista de especies y clave de identificación. [The Coreoidea Leach, 1815 (Hemiptera: Heteroptera) of Magallanes region: checklist and identification key to the species]. *An Inst Patagon* 2016;944:39-42.
32. Burdfield-Steel ER, Shuker DM. The evolutionary ecology of the Lygaeidae. *Ecol Evol* 2014;4:278-301.
33. Cassis G, Vanags L. Jewel bugs of Australia (Insecta, Heteroptera, Scutelleridae). In: W. Rabitsch (Eds.). *Denisia* 2006;19:275-398.
34. Fowles TM, Coscarón MC, Panizzi AR, Carroll SP. Scentless plant bugs (Rhopalidae). In: true bugs (Heteroptera) of the Neotropics. *Entomology in Focus*. Dordrecht: Springer Nature. 2015;607-37.
35. Henry TJ, Adamski D. 1998 *Rhyparochromus saturnius* (Rossi) (Heteroptera: Lygaeoidea: Rhyparochromidae), a palearctic seed bug newly discovered in North America. *J.N.Y. Entomol Soc* 1998;106:132-40.
36. Schaefer WC. The higher classification of the Alydidae (Hemiptera:Heteroptera). *Proceedings of the WA. Entomol Soc* 1999;101:94–8.
37. Kuznetsova V, Grozeva S, Nokkala S, Nokkala C. Cytogenetics of the true bug infraorder Cimicomorpha (Hemiptera, Heteroptera): a review. *ZooKeys* 2011;154:31–70.
38. Henry TJ, Wheeler AG. Family Miridae Hahn, 1833 (= Capsidae Burmeister, 1835). The plant bugs. 1<sup>st</sup> ed. Florida: CRC Press;1988. pp 251-507.
39. Weirauch C, Bérenger JM, Berniker L. et al. An illustrated identification key to assassin bug subfamilies and tribes (Hemiptera: Reduviidae). *Can J Arthropod Identif* 2014;26.

## **Supplementary Materials**

**Annex 1.** List of species referred by Josifov.<sup>26</sup>

**Table 1.** Heteroptera species collected in the Berat area.

Submitted: 15 March 2024

Accepted: 13 July 2024

Early access: 9 September 2024